

Space-time error estimates for deep neural network approximations for differential equations

Philipp Grohs¹, Fabian Hornung^{2,3},
Arnulf Jentzen⁴, and Philipp Zimmermann^{5,6}

¹Faculty of Mathematics and Research Platform Data Science, University of Vienna

Austria, e-mail: philipp.grohs@univie.ac.at

² Department of Mathematics, ETH Zurich, Zürich,

Switzerland, e-mail: fabianhornung89@gmail.com

³ Faculty of Mathematics, Karlsruhe Institute of Technology, Karlsruhe,

Germany, e-mail: fabianhornung89@gmail.com

⁴ Department of Mathematics, ETH Zurich, Zürich,

Switzerland, e-mail: arnulf.jentzen@sam.math.ethz.ch

⁵ Department of Mathematics, ETH Zurich, Zürich,

Switzerland, e-mail: philipp.zimmermann@math.ethz.ch

⁶ Faculty of Mathematics, University of Regensburg, Regensburg,

Germany, e-mail: philipp.zimmermann@ur.de

August 13, 2019

Abstract

Over the last few years deep artificial neural networks (DNNs) have very successfully been used in numerical simulations for a wide variety of computational problems including computer vision, image classification, speech recognition, natural language processing, as well as computational advertisement. In addition, it has recently been proposed to approximate solutions of partial differential equations (PDEs) by means of stochastic learning problems involving DNNs. There are now also a few rigorous mathematical results in the scientific literature which provide error estimates for such deep learning based approximation methods for PDEs. All of these articles provide spatial error estimates for neural network approximations for PDEs but do not provide error estimates for the entire space-time error for the

considered neural network approximations. It is the subject of the main result of this article to provide space-time error estimates for DNN approximations of Euler approximations of certain perturbed differential equations. Our proof of this result is based (i) on a certain artificial neural network (ANN) calculus and (ii) on ANN approximation results for products of the form $[0, T] \times \mathbb{R}^d \ni (t, x) \mapsto tx \in \mathbb{R}^d$ where $T \in (0, \infty)$, $d \in \mathbb{N}$, which we both develop within this article.

Contents

1	Introduction	3
2	Artificial neural network (ANN) calculus	5
2.1	Artificial neural networks (ANNs) and their realizations	5
2.2	Compositions of ANNs	7
2.2.1	Standard compositions of ANNs	7
2.2.2	Associativity of standard compositions of ANNs	11
2.2.3	Compositions of ANNs and affine linear transformations	14
2.2.4	Powers and extensions of ANNs	15
2.2.5	Compositions of ANNs involving artificial identities	19
2.3	Parallelizations of ANNs	21
2.3.1	Parallelizations of ANNs with the same length	21
2.3.2	Parallelizations of ANNs with different lengths	25
2.4	Sums of ANNs	28
2.4.1	Sums of ANNs with the same length	28
2.4.2	Sums of ANNs with different lengths	30
2.5	ANN representations for Euler approximations	32
2.5.1	ANN representations for one Euler step	32
2.5.2	ANN representations for multiple nested Euler steps	37
2.5.3	ANN representations for multiple perturbed nested Euler steps	42
3	ANN approximation results	45
3.1	ANN approximations for the square function	45
3.1.1	Explicit approximations for the square function on $[0, 1]$	45
3.1.2	ANN approximations for the square function on $[0, 1]$	50
3.1.3	ANN approximations for the square function on \mathbb{R}	54
3.2	ANN approximations for products	57
3.2.1	ANN approximations for one-dimensional products	57
3.2.2	ANN approximations for multi-dimensional products	62
3.3	Space-time ANN approximations for Euler approximations	65
3.3.1	Space-time representations for Euler approximations	65

3.3.2	ANN representations for hat functions	67
3.3.3	A posteriori error estimates for ANN approximations	68
3.3.4	A priori estimates for Euler approximations	75
3.3.5	A priori error estimates for ANN approximations	77

1 Introduction

Over the last few years deep artificial neural networks (DNNs) have very successfully been used in numerical simulations for a wide variety of computational problems including computer vision, image classification, speech recognition, natural language processing, as well as computational advertisement (cf., e.g., the references mentioned in [14, 17, 25]). In addition, the articles [9, 18] suggest to approximate solutions of partial differential equations (PDEs) by means of stochastic learning problems involving DNNs. We also refer to [1, 2, 3, 4, 5, 6, 8, 10, 12, 13, 15, 19, 20, 21, 24, 28, 29, 30, 33, 34, 36] for extensions and improvements of such deep learning based approximation methods for PDEs. There are now also a few rigorous mathematical results in the scientific literature which provide error estimates for such deep learning based approximation methods for PDEs; see, e.g., [7, 11, 16, 19, 22, 25, 26, 35, 36]. The articles in this reference list all provide spatial error estimates for neural network approximations for PDEs but do not provide error estimates for the entire space-time error for the considered neural network approximations. It is the subject of Theorem 3.12 in this article, which is the main result of this article, to provide space-time error estimates for DNN approximations of Euler approximations of certain perturbed differential equations. To illustrate the findings of the main result of this article in more details, we now formulate in Theorem 1.1 below a special case of Theorem 3.12.

Theorem 1.1. *Let $\mathfrak{C}, T, \mathfrak{d} \in (0, \infty)$, let $A_d \in C(\mathbb{R}^d, \mathbb{R}^d)$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ that $A_d(x) = (\max\{x_1, 0\}, \max\{x_2, 0\}, \dots, \max\{x_d, 0\})$, let $\mathbf{N} = \cup_{L \in \mathbb{N}} \cup_{(l_0, l_1, \dots, l_L) \in \mathbb{N}^{L+1}} (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$, let $R: \mathbf{N} \rightarrow \cup_{k, l \in \mathbb{N}} C(\mathbb{R}^k, \mathbb{R}^l)$ and $P: \mathbf{N} \rightarrow \mathbb{N}$ satisfy for all $L \in \mathbb{N}$, $l_0, l_1, \dots, l_L \in \mathbb{N}$, $\Phi \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$, $\Psi = ((W_1, B_1), (W_2, B_2), \dots, (W_L, B_L)) \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$, $x_0 \in \mathbb{R}^{l_0}$, $x_1 \in \mathbb{R}^{l_1}, \dots, x_{L-1} \in \mathbb{R}^{l_{L-1}}$ with $\forall k \in \mathbb{N} \cap (0, L): x_k = A_{l_k}(W_k x_{k-1} + B_k)$ that $P(\Phi) = \sum_{k=1}^L l_k(l_{k-1}+1)$, $R(\Psi) \in C(\mathbb{R}^{l_0}, \mathbb{R}^{l_L})$, and $(R(\Psi))(x_0) = W_L x_{L-1} + B_L$, let $\Phi_d \in \mathbf{N}$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$ that $R(\Phi_d) \in C(\mathbb{R}^d, \mathbb{R}^d)$, $\|(R(\Phi_d))(x)\| \leq \mathfrak{C}(1 + \|x\|)$, and $P(\Phi_d) \leq \mathfrak{C}d^{\mathfrak{d}}$, let $Y^{d,N} = (Y_{t,x}^{d,N})_{(t,x) \in [0,T] \times \mathbb{R}^d}: [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$, $N, d \in \mathbb{N}$, be the functions which satisfy for all $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$ that $Y_{0,x}^{d,N} = x$ and*

$$Y_{t,x}^{d,N} = Y_{\frac{nT}{N}, x}^{d,N} + (t - \frac{nT}{N}) (R(\Phi_d))(Y_{\frac{nT}{N}, x}^{d,N}). \quad (1)$$

Then there exist $C \in \mathbb{R}$ and $\Psi_{\varepsilon,d,N} \in \mathbf{N}$, $N, d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, such that

(i) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$ that $R(\Psi_{\varepsilon,d,N}) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(ii) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\|Y_{t,x}^{d,N} - (R(\Psi_{\varepsilon,d,N}))(t, x)\| \leq Cd^{1/2}N^{3/2}\varepsilon(1 + \|x\|^3), \quad (2)$$

(iii) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\|(R(\Psi_{\varepsilon,d,N}))(t, x)\| \leq Cd^{1/2}N(1 + \|x\|^2), \quad (3)$$

and

(iv) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$ that

$$P(\Psi_{\varepsilon,d,N}) \leq Cd^{16+8d}N^6[1 + |\ln(\varepsilon)|^2]. \quad (4)$$

Theorem 1.1 is an immediate consequence of Corollary 3.13 in Subsection 3.3.5 below. Corollary 3.13, in turn, follows from Theorem 3.12 in Subsection 3.3.5, which is the main result of this article. Our proof of Theorem 1.1 and Theorem 3.12, respectively, is based on a certain artificial neural network (ANN) calculus, which we develop in Section 2. Section 2 is in parts based on several well-known concepts and results in the scientific literature (cf., e.g., [11, 25, 32, 37]). We refer to the beginning of Section 2 for a more detailed comparison of the content of Section 2 with the material in related articles in the scientific literature. Our proof of Theorem 1.1 and Theorem 3.12, respectively, is mainly inspired by [25], [11, Section 6], and [37, Section 3.2]. Theorem 1.1 and Theorem 3.12, respectively, provide error estimates for rectified DNN approximations of Euler approximations of certain perturbed differential equations. Many of the DNN approximation and representation results of this work, however, apply to DNNs with more general activation functions than only the rectifier function (cf., e.g., Li et al. [27, Section 1] and Petersen et al. [31, Section 2] for further activation functions). The error estimates for rectified DNN approximations of Euler approximations of perturbed differential equations, which we establish in Theorem 1.1 and Theorem 3.12, respectively, can then be used to establish space-time error estimates for DNN approximations for PDEs. This will be the subject of a future research article, which will be based on this article.

The remainder of this article is organized as follows. In Section 2 we develop the above mentioned ANN calculus and, in particular, we establish in Subsection 2.5 ANN representation results for Euler approximations. In Subsection 3.1 we develop ANN approximation results for the square function $\mathbb{R} \ni x \mapsto x^2 \in \mathbb{R}$. These ANN

approximation results for the square function are then used in Subsection 3.2 to develop ANN approximation results for products of the form $[0, T] \times \mathbb{R}^d \ni (t, x) \mapsto tx \in \mathbb{R}^d$ where $T \in (0, \infty)$, $d \in \mathbb{N}$. In Subsection 3.3 we then combine the ANN representation results in Subsection 2.5 with the ANN approximation results for products in Subsection 3.2 to establish in Theorem 3.12 the main result of this article.

2 Artificial neural network (ANN) calculus

This section develops a certain calculus for ANNs. Some of the notions and results which we present here are rather elementary, but for convenience of the reader we present here all details and we include the proof of every result. The material in this section is also in parts based on several well-known concepts and results in the scientific literature. In particular, Definition 2.1, Definition 2.2, and Definition 2.3 are slight reformulations of Petersen & Voigtlaender [32, Definition 2.1]. Moreover, Lemma 2.4 is elementary and well-known in the scientific literature. Furthermore, Definition 2.5 is also a slight reformulation of Petersen & Voigtlaender [32, Definition 2.2]. In addition, Proposition 2.6, Corollary 2.7, and Lemma 2.8 are elementary and essentially well-known in the scientific literature (cf., e.g., Petersen & Voigtlaender [32]). Moreover, Definition 2.11 is an extension of Elbrächter et al. [11, Setting 5.2] and Proposition 2.16 is in parts an extension of Elbrächter et al. [11, Lemma 5.3]. Furthermore, Definition 2.17 and Definition 2.22 extend Elbrächter et al. [11, Setting 5.2] (cf., e.g., Petersen & Voigtlaender [32, Definition 2.7]). In addition, Proposition 2.25 is a reformulation of [25, Lemma 5.1]. Moreover, Lemma 2.27 and Proposition 2.28 are significantly inspired by [25, Proposition 5.3]. Furthermore, item (iv) in Lemma 2.27 and item (iv) in Proposition 2.28, respectively, improve the parameter estimates in [25, Proposition 5.3]. In addition, Corollary 2.31 in Subsection 2.5.2 below is also in parts inspired by [25, Proposition 6.1].

2.1 Artificial neural networks (ANNs) and their realizations

Definition 2.1 (Artificial neural networks (ANNs)). *We denote by \mathbf{N} the set given by*

$$\mathbf{N} = \cup_{L \in \mathbb{N}} \cup_{(l_0, l_1, \dots, l_L) \in \mathbb{N}^{L+1}} \left(\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}) \right) \quad (5)$$

and we denote by $\mathcal{P}, \mathcal{L}, \mathcal{I}, \mathcal{O}: \mathbf{N} \rightarrow \mathbb{N}$, $\mathcal{H}: \mathbf{N} \rightarrow \mathbb{N}_0$, and $\mathcal{D}: \mathbf{N} \rightarrow \cup_{L=2}^{\infty} \mathbb{N}^L$ the functions which satisfy for all $L \in \mathbb{N}$, $l_0, l_1, \dots, l_L \in \mathbb{N}$, $\Phi \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$

that $\mathcal{P}(\Phi) = \sum_{k=1}^L l_k(l_{k-1} + 1)$, $\mathcal{L}(\Phi) = L$, $\mathcal{I}(\Phi) = l_0$, $\mathcal{O}(\Phi) = l_L$, $\mathcal{H}(\Phi) = L - 1$, and $\mathcal{D}(\Phi) = (l_0, l_1, \dots, l_L)$.

Definition 2.2 (Multidimensional versions). *Let $d \in \mathbb{N}$ and let $\psi: \mathbb{R} \rightarrow \mathbb{R}$ be a function. Then we denote by $\mathfrak{M}_{\psi,d}: \mathbb{R}^d \rightarrow \mathbb{R}^d$ the function which satisfies for all $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ that*

$$\mathfrak{M}_{\psi,d}(x) = (\psi(x_1), \dots, \psi(x_d)). \quad (6)$$

Definition 2.3 (Realizations associated to ANNs). *Let $a \in C(\mathbb{R}, \mathbb{R})$. Then we denote by $\mathcal{R}_a: \mathbf{N} \rightarrow \cup_{k,l \in \mathbb{N}} C(\mathbb{R}^k, \mathbb{R}^l)$ the function which satisfies for all $L \in \mathbb{N}$, $l_0, l_1, \dots, l_L \in \mathbb{N}$, $\Phi = ((W_1, B_1), (W_2, B_2), \dots, (W_L, B_L)) \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$, $x_0 \in \mathbb{R}^{l_0}, x_1 \in \mathbb{R}^{l_1}, \dots, x_{L-1} \in \mathbb{R}^{l_{L-1}}$ with $\forall k \in \mathbb{N} \cap (0, L): x_k = \mathfrak{M}_{a,l_k}(W_k x_{k-1} + B_k)$ that*

$$\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{l_0}, \mathbb{R}^{l_L}) \quad \text{and} \quad (\mathcal{R}_a(\Phi))(x_0) = W_L x_{L-1} + B_L \quad (7)$$

(cf. Definition 2.2 and Definition 2.1).

Lemma 2.4. *Let $\Phi \in \mathbf{N}$ (cf. Definition 2.1). Then*

(i) *it holds that $\mathcal{D}(\Phi) \in \mathbb{N}^{\mathcal{L}(\Phi)+1}$ and*

(ii) *it holds for all $a \in C(\mathbb{R}, \mathbb{R})$ that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{O}(\Phi)})$*

(cf. Definition 2.3).

Proof of Lemma 2.4. Note that the assumption that $\Phi \in \mathbf{N} = \cup_{L \in \mathbb{N} \cup (l_0, l_1, \dots, l_L) \in \mathbb{N}^{L+1}} (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$ ensures that there exist $L \in \mathbb{N}$, $l_0, l_1, \dots, l_L \in \mathbb{N}$ such that

$$\Phi \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k})). \quad (8)$$

Observe that (8) assures that

$$\mathcal{L}(\Phi) = L, \quad \mathcal{I}(\Phi) = l_0, \quad \mathcal{O}(\Phi) = l_L, \quad (9)$$

$$\text{and} \quad \mathcal{D}(\Phi) = (l_0, l_1, \dots, l_L) \in \mathbb{N}^{L+1} = \mathbb{N}^{\mathcal{L}(\Phi)+1}. \quad (10)$$

This establishes item (i). Moreover, note that (9) and (7) show that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{O}(\Phi)})$. This establishes item (ii). The proof of Lemma 2.4 is thus completed. \square

2.2 Compositions of ANNs

2.2.1 Standard compositions of ANNs

Definition 2.5 (Standard compositions of ANNs). *We denote by $(\cdot) \bullet (\cdot) : \{(\Phi_1, \Phi_2) \in \mathbf{N} \times \mathbf{N} : \mathcal{I}(\Phi_1) = \mathcal{O}(\Phi_2)\} \rightarrow \mathbf{N}$ the function which satisfies for all $L, \mathcal{L} \in \mathbb{N}$, $l_0, l_1, \dots, l_L, l_0, l_1, \dots, l_{\mathcal{L}} \in \mathbb{N}$, $\Phi_1 = ((W_1, B_1), (W_2, B_2), \dots, (W_L, B_L)) \in (\times_{k=1}^L \mathbb{N}^{\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}})$, $\Phi_2 = ((\mathcal{W}_1, \mathfrak{B}_1), (\mathcal{W}_2, \mathfrak{B}_2), \dots, (\mathcal{W}_{\mathcal{L}}, \mathfrak{B}_{\mathcal{L}})) \in (\times_{k=1}^{\mathcal{L}} (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$ with $l_0 = \mathcal{I}(\Phi_1) = \mathcal{O}(\Phi_2) = l_{\mathcal{L}}$ that*

$$\Phi_1 \bullet \Phi_2 = \begin{cases} ((\mathcal{W}_1, \mathfrak{B}_1), (\mathcal{W}_2, \mathfrak{B}_2), \dots, (\mathcal{W}_{\mathcal{L}-1}, \mathfrak{B}_{\mathcal{L}-1}), (W_1 \mathcal{W}_{\mathcal{L}}, W_1 \mathfrak{B}_{\mathcal{L}} + B_1), \\ \quad (W_2, B_2), (W_3, B_3), \dots, (W_L, B_L)) & : L > 1 < \mathcal{L} \\ ((W_1 \mathcal{W}_1, W_1 \mathfrak{B}_1 + B_1), (W_2, B_2), (W_3, B_3), \dots, (W_L, B_L)) & : L > 1 = \mathcal{L} \\ ((\mathcal{W}_1, \mathfrak{B}_1), (\mathcal{W}_2, \mathfrak{B}_2), \dots, (\mathcal{W}_{\mathcal{L}-1}, \mathfrak{B}_{\mathcal{L}-1}), (W_1 \mathcal{W}_{\mathcal{L}}, W_1 \mathfrak{B}_{\mathcal{L}} + B_1)) & : L = 1 < \mathcal{L} \\ (W_1 \mathcal{W}_1, W_1 \mathfrak{B}_1 + B_1) & : L = 1 = \mathcal{L} \end{cases} \quad (11)$$

(cf. Definition 2.1).

Proposition 2.6. *Let $\Phi_1, \Phi_2 \in \mathbf{N}$, $l_{1,0}, l_{1,1}, \dots, l_{1,\mathcal{L}(\Phi_1)}, l_{2,0}, l_{2,1}, \dots, l_{2,\mathcal{L}(\Phi_2)} \in \mathbb{N}$ satisfy for all $k \in \{1, 2\}$ that $\mathcal{I}(\Phi_1) = \mathcal{O}(\Phi_2)$ and $\mathcal{D}(\Phi_k) = (l_{k,0}, l_{k,1}, \dots, l_{k,\mathcal{L}(\Phi_k)})$ (cf. Definition 2.1). Then*

(i) *it holds that*

$$\mathcal{D}(\Phi_1 \bullet \Phi_2) = (l_{2,0}, l_{2,1}, \dots, l_{2,\mathcal{L}(\Phi_2)-1}, l_{1,1}, l_{1,2}, \dots, l_{1,\mathcal{L}(\Phi_1)}), \quad (12)$$

(ii) *it holds that*

$$[\mathcal{L}(\Phi_1 \bullet \Phi_2) - 1] = [\mathcal{L}(\Phi_1) - 1] + [\mathcal{L}(\Phi_2) - 1], \quad (13)$$

(iii) *it holds that*

$$\mathcal{H}(\Phi_1 \bullet \Phi_2) = \mathcal{H}(\Phi_1) + \mathcal{H}(\Phi_2), \quad (14)$$

(iv) *it holds that*

$$\begin{aligned} \mathcal{P}(\Phi_1 \bullet \Phi_2) &= \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}(l_{2,\mathcal{L}(\Phi_2)-1} + 1) \\ &\quad - l_{1,1}(l_{1,0} + 1) - l_{2,\mathcal{L}(\Phi_2)}(l_{2,\mathcal{L}(\Phi_2)-1} + 1) \\ &\leq \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}l_{2,\mathcal{L}(\Phi_2)-1}, \end{aligned} \quad (15)$$

and

(v) it holds for all $a \in C(\mathbb{R}, \mathbb{R})$ that $\mathcal{R}_a(\Phi_1 \bullet \Phi_2) \in C(\mathbb{R}^{\mathcal{I}(\Phi_2)}, \mathbb{R}^{\mathcal{O}(\Phi_1)})$ and

$$\mathcal{R}_a(\Phi_1 \bullet \Phi_2) = [\mathcal{R}_a(\Phi_1)] \circ [\mathcal{R}_a(\Phi_2)] \quad (16)$$

(cf. Definition 2.3 and Definition 2.5).

Proof of Proposition 2.6. Throughout this proof let $a \in C(\mathbb{R}, \mathbb{R})$, let $L_k \in \mathbb{N}$, $k \in \{1, 2\}$, satisfy for all $k \in \{1, 2\}$ that $L_k = \mathcal{L}(\Phi_k)$, let $((W_{k,1}, B_{k,1}), (W_{k,2}, B_{k,2}), \dots, (W_{k,L_k}, B_{k,L_k})) \in (\times_{j=1}^{L_k} (\mathbb{R}^{l_{k,j} \times l_{k,j-1}} \times \mathbb{R}^{l_{k,j}}))$, $k \in \{1, 2\}$, satisfy for all $k \in \{1, 2\}$ that

$$\Phi_k = ((W_{k,1}, B_{k,1}), (W_{k,2}, B_{k,2}), \dots, (W_{k,L_k}, B_{k,L_k})), \quad (17)$$

let $L_3 \in \mathbb{N}$, $l_{3,0}, l_{3,1}, \dots, l_{3,L_3} \in \mathbb{N}$, $\Phi_3 = ((W_{3,1}, B_{3,1}), \dots, (W_{3,L_3}, B_{3,L_3})) \in (\times_{j=1}^{L_3} (\mathbb{R}^{l_{3,j} \times l_{3,j-1}} \times \mathbb{R}^{l_{3,j}}))$ satisfy that $\Phi_3 = \Phi_1 \bullet \Phi_2$, let $x_0 \in \mathbb{R}^{l_{2,0}}$, $x_1 \in \mathbb{R}^{l_{2,1}}, \dots, x_{L_2-1} \in \mathbb{R}^{l_{2,L_2-1}}$ satisfy that

$$\forall j \in \mathbb{N} \cap (0, L_2): x_j = \mathfrak{M}_{a,l_{2,j}}(W_{2,j}x_{j-1} + B_{2,j}) \quad (18)$$

(cf. Definition 2.2), let $y_0 \in \mathbb{R}^{l_{1,0}}$, $y_1 \in \mathbb{R}^{l_{1,1}}, \dots, y_{L_1-1} \in \mathbb{R}^{l_{1,L_1-1}}$ satisfy that $y_0 = W_{2,L_2}x_{L_2-1} + B_{2,L_2}$ and

$$\forall j \in \mathbb{N} \cap (0, L_1): y_j = \mathfrak{M}_{a,l_{1,j}}(W_{1,j}y_{j-1} + B_{1,j}), \quad (19)$$

and let $z_0 \in \mathbb{R}^{l_{3,0}}$, $z_1 \in \mathbb{R}^{l_{3,1}}, \dots, z_{L_3-1} \in \mathbb{R}^{l_{3,L_3-1}}$ satisfy that $z_0 = x_0$ and

$$\forall j \in \mathbb{N} \cap (0, L_3): z_j = \mathfrak{M}_{a,l_{3,j}}(W_{3,j}z_{j-1} + B_{3,j}). \quad (20)$$

Note that (11) ensures that

$$\Phi_3 = \Phi_1 \bullet \Phi_2 = \left\{ \begin{array}{ll} ((W_{2,1}, B_{2,1}), (W_{2,2}, B_{2,2}), \dots, (W_{2,L_2-1}, B_{2,L_2-1}), \\ (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}), (W_{1,2}, B_{1,2}), \\ (W_{1,3}, B_{1,3}), \dots, (W_{1,L_1}, B_{1,L_1})) & : L_1 > 1 < L_2 \\ ((W_{1,1}W_{2,1}, W_{1,1}B_{2,1} + B_{1,1}), (W_{1,2}, B_{1,2}), \\ (W_{1,3}, B_{1,3}), \dots, (W_{1,L_1}, B_{1,L_1})) & : L_1 > 1 = L_2 \\ ((W_{2,1}, B_{2,1}), (W_{2,2}, B_{2,2}), \dots, (W_{2,L_2-1}, B_{2,L_2-1}), \\ (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1})) & : L_1 = 1 < L_2 \\ (W_{1,1}W_{2,1}, W_{1,1}B_{2,1} + B_{1,1}) & : L_1 = 1 = L_2 \end{array} \right. \quad (21)$$

Hence, we obtain that

$$\begin{aligned} [\mathcal{L}(\Phi_1 \bullet \Phi_2) - 1] &= [(L_2 - 1) + 1 + (L_1 - 1)] - 1 \\ &= [L_1 - 1] + [L_2 - 1] = [\mathcal{L}(\Phi_1) - 1] + [\mathcal{L}(\Phi_2) - 1] \end{aligned} \quad (22)$$

$$\text{and} \quad \mathcal{D}(\Phi_1 \bullet \Phi_2) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1}, l_{1,2}, \dots, l_{1,L_1}). \quad (23)$$

This establishes items (i)–(iii). In addition, observe that (23) demonstrates that

$$\begin{aligned} \mathcal{P}(\Phi_1 \bullet \Phi_2) &= \sum_{j=1}^{L_3} l_{3,j}(l_{3,j-1} + 1) \\ &= \left[\sum_{j=1}^{L_2-1} l_{3,j}(l_{3,j-1} + 1) \right] + l_{3,L_2}(l_{3,L_2-1} + 1) + \left[\sum_{j=L_2+1}^{L_3} l_{3,j}(l_{3,j-1} + 1) \right] \\ &= \left[\sum_{j=1}^{L_2-1} l_{2,j}(l_{2,j-1} + 1) \right] + l_{1,1}(l_{2,L_2-1} + 1) + \left[\sum_{j=L_2+1}^{L_3} l_{1,j-L_2+1}(l_{1,j-L_2} + 1) \right] \\ &= \left[\sum_{j=1}^{L_2-1} l_{2,j}(l_{2,j-1} + 1) \right] + \left[\sum_{j=2}^{L_1} l_{1,j}(l_{1,j-1} + 1) \right] + l_{1,1}(l_{2,L_2-1} + 1) \\ &= \left[\sum_{j=1}^{L_2} l_{2,j}(l_{2,j-1} + 1) \right] + \left[\sum_{j=1}^{L_1} l_{1,j}(l_{1,j-1} + 1) \right] + l_{1,1}(l_{2,L_2-1} + 1) \\ &\quad - l_{2,L_2}(l_{2,L_2-1} + 1) - l_{1,1}(l_{1,0} + 1) \\ &= \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}(l_{2,L_2-1} + 1) - l_{2,L_2}(l_{2,L_2-1} + 1) \\ &\quad - l_{1,1}(l_{1,0} + 1) \\ &\leq \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}l_{2,L_2-1}. \end{aligned} \quad (24)$$

This establishes item (iv). Moreover, observe that (21) and the fact that $a \in C(\mathbb{R}, \mathbb{R})$ ensure that

$$\mathcal{R}_a(\Phi_1 \bullet \Phi_2) \in C(\mathbb{R}^{l_{2,0}}, \mathbb{R}^{l_{1,L_1}}) = C(\mathbb{R}^{\mathcal{I}(\Phi_2)}, \mathbb{R}^{\mathcal{O}(\Phi_1)}). \quad (25)$$

Next note that (22) implies that $L_3 = L_1 + L_2 - 1$. This, (21), and (23) ensure that

$$(l_{3,0}, l_{3,1}, \dots, l_{3,L_1+L_2-1}) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1}, l_{1,2}, \dots, l_{1,L_1}), \quad (26)$$

$$[\forall j \in \mathbb{N} \cap (0, L_2): (W_{3,j}, B_{3,j}) = (W_{2,j}, B_{2,j})], \quad (27)$$

$$(W_{3,L_2}, B_{3,L_2}) = (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}), \quad (28)$$

$$\text{and} \quad [\forall j \in \mathbb{N} \cap (L_2, L_1 + L_2): (W_{3,j}, B_{3,j}) = (W_{1,j+1-L_2}, B_{1,j+1-L_2})]. \quad (29)$$

This, (18), (20), and induction imply that for all $j \in \mathbb{N}_0 \cap [0, L_2)$ it holds that $z_j = x_j$. Combining this with (28) and the fact that $y_0 = W_{2,L_2}x_{L_2-1} + B_{2,L_2}$ ensures that

$$\begin{aligned} W_{3,L_2}z_{L_2-1} + B_{3,L_2} &= W_{3,L_2}x_{L_2-1} + B_{3,L_2} \\ &= W_{1,1}W_{2,L_2}x_{L_2-1} + W_{1,1}B_{2,L_2} + B_{1,1} \\ &= W_{1,1}(W_{2,L_2}x_{L_2-1} + B_{2,L_2}) + B_{1,1} = W_{1,1}y_0 + B_{1,1}. \end{aligned} \quad (30)$$

Next we claim that for all $j \in \mathbb{N} \cap [L_2, L_1 + L_2)$ it holds that

$$W_{3,j}z_{j-1} + B_{3,j} = W_{1,j+1-L_2}y_{j-L_2} + B_{1,j+1-L_2}. \quad (31)$$

We prove (31) by induction on $j \in \mathbb{N} \cap [L_2, L_1 + L_2)$. Note that (30) establishes (31) in the base case $j = L_2$. For the induction step note that the fact that $L_3 = L_1 + L_2 - 1$, (19), (20), (26), and (29) imply that for all $j \in \mathbb{N} \cap [L_2, \infty) \cap (0, L_1 + L_2 - 1)$ with

$$W_{3,j}z_{j-1} + B_{3,j} = W_{1,j+1-L_2}y_{j-L_2} + B_{1,j+1-L_2} \quad (32)$$

it holds that

$$\begin{aligned} W_{3,j+1}z_j + B_{3,j+1} &= W_{3,j+1}\mathfrak{M}_{a,l_3,j}(W_{3,j}z_{j-1} + B_{3,j}) + B_{3,j+1} \\ &= W_{1,j+2-L_2}\mathfrak{M}_{a,l_1,j+1-L_2}(W_{1,j+1-L_2}y_{j-L_2} + B_{1,j+1-L_2}) + B_{1,j+2-L_2} \\ &= W_{1,j+2-L_2}y_{j+1-L_2} + B_{1,j+2-L_2}. \end{aligned} \quad (33)$$

Induction hence proves (31). Next observe that (31) and the fact that $L_3 = L_1 + L_2 - 1$ assure that

$$W_{3,L_3}z_{L_3-1} + B_{3,L_3} = W_{3,L_1+L_2-1}z_{L_1+L_2-2} + B_{3,L_1+L_2-1} = W_{1,L_1}y_{L_1-1} + B_{1,L_1}. \quad (34)$$

The fact that $\Phi_3 = \Phi_1 \bullet \Phi_2$, (18), (19), and (20) therefore prove that

$$\begin{aligned} [\mathcal{R}_a(\Phi_1 \bullet \Phi_2)](x_0) &= [\mathcal{R}_a(\Phi_3)](x_0) = [\mathcal{R}_a(\Phi_3)](z_0) = W_{3,L_3}z_{L_3-1} + B_{3,L_3} \\ &= W_{1,L_1}y_{L_1-1} + B_{1,L_1} = [\mathcal{R}_a(\Phi_1)](y_0) \\ &= [\mathcal{R}_a(\Phi_1)](W_{2,L_2}x_{L_2-1} + B_{2,L_2}) \\ &= [\mathcal{R}_a(\Phi_1)]([\mathcal{R}_a(\Phi_2)](x_0)) = [(\mathcal{R}_a(\Phi_1)) \circ (\mathcal{R}_a(\Phi_2))](x_0). \end{aligned} \quad (35)$$

Combining this with (25) establishes item (v). The proof of Proposition 2.6 is thus completed. \square

Corollary 2.7. *Let $L_1, L_2, L_3 \in \mathbb{N}$, $l_{1,0}, l_{1,1}, \dots, l_{1,L_1}, l_{2,0}, l_{2,1}, \dots, l_{2,L_2}, l_{3,0}, l_{3,1}, \dots, l_{3,L_3} \in \mathbb{N}$ satisfy that $l_{1,0} = l_{2,L_2}$ and let $\Phi_k = ((W_{k,1}, B_{k,1}), (W_{k,2}, B_{k,2}), \dots, (W_{k,L_k}, B_{k,L_k})) \in (\times_{j=1}^{L_k} (\mathbb{R}^{l_{k,j} \times l_{k,j-1}} \times \mathbb{R}^{l_{k,j}}))$, $k \in \{1, 2, 3\}$, satisfy that $\Phi_3 = \Phi_1 \bullet \Phi_2$ (cf. Definition 2.1 and Definition 2.5). Then*

(i) it holds that

$$L_3 = \mathcal{L}(\Phi_3) = \mathcal{L}(\Phi_1) + \mathcal{L}(\Phi_2) - 1 = L_1 + L_2 - 1 \geq \max\{L_1, L_2\}, \quad (36)$$

(ii) it holds for all $j \in \mathbb{N} \cap (0, L_2)$ that

$$(W_{3,j}, B_{3,j}) = (W_{2,j}, B_{2,j}), \quad (37)$$

(iii) it holds that

$$(W_{3,L_2}, B_{3,L_2}) = (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}), \quad (38)$$

and

(iv) it holds for all $j \in \mathbb{N} \cap (L_2, L_1 + L_2) = \mathbb{N} \cap (L_2, \infty) \cap [1, L_3]$ that

$$(W_{3,j}, B_{3,j}) = (W_{1,j-L_2+1}, B_{1,j-L_2+1}). \quad (39)$$

Proof of Corollary 2.7. Observe that item (ii) in Proposition 2.6 proves item (i). Moreover, note that (11) establishes items (ii)–(iv). The proof of Corollary 2.7 is thus completed. \square

2.2.2 Associativity of standard compositions of ANNs

Lemma 2.8. *Let $\Phi_1, \Phi_2, \Phi_3 \in \mathbf{N}$ satisfy that $\mathcal{I}(\Phi_1) = \mathcal{O}(\Phi_2)$ and $\mathcal{I}(\Phi_2) = \mathcal{O}(\Phi_3)$ (cf. Definition 2.1). Then it holds that*

$$(\Phi_1 \bullet \Phi_2) \bullet \Phi_3 = \Phi_1 \bullet (\Phi_2 \bullet \Phi_3) \quad (40)$$

(cf. Definition 2.5).

Proof of Lemma 2.8. Throughout this proof let $\Phi_4, \Phi_5, \Phi_6, \Phi_7 \in \mathbf{N}$ satisfy that $\Phi_4 = \Phi_1 \bullet \Phi_2$, $\Phi_5 = \Phi_2 \bullet \Phi_3$, $\Phi_6 = \Phi_4 \bullet \Phi_3$, and $\Phi_7 = \Phi_1 \bullet \Phi_5$, let $L_k \in \mathbb{N}$, $k \in \{1, 2, \dots, 7\}$, satisfy for all $k \in \{1, 2, \dots, 7\}$ that $L_k = \mathcal{L}(\Phi_k)$, let $l_{k,0}, l_{k,1}, \dots, l_{k,L_k} \in \mathbb{N}$, $k \in \{1, 2, \dots, 7\}$, and let $((W_{k,1}, B_{k,1}), (W_{k,2}, B_{k,2}), \dots, (W_{k,L_k}, B_{k,L_k})) \in (\times_{j=1}^{L_k} (\mathbb{R}^{l_{k,j} \times l_{k,j-1}} \times \mathbb{R}^{l_{k,j}}))$, $k \in \{1, 2, \dots, 7\}$, satisfy for all $k \in \{1, 2, \dots, 7\}$ that

$$\Phi_k = ((W_{k,1}, B_{k,1}), (W_{k,2}, B_{k,2}), \dots, (W_{k,L_k}, B_{k,L_k})). \quad (41)$$

Observe that item (ii) in Proposition 2.6 and the fact that for all $k \in \{1, 2, 3\}$ it holds that $\mathcal{L}(\Phi_k) = L_k$ proves that

$$\begin{aligned} \mathcal{L}(\Phi_6) &= \mathcal{L}((\Phi_1 \bullet \Phi_2) \bullet \Phi_3) = \mathcal{L}(\Phi_1 \bullet \Phi_2) + \mathcal{L}(\Phi_3) - 1 \\ &= \mathcal{L}(\Phi_1) + \mathcal{L}(\Phi_2) + \mathcal{L}(\Phi_3) - 2 = L_1 + L_2 + L_3 - 2 \\ &= \mathcal{L}(\Phi_1) + \mathcal{L}(\Phi_2 \bullet \Phi_3) - 1 = \mathcal{L}(\Phi_1 \bullet (\Phi_2 \bullet \Phi_3)) = \mathcal{L}(\Phi_7). \end{aligned} \quad (42)$$

Next note that Corollary 2.7, (41), and the fact that $\Phi_4 = \Phi_1 \bullet \Phi_2$ imply that

$$[\forall j \in \mathbb{N} \cap (0, L_2): (W_{4,j}, B_{4,j}) = (W_{2,j}, B_{2,j})], \quad (43)$$

$$(W_{4,L_2}, B_{4,L_2}) = (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}), \quad (44)$$

$$\text{and } [\forall j \in \mathbb{N} \cap (L_2, L_1 + L_2): (W_{4,j}, B_{4,j}) = (W_{1,j+1-L_2}, B_{1,j+1-L_2})]. \quad (45)$$

Hence, we obtain that

$$[\forall j \in \mathbb{N} \cap (L_3 - 1, L_2 + L_3 - 1): \\ (W_{4,j+1-L_3}, B_{4,j+1-L_3}) = (W_{2,j+1-L_3}, B_{2,j+1-L_3})], \quad (46)$$

$$(W_{4,L_2}, B_{4,L_2}) = (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}), \quad (47)$$

and

$$[\forall j \in \mathbb{N} \cap (L_2 + L_3 - 1, L_1 + L_2 + L_3 - 1): \\ (W_{4,j+1-L_3}, B_{4,j+1-L_3}) = (W_{1,j+2-L_2-L_3}, B_{1,j+2-L_2-L_3})]. \quad (48)$$

In addition, observe that Corollary 2.7, (41), and the fact that $\Phi_5 = \Phi_2 \bullet \Phi_3$ demonstrate that

$$[\forall j \in \mathbb{N} \cap (0, L_3): (W_{5,j}, B_{5,j}) = (W_{3,j}, B_{3,j})], \quad (49)$$

$$(W_{5,L_3}, B_{5,L_3}) = (W_{2,1}W_{3,L_3}, W_{2,1}B_{3,L_3} + B_{2,1}), \quad (50)$$

$$\text{and } [\forall j \in \mathbb{N} \cap (L_3, L_2 + L_3): (W_{5,j}, B_{5,j}) = (W_{2,j+1-L_3}, B_{2,j+1-L_3})]. \quad (51)$$

Moreover, note that Corollary 2.7, (41), and the fact that $\Phi_6 = \Phi_4 \bullet \Phi_3$ ensure that

$$[\forall j \in \mathbb{N} \cap (0, L_3): (W_{6,j}, B_{6,j}) = (W_{3,j}, B_{3,j})], \quad (52)$$

$$(W_{6,L_3}, B_{6,L_3}) = (W_{4,1}W_{3,L_3}, W_{4,1}B_{3,L_3} + B_{4,1}), \quad (53)$$

$$\text{and } [\forall j \in \mathbb{N} \cap (L_3, L_4 + L_3): (W_{6,j}, B_{6,j}) = (W_{4,j+1-L_3}, B_{4,j+1-L_3})]. \quad (54)$$

Furthermore, observe that Corollary 2.7, (41), and the fact that $\Phi_7 = \Phi_1 \bullet \Phi_5$ show that

$$[\forall j \in \mathbb{N} \cap (0, L_5): (W_{7,j}, B_{7,j}) = (W_{5,j}, B_{5,j})], \quad (55)$$

$$(W_{7,L_5}, B_{7,L_5}) = (W_{1,1}W_{5,L_5}, W_{1,1}B_{5,L_5} + B_{1,1}), \quad (56)$$

$$\text{and } [\forall j \in \mathbb{N} \cap (L_5, L_1 + L_5): (W_{7,j}, B_{7,j}) = (W_{1,j+1-L_5}, B_{1,j+1-L_5})]. \quad (57)$$

This, the fact that $L_3 \leq L_2 + L_3 - 1 = L_5$, (49), and (52) imply that for all $j \in \mathbb{N} \cap (0, L_3)$ it holds that

$$(W_{6,j}, B_{6,j}) = (W_{3,j}, B_{3,j}) = (W_{5,j}, B_{5,j}) = (W_{7,j}, B_{7,j}). \quad (58)$$

In addition, observe that (43), (44), (49), (50), (53), (55), (56), and the fact that $L_5 = L_2 + L_3 - 1$ demonstrate that

$$\begin{aligned}
& (W_{6,L_3}, B_{6,L_3}) = (W_{4,1}W_{3,L_3}, W_{4,1}B_{3,L_3} + B_{4,1}) \\
& = \begin{cases} (W_{2,1}W_{3,L_3}, W_{2,1}B_{3,L_3} + B_{2,1}) & : L_2 > 1 \\ (W_{1,1}W_{2,1}W_{3,L_3}, W_{1,1}W_{2,1}B_{3,L_3} + W_{1,1}B_{2,1} + B_{1,1}) & : L_2 = 1 \end{cases} \\
& = \begin{cases} (W_{2,1}W_{3,L_3}, W_{2,1}B_{3,L_3} + B_{2,1}) & : L_2 > 1 \\ (W_{1,1}(W_{2,1}W_{3,L_3}), W_{1,1}(W_{2,1}B_{3,L_3} + B_{2,1}) + B_{1,1}) & : L_2 = 1 \end{cases} \quad (59) \\
& = \begin{cases} (W_{5,L_3}, B_{5,L_3}) & : L_2 > 1 \\ (W_{1,1}W_{5,L_3}, W_{1,1}B_{5,L_3} + B_{1,1}) & : L_2 = 1 \end{cases} \\
& = (W_{7,L_3}, B_{7,L_3}).
\end{aligned}$$

Next note that the fact that $L_5 = L_2 + L_3 - 1 < L_1 + L_2 + L_3 - 1 = L_3 + L_4$, (54), (46), (51), and (55) ensure that for all $j \in \mathbb{N}$ with $L_3 < j < L_5$ it holds that

$$\begin{aligned}
(W_{6,j}, B_{6,j}) &= (W_{4,j+1-L_3}, B_{4,j+1-L_3}) = (W_{2,j+1-L_3}, B_{2,j+1-L_3}) \\
&= (W_{5,j}, B_{5,j}) = (W_{7,j}, B_{7,j}). \quad (60)
\end{aligned}$$

Moreover, observe that the fact that $L_5 = L_2 + L_3 - 1 < L_1 + L_2 + L_3 - 1 = L_3 + L_4$, (54), (59), (44), (51), and (56) prove that

$$\begin{aligned}
(W_{6,L_5}, B_{6,L_5}) &= \begin{cases} (W_{4,L_5+1-L_3}, B_{4,L_5+1-L_3}) & : L_2 > 1 \\ (W_{6,L_3}, B_{6,L_3}) & : L_2 = 1 \end{cases} \\
&= \begin{cases} (W_{4,L_2}, B_{4,L_2}) & : L_2 > 1 \\ (W_{7,L_3}, B_{7,L_3}) & : L_2 = 1 \end{cases} \\
&= \begin{cases} (W_{1,1}W_{2,L_2}, W_{1,1}B_{2,L_2} + B_{1,1}) & : L_2 > 1 \\ (W_{7,L_5}, B_{7,L_5}) & : L_2 = 1 \end{cases} \quad (61) \\
&= \begin{cases} (W_{1,1}W_{5,L_5}, W_{1,1}B_{5,L_5} + B_{1,1}) & : L_2 > 1 \\ (W_{7,L_5}, B_{7,L_5}) & : L_2 = 1 \end{cases} \\
&= (W_{7,L_5}, B_{7,L_5}).
\end{aligned}$$

Furthermore, note that (54), (48), (57), and the fact that $L_5 = L_2 + L_3 - 1 \geq L_3$ assure that for all $j \in \mathbb{N}$ with $L_5 < j \leq L_6$ it holds that

$$\begin{aligned}
(W_{6,j}, B_{6,j}) &= (W_{4,j+1-L_3}, B_{4,j+1-L_3}) = (W_{1,j+2-L_2-L_3}, B_{1,j+2-L_2-L_3}) \\
&= (W_{1,j+1-L_5}, B_{1,j+1-L_5}) = (W_{7,j}, B_{7,j}). \quad (62)
\end{aligned}$$

Combining this with (42), (58), (59), (60), and (61) establishes that

$$(\Phi_1 \bullet \Phi_2) \bullet \Phi_3 = \Phi_4 \bullet \Phi_3 = \Phi_6 = \Phi_7 = \Phi_1 \bullet \Phi_5 = \Phi_1 \bullet (\Phi_2 \bullet \Phi_3). \quad (63)$$

The proof of Lemma 2.8 is thus completed. \square

2.2.3 Compositions of ANNs and affine linear transformations

Corollary 2.9. *Let $\Phi \in \mathbf{N}$ (cf. Definition 2.1). Then*

(i) *it holds for all $\mathbb{A} \in \mathbf{N}$ with $\mathcal{L}(\mathbb{A}) = 1$ and $\mathcal{I}(\mathbb{A}) = \mathcal{O}(\Phi)$ that*

$$\mathcal{P}(\mathbb{A} \bullet \Phi) \leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A})}{\mathcal{O}(\Phi)} \right\} \right] \mathcal{P}(\Phi) \quad (64)$$

and

(ii) *it holds for all $\mathbb{A} \in \mathbf{N}$ with $\mathcal{L}(\mathbb{A}) = 1$ and $\mathcal{I}(\Phi) = \mathcal{O}(\mathbb{A})$ that*

$$\mathcal{P}(\Phi \bullet \mathbb{A}) \leq \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A})+1}{\mathcal{I}(\Phi)+1} \right\} \right] \mathcal{P}(\Phi) \quad (65)$$

(cf. Definition 2.5).

Proof of Corollary 2.9. Throughout this proof let $L \in \mathbf{N}$, $l_0, l_1, \dots, l_L \in \mathbf{N}$, $\mathbb{A}_1, \mathbb{A}_2 \in \mathbf{N}$ satisfy that $\mathcal{L}(\mathbb{A}_1) = \mathcal{L}(\mathbb{A}_2) = 1$, $\mathcal{I}(\mathbb{A}_1) = \mathcal{O}(\Phi)$, $\mathcal{I}(\Phi) = \mathcal{O}(\mathbb{A}_2)$, and $\mathcal{D}(\Phi) = (l_0, l_1, \dots, l_L)$. Observe that item (iv) in Proposition 2.6, the fact that $\mathcal{O}(\Phi) = l_L$, the fact that $\mathcal{I}(\Phi) = l_0$, and the fact that for all $k \in \{1, 2\}$ it holds that $\mathcal{D}(\mathbb{A}_k) = (\mathcal{I}(\mathbb{A}_k), \mathcal{O}(\mathbb{A}_k))$ ensure that

$$\begin{aligned} \mathcal{P}(\mathbb{A}_1 \bullet \Phi) &= \left[\sum_{m=1}^{L-1} l_m(l_{m-1} + 1) \right] + [\mathcal{O}(\mathbb{A}_1)](l_{L-1} + 1) \\ &= \left[\sum_{m=1}^{L-1} l_m(l_{m-1} + 1) \right] + \left[\frac{\mathcal{O}(\mathbb{A}_1)}{l_L} \right] l_L(l_{L-1} + 1) \\ &\leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{l_L} \right\} \right] \left[\sum_{m=1}^{L-1} l_m(l_{m-1} + 1) \right] + \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{l_L} \right\} \right] l_L(l_{L-1} + 1) \\ &= \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{l_L} \right\} \right] \left[\sum_{m=1}^L l_m(l_{m-1} + 1) \right] = \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{\mathcal{O}(\Phi)} \right\} \right] \mathcal{P}(\Phi) \end{aligned} \quad (66)$$

and

$$\begin{aligned}
\mathcal{P}(\Phi \bullet \mathbb{A}_2) &= \left[\sum_{m=2}^L l_m(l_{m-1} + 1) \right] + l_1[\mathcal{I}(\mathbb{A}_2) + 1] \\
&= \left[\sum_{m=2}^L l_m(l_{m-1} + 1) \right] + \left[\frac{\mathcal{I}(\mathbb{A}_2)+1}{l_0+1} \right] l_1(l_0 + 1) \\
&\leq \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{l_0+1} \right\} \right] \left[\sum_{m=2}^L l_m(l_{m-1} + 1) \right] + \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{l_0+1} \right\} \right] l_1(l_0 + 1) \\
&= \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{l_0+1} \right\} \right] \left[\sum_{m=1}^L l_m(l_{m-1} + 1) \right] = \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{\mathcal{I}(\Phi)+1} \right\} \right] \mathcal{P}(\Phi).
\end{aligned} \tag{67}$$

This establishes items (i)–(ii). The proof of Corollary 2.9 is thus completed. \square

2.2.4 Powers and extensions of ANNs

Definition 2.10. Let $d \in \mathbb{N}$. Then we denote by $I_d \in \mathbb{R}^{d \times d}$ the identity matrix in $\mathbb{R}^{d \times d}$.

Definition 2.11. We denote by $(\cdot)^{\bullet n}: \{\Phi \in \mathbf{N}: \mathcal{I}(\Phi) = \mathcal{O}(\Phi)\} \rightarrow \mathbf{N}$, $n \in \mathbb{N}_0$, the functions which satisfy for all $n \in \mathbb{N}_0$, $\Phi \in \mathbf{N}$ with $\mathcal{I}(\Phi) = \mathcal{O}(\Phi)$ that

$$\Phi^{\bullet n} = \begin{cases} (I_{\mathcal{O}(\Phi)}, (0, 0, \dots, 0)) \in \mathbb{R}^{\mathcal{O}(\Phi) \times \mathcal{O}(\Phi)} \times \mathbb{R}^{\mathcal{O}(\Phi)} & : n = 0 \\ \Phi \bullet (\Phi^{\bullet(n-1)}) & : n \in \mathbb{N} \end{cases} \tag{68}$$

(cf. Definition 2.1, Definition 2.5, and Definition 2.10).

Definition 2.12 (Extension of ANNs). Let $L \in \mathbb{N}$, $\Psi \in \mathbf{N}$ satisfy that $\mathcal{I}(\Psi) = \mathcal{O}(\Psi)$. Then we denote by $\mathcal{E}_{L,\Psi}: \{\Phi \in \mathbf{N}: (\mathcal{L}(\Phi) \leq L \text{ and } \mathcal{O}(\Phi) = \mathcal{I}(\Psi))\} \rightarrow \mathbf{N}$ the function which satisfies for all $\Phi \in \mathbf{N}$ with $\mathcal{L}(\Phi) \leq L$ and $\mathcal{O}(\Phi) = \mathcal{I}(\Psi)$ that

$$\mathcal{E}_{L,\Psi}(\Phi) = (\Psi^{\bullet(L-\mathcal{L}(\Phi))}) \bullet \Phi \tag{69}$$

(cf. Definition 2.1, Definition 2.5, and Definition 2.11).

Lemma 2.13. Let $d, \mathbf{i} \in \mathbb{N}$, $\Psi \in \mathbf{N}$ satisfy that $\mathcal{D}(\Psi) = (d, \mathbf{i}, d)$ (cf. Definition 2.1). Then

(i) it holds for all $n \in \mathbb{N}_0$ that $\mathcal{L}(\Psi^{\bullet n}) = n + 1$, $\mathcal{D}(\Psi^{\bullet n}) \in \mathbb{N}^{n+2}$, and

$$\mathcal{D}(\Psi^{\bullet n}) = \begin{cases} (d, d) & : n = 0 \\ (d, \mathbf{i}, \mathbf{i}, \dots, \mathbf{i}, d) & : n \in \mathbb{N} \end{cases} \tag{70}$$

and

(ii) it holds for all $\Phi \in \mathbf{N}$, $L \in \mathbb{N} \cap [\mathcal{L}(\Phi), \infty)$ with $\mathcal{O}(\Phi) = d$ that $\mathcal{L}(\mathcal{E}_{L,\Psi}(\Phi)) = L$ and

$$\begin{aligned} & \mathcal{P}(\mathcal{E}_{L,\Psi}(\Phi)) \\ & \leq \begin{cases} \mathcal{P}(\Phi) & : \mathcal{L}(\Phi) = L \\ [(\max\{1, \frac{i}{d}\})\mathcal{P}(\Phi) + ((L - \mathcal{L}(\Phi) - 1)\mathbf{i} + d)(\mathbf{i} + 1)] & : \mathcal{L}(\Phi) < L \end{cases} \quad (71) \end{aligned}$$

(cf. Definition 2.11 and Definition 2.12).

Proof of Lemma 2.13. Throughout this proof let $\Phi \in \mathbf{N}$, $l_0, l_1, \dots, l_{\mathcal{L}(\Phi)} \in \mathbb{N}$ satisfy that $\mathcal{O}(\Phi) = d$ and $\mathcal{D}(\Phi) = (l_0, l_1, \dots, l_{\mathcal{L}(\Phi)}) \in \mathbb{N}^{\mathcal{L}(\Phi)+1}$ and let $a_{L,k} \in \mathbb{N}$, $k \in \mathbb{N}_0 \cap [0, L]$, $L \in \mathbb{N} \cap [\mathcal{L}(\Phi), \infty)$, satisfy for all $L \in \mathbb{N} \cap [\mathcal{L}(\Phi), \infty)$, $k \in \mathbb{N}_0 \cap [0, L]$ that

$$a_{L,k} = \begin{cases} l_k & : k < \mathcal{L}(\Phi) \\ \mathbf{i} & : \mathcal{L}(\Phi) \leq k < L \\ d & : k = L \end{cases} \quad (72)$$

We claim that for all $n \in \mathbb{N}_0$ it holds that

$$\mathcal{L}(\Psi^{\bullet n}) = n + 1 \quad \text{and} \quad \mathbb{N}^{n+2} \ni \mathcal{D}(\Psi^{\bullet n}) = \begin{cases} (d, d) & : n = 0 \\ (d, \mathbf{i}, \mathbf{i}, \dots, \mathbf{i}, d) & : n \in \mathbb{N} \end{cases} \quad (73)$$

We now prove (73) by induction on $n \in \mathbb{N}_0$. Note that the fact that $\Psi^{\bullet 0} = (\mathbf{I}_d, 0) \in \mathbb{R}^{d \times d} \times \mathbb{R}^d$ (cf. Definition 2.10) establishes (70) in the base case $n = 0$. For the induction step $\mathbb{N}_0 \ni n \rightarrow n + 1 \in \mathbb{N}$ assume that there exists $n \in \mathbb{N}_0$ such that

$$\mathcal{L}(\Psi^{\bullet n}) = n + 1 \quad \text{and} \quad \mathbb{N}^{n+2} \ni \mathcal{D}(\Psi^{\bullet n}) = \begin{cases} (d, d) & : n = 0 \\ (d, \mathbf{i}, \mathbf{i}, \dots, \mathbf{i}, d) & : n \in \mathbb{N} \end{cases} \quad (74)$$

Observe that Lemma 2.4, (68), items (i)–(ii) in Proposition 2.6, (74), and the hypothesis that $\mathcal{D}(\Psi) = (d, \mathbf{i}, d)$ imply that

$$\begin{aligned} \mathcal{L}(\Psi^{\bullet(n+1)}) &= \mathcal{L}(\Psi \bullet (\Psi^{\bullet n})) = \mathcal{L}(\Psi) + \mathcal{L}(\Psi^{\bullet n}) - 1 = 2 + (n + 1) - 1 = (n + 1) + 1 \\ \text{and} \quad \mathcal{D}(\Psi^{\bullet(n+1)}) &= \mathcal{D}(\Psi \bullet (\Psi^{\bullet n})) = (d, \mathbf{i}, \mathbf{i}, \dots, \mathbf{i}, d) \in \mathbb{N}^{n+3}. \end{aligned} \quad (75)$$

Induction thus proves (73). Next note that (73) establishes item (i). In addition, observe that items (i)–(ii) in Proposition 2.6, item (i), (69), and (72) ensure that for all $L \in \mathbb{N} \cap [\mathcal{L}(\Phi), \infty)$ it holds that

$$\begin{aligned} \mathcal{L}(\mathcal{E}_{L,\Psi}(\Phi)) &= \mathcal{L}((\Psi^{\bullet(L-\mathcal{L}(\Phi))}) \bullet \Phi) = \mathcal{L}(\Psi^{\bullet(L-\mathcal{L}(\Phi))}) + \mathcal{L}(\Phi) - 1 \\ &= (L - \mathcal{L}(\Phi) + 1) + \mathcal{L}(\Phi) - 1 = L \end{aligned} \quad (76)$$

and

$$\mathcal{D}(\mathcal{E}_{L,\Psi}(\Phi)) = \mathcal{D}((\Psi^{\bullet(L-\mathcal{L}(\Phi))}) \bullet \Phi) = (a_{L,0}, a_{L,1}, \dots, a_{L,L}). \quad (77)$$

Combining this with (72) demonstrates that

$$\mathcal{L}(\mathcal{E}_{\mathcal{L}(\Phi),\Psi}(\Phi)) = \mathcal{L}(\Phi) \quad (78)$$

and

$$\begin{aligned} \mathcal{D}(\mathcal{E}_{\mathcal{L}(\Phi),\Psi}(\Phi)) &= (a_{\mathcal{L}(\Phi),0}, a_{\mathcal{L}(\Phi),1}, \dots, a_{\mathcal{L}(\Phi),\mathcal{L}(\Phi)}) \\ &= (l_0, l_1, \dots, l_{\mathcal{L}(\Phi)}) = \mathcal{D}(\Phi). \end{aligned} \quad (79)$$

Hence, we obtain that

$$\mathcal{P}(\mathcal{E}_{\mathcal{L}(\Phi),\Psi}(\Phi)) = \mathcal{P}(\Phi). \quad (80)$$

Next note that (72), (77), and the fact that $l_{\mathcal{L}(\Phi)} = \mathcal{O}(\Phi) = d$ imply that for all

$L \in \mathbb{N} \cap (\mathcal{L}(\Phi), \infty)$ it holds that

$$\begin{aligned}
\mathcal{P}(\mathcal{E}_{L,\Psi}(\Phi)) &= \sum_{k=1}^L a_{L,k}(a_{L,k-1} + 1) \\
&= \left[\sum_{k=1}^{\mathcal{L}(\Phi)-1} a_{L,k}(a_{L,k-1} + 1) \right] + \left[\sum_{k=\mathcal{L}(\Phi)}^L a_{L,k}(a_{L,k-1} + 1) \right] \\
&= \left[\sum_{k=1}^{\mathcal{L}(\Phi)-1} l_k(l_{k-1} + 1) \right] + \left[\sum_{k=\mathcal{L}(\Phi)}^{\mathcal{L}(\Phi)} a_{L,k}(a_{L,k-1} + 1) \right] \\
&\quad + \left[\sum_{k=\mathcal{L}(\Phi)+1}^L a_{L,k}(a_{L,k-1} + 1) \right] \\
&= \left[\sum_{k=1}^{\mathcal{L}(\Phi)-1} l_k(l_{k-1} + 1) \right] + a_{L,\mathcal{L}(\Phi)}(a_{L,\mathcal{L}(\Phi)-1} + 1) \\
&\quad + \left[\sum_{k=\mathcal{L}(\Phi)+1}^{L-1} a_{L,k}(a_{L,k-1} + 1) \right] + \left[\sum_{k=L}^L a_{L,k}(a_{L,k-1} + 1) \right] \tag{81} \\
&= \left[\sum_{k=1}^{\mathcal{L}(\Phi)-1} l_k(l_{k-1} + 1) \right] + \mathbf{i}(l_{\mathcal{L}(\Phi)-1} + 1) \\
&\quad + (L - 1 - (\mathcal{L}(\Phi) + 1) + 1)\mathbf{i}(\mathbf{i} + 1) + a_{L,L}(a_{L,L-1} + 1) \\
&= \left[\sum_{k=1}^{\mathcal{L}(\Phi)-1} l_k(l_{k-1} + 1) \right] + \frac{\mathbf{i}}{d}[l_{\mathcal{L}(\Phi)}(l_{\mathcal{L}(\Phi)-1} + 1)] \\
&\quad + (L - \mathcal{L}(\Phi) - 1)\mathbf{i}(\mathbf{i} + 1) + d(\mathbf{i} + 1) \\
&\leq \left[\max\{1, \frac{\mathbf{i}}{d}\} \right] \left[\sum_{k=1}^{\mathcal{L}(\Phi)} l_k(l_{k-1} + 1) \right] + (L - \mathcal{L}(\Phi) - 1)\mathbf{i}(\mathbf{i} + 1) + d(\mathbf{i} + 1) \\
&= \left[\max\{1, \frac{\mathbf{i}}{d}\} \right] \mathcal{P}(\Phi) + (L - \mathcal{L}(\Phi) - 1)\mathbf{i}(\mathbf{i} + 1) + d(\mathbf{i} + 1).
\end{aligned}$$

Combining this with (80) establishes (71). The proof of Lemma 2.13 is thus completed. \square

Lemma 2.14. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $\mathbb{I} \in \mathbb{N}$ satisfy for all $x \in \mathbb{R}^{\mathcal{I}(\mathbb{I})}$ that $\mathcal{I}(\mathbb{I}) = \mathcal{O}(\mathbb{I})$ and $(\mathcal{R}_a(\mathbb{I}))(x) = x$ (cf. Definition 2.1 and Definition 2.3). Then*

(i) *it holds for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^{\mathcal{I}(\mathbb{I})}$ that*

$$\mathcal{R}_a(\mathbb{I}^{\bullet n}) \in C(\mathbb{R}^{\mathcal{I}(\mathbb{I})}, \mathbb{R}^{\mathcal{I}(\mathbb{I})}) \quad \text{and} \quad (\mathcal{R}_a(\mathbb{I}^{\bullet n}))(x) = x \tag{82}$$

and

(ii) it holds for all $\Phi \in \mathbf{N}$, $L \in \mathbb{N} \cap [\mathcal{L}(\Phi), \infty)$, $x \in \mathbb{R}^{\mathcal{I}(\Phi)}$ with $\mathcal{O}(\Phi) = \mathcal{I}(\mathbb{I})$ that

$$\mathcal{R}_a(\mathcal{E}_{L,\mathbb{I}}(\Phi)) \in C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{O}(\Phi)}) \quad \text{and} \quad (\mathcal{R}_a(\mathcal{E}_{L,\mathbb{I}}(\Phi)))(x) = (\mathcal{R}_a(\Phi))(x) \quad (83)$$

(cf. Definition 2.11 and Definition 2.12).

Proof of Lemma 2.14. Throughout this proof let $\Phi \in \mathbf{N}$, $L, d \in \mathbb{N}$ satisfy that $\mathcal{L}(\Phi) \leq L$ and $\mathcal{I}(\mathbb{I}) = \mathcal{O}(\Phi) = d$. We claim that for all $n \in \mathbb{N}_0$ it holds that

$$\mathcal{R}_a(\mathbb{I}^{\bullet n}) \in C(\mathbb{R}^d, \mathbb{R}^d) \quad \text{and} \quad \forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{I}^{\bullet n}))(x) = x. \quad (84)$$

We now prove (84) by induction on $n \in \mathbb{N}_0$. Note that (68) and the fact that $\mathcal{O}(\mathbb{I}) = d$ demonstrate that $\mathcal{R}_a(\mathbb{I}^{\bullet 0}) \in C(\mathbb{R}^d, \mathbb{R}^d)$ and $\forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{I}^{\bullet 0}))(x) = x$. This establishes (84) in the base case $n = 0$. For the induction step observe that for all $n \in \mathbb{N}_0$ with $\mathcal{R}_a(\mathbb{I}^{\bullet n}) \in C(\mathbb{R}^d, \mathbb{R}^d)$ and $\forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{I}^{\bullet n}))(x) = x$ it holds that

$$\mathcal{R}_a(\mathbb{I}^{\bullet(n+1)}) = \mathcal{R}_a(\mathbb{I} \bullet (\mathbb{I}^{\bullet n})) = (\mathcal{R}_a(\mathbb{I})) \circ (\mathcal{R}_a(\mathbb{I}^{\bullet n})) \in C(\mathbb{R}^d, \mathbb{R}^d) \quad (85)$$

and

$$\begin{aligned} \forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{I}^{\bullet(n+1)}))(x) &= ([\mathcal{R}_a(\mathbb{I})] \circ [\mathcal{R}_a(\mathbb{I}^{\bullet n})])(x) \\ &= (\mathcal{R}_a(\mathbb{I}))((\mathcal{R}_a(\mathbb{I}^{\bullet n}))(x)) = (\mathcal{R}_a(\mathbb{I}))(x) = x. \end{aligned} \quad (86)$$

Induction thus proves (84). Next observe that (84) establishes item (i). Moreover, note that (69), item (v) in Proposition 2.6, item (i), and the fact that $\mathcal{I}(\mathbb{I}) = \mathcal{O}(\Phi)$ ensure that

$$\begin{aligned} \mathcal{R}_a(\mathcal{E}_{L,\mathbb{I}}(\Phi)) &= \mathcal{R}_a((\mathbb{I}^{\bullet(L-\mathcal{L}(\Phi))} \bullet \Phi)) \\ &\in C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{O}(\mathbb{I})}) = C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{I}(\mathbb{I})}) = C(\mathbb{R}^{\mathcal{I}(\Phi)}, \mathbb{R}^{\mathcal{O}(\Phi)}) \end{aligned} \quad (87)$$

and

$$\begin{aligned} \forall x \in \mathbb{R}^{\mathcal{I}(\Phi)}: (\mathcal{R}_a(\mathcal{E}_{L,\mathbb{I}}(\Phi)))(x) &= (\mathcal{R}_a(\mathbb{I}^{\bullet(L-\mathcal{L}(\Phi))})((\mathcal{R}_a(\Phi))(x))) \\ &= (\mathcal{R}_a(\Phi))(x). \end{aligned} \quad (88)$$

This establishes item (ii). The proof of Lemma 2.14 is thus completed. \square

2.2.5 Compositions of ANNs involving artificial identities

Definition 2.15 (Composition of ANNs involving artificial identities). *Let $\Psi \in \mathbf{N}$. Then we denote by*

$$(\cdot) \odot_{\Psi} (\cdot): \{(\Phi_1, \Phi_2) \in \mathbf{N} \times \mathbf{N}: \mathcal{I}(\Phi_1) = \mathcal{O}(\Psi) \text{ and } \mathcal{O}(\Phi_2) = \mathcal{I}(\Psi)\} \rightarrow \mathbf{N} \quad (89)$$

the function which satisfies for all $\Phi_1, \Phi_2 \in \mathbf{N}$ with $\mathcal{I}(\Phi_1) = \mathcal{O}(\Psi)$ and $\mathcal{O}(\Phi_2) = \mathcal{I}(\Psi)$ that

$$\Phi_1 \odot_{\Psi} \Phi_2 = \Phi_1 \bullet (\Psi \bullet \Phi_2) = (\Phi_1 \bullet \Psi) \bullet \Phi_2 \quad (90)$$

(cf. Definition 2.1, Definition 2.5, and Lemma 2.8).

Proposition 2.16. *Let $\Psi, \Phi_1, \Phi_2 \in \mathbf{N}$, $\mathbf{i}, l_{1,0}, l_{1,1}, \dots, l_{1,\mathcal{L}(\Phi_1)}, l_{2,0}, l_{2,1}, \dots, l_{2,\mathcal{L}(\Phi_2)} \in \mathbb{N}$ satisfy for all $k \in \{1, 2\}$ that $\mathcal{D}(\Psi) = (\mathcal{I}(\Psi), \mathbf{i}, \mathcal{O}(\Psi))$, $\mathcal{I}(\Phi_1) = \mathcal{O}(\Psi)$, $\mathcal{O}(\Phi_2) = \mathcal{I}(\Psi)$, and $\mathcal{D}(\Phi_k) = (l_{k,0}, l_{k,1}, \dots, l_{k,\mathcal{L}(\Phi_k)})$ (cf. Definition 2.1). Then*

(i) *it holds that*

$$\mathcal{D}(\Phi_1 \odot_{\Psi} \Phi_2) = (l_{2,0}, l_{2,1}, \dots, l_{2,\mathcal{L}(\Phi_2)-1}, \mathbf{i}, l_{1,1}, l_{1,2}, \dots, l_{1,\mathcal{L}(\Phi_1)}), \quad (91)$$

(ii) *it holds that*

$$\mathcal{L}(\Phi_1 \odot_{\Psi} \Phi_2) = \mathcal{L}(\Phi_1) + \mathcal{L}(\Phi_2), \quad (92)$$

(iii) *it holds that*

$$\mathcal{P}(\Phi_1 \odot_{\Psi} \Phi_2) \leq \left[\max \left\{ 1, \frac{\mathbf{i}}{\mathcal{I}(\Psi)}, \frac{\mathbf{i}}{\mathcal{O}(\Psi)} \right\} \right] (\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2)), \quad (93)$$

and

(iv) *it holds for all $a \in C(\mathbb{R}, \mathbb{R})$ that $\mathcal{R}_a(\Phi_1 \odot_{\Psi} \Phi_2) \in C(\mathbb{R}^{\mathcal{I}(\Phi_2)}, \mathbb{R}^{\mathcal{O}(\Phi_1)})$ and*

$$\mathcal{R}_a(\Phi_1 \odot_{\Psi} \Phi_2) = [\mathcal{R}_a(\Phi_1)] \circ [\mathcal{R}_a(\Psi)] \circ [\mathcal{R}_a(\Phi_2)] \quad (94)$$

(cf. Definition 2.3 and Definition 2.15).

Proof of Propositions 2.16. Throughout this proof let $a \in C(\mathbb{R}, \mathbb{R})$, $L_1, L_2 \in \mathbb{N}$ satisfy that $L_1 = \mathcal{L}(\Phi_1)$ and $L_2 = \mathcal{L}(\Phi_2)$. Note that item (i) in Proposition 2.6, the hypothesis that $\mathcal{D}(\Phi_2) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2})$, the hypothesis that $\mathcal{D}(\Psi) = (\mathcal{I}(\Psi), \mathbf{i}, \mathcal{O}(\Psi))$, and the hypothesis that $\mathcal{I}(\Psi) = \mathcal{O}(\Phi_2)$ show that

$$\mathcal{D}(\Psi \bullet \Phi_2) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, \mathbf{i}, \mathcal{O}(\Psi)) \quad (95)$$

(cf. Definition 2.5). Combining this with item (i) in Proposition 2.6, the hypothesis that $\mathcal{D}(\Phi_1) = (l_{1,0}, l_{1,1}, \dots, l_{1,L_1})$, and the hypothesis that $\mathcal{I}(\Phi_1) = \mathcal{O}(\Psi)$ proves that

$$\mathcal{D}(\Phi_1 \odot_{\Psi} \Phi_2) = \mathcal{D}(\Phi_1 \bullet (\Psi \bullet \Phi_2)) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, \mathbf{i}, l_{1,1}, l_{1,2}, \dots, l_{1,L_1}). \quad (96)$$

This establishes item (i). Moreover, observe that item (ii) in Proposition 2.6 and the fact that $\mathcal{L}(\Psi) = 2$ ensure that

$$\begin{aligned}\mathcal{L}(\Phi_1 \odot_{\Psi} \Phi_2) &= \mathcal{L}(\Phi_1 \bullet (\Psi \bullet \Phi_2)) = \mathcal{L}(\Phi_1) + \mathcal{L}(\Psi \bullet \Phi_2) - 1 \\ &= \mathcal{L}(\Phi_1) + \mathcal{L}(\Psi) + \mathcal{L}(\Phi_2) - 2 = \mathcal{L}(\Phi_1) + \mathcal{L}(\Phi_2).\end{aligned}\quad (97)$$

This establishes item (ii). In addition, observe that (96), the fact that $\mathcal{I}(\Psi) = \mathcal{O}(\Phi_2) = l_{2,L_2}$, and the fact that $\mathcal{O}(\Psi) = \mathcal{I}(\Phi_1) = l_{1,0}$ demonstrate that

$$\begin{aligned}\mathcal{P}(\Phi_1 \odot_{\Psi} \Phi_2) &= \left[\sum_{m=1}^{L_2-1} l_{2,m}(l_{2,m-1} + 1) \right] + \left[\sum_{m=2}^{L_1} l_{1,m}(l_{1,m-1} + 1) \right] \\ &\quad + \mathbf{i}(l_{2,L_2-1} + 1) + l_{1,1}(\mathbf{i} + 1) \\ &= \left[\sum_{m=1}^{L_2-1} l_{2,m}(l_{2,m-1} + 1) \right] + \left[\sum_{m=2}^{L_1} l_{1,m}(l_{1,m-1} + 1) \right] \\ &\quad + \frac{\mathbf{i}}{\mathcal{I}(\Psi)} l_{2,L_2}(l_{2,L_2-1} + 1) + l_{1,1} \left(\frac{\mathbf{i}}{\mathcal{O}(\Psi)} l_{1,0} + 1 \right) \\ &\leq \left[\max \left\{ 1, \frac{\mathbf{i}}{\mathcal{I}(\Psi)} \right\} \right] \left[\sum_{m=1}^{L_2} l_{2,m}(l_{2,m-1} + 1) \right] \\ &\quad + \left[\max \left\{ 1, \frac{\mathbf{i}}{\mathcal{O}(\Psi)} \right\} \right] \left[\sum_{m=1}^{L_1} l_{1,m}(l_{1,m-1} + 1) \right] \\ &\leq \left[\max \left\{ 1, \frac{\mathbf{i}}{\mathcal{I}(\Psi)}, \frac{\mathbf{i}}{\mathcal{O}(\Psi)} \right\} \right] (\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2)).\end{aligned}\quad (98)$$

This establishes item (iii). Next note that item (v) in Proposition 2.6 implies that

$$\begin{aligned}\mathcal{R}_a(\Phi_1 \odot_{\Psi} \Phi_2) &= \mathcal{R}_a(\Phi_1 \bullet (\Psi \bullet \Phi_2)) \\ &= [\mathcal{R}_a(\Phi_1)] \circ [\mathcal{R}_a(\Psi \bullet \Phi_2)] \\ &= ([\mathcal{R}_a(\Phi_1)] \circ [\mathcal{R}_a(\Psi)] \circ [\mathcal{R}_a(\Phi_2)]) \in C(\mathbb{R}^{\mathcal{I}(\Phi_2)}, \mathbb{R}^{\mathcal{O}(\Phi_1)}).\end{aligned}\quad (99)$$

This establishes item (iv). The proof of Proposition 2.16 is thus completed. \square

2.3 Parallelizations of ANNs

2.3.1 Parallelizations of ANNs with the same length

Definition 2.17 (Parallelization of ANNs with the same length). *Let $n \in \mathbb{N}$. Then we denote by*

$$\mathbf{P}_n: \{(\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n: \mathcal{L}(\Phi_1) = \mathcal{L}(\Phi_2) = \dots = \mathcal{L}(\Phi_n)\} \rightarrow \mathbf{N} \quad (100)$$

the function which satisfies for all $L \in \mathbb{N}$, $(l_{1,0}, l_{1,1}, \dots, l_{1,L}), (l_{2,0}, l_{2,1}, \dots, l_{2,L}), \dots, (l_{n,0}, l_{n,1}, \dots, l_{n,L}) \in \mathbb{N}^{L+1}$, $\Phi_1 = ((W_{1,1}, B_{1,1}), (W_{1,2}, B_{1,2}), \dots, (W_{1,L}, B_{1,L})) \in (\times_{k=1}^L$

$(\mathbb{R}^{l_{1,k} \times l_{1,k-1}} \times \mathbb{R}^{l_{1,k}})$, $\Phi_2 = ((W_{2,1}, B_{2,1}), (W_{2,2}, B_{2,2}), \dots, (W_{2,L}, B_{2,L})) \in (\times_{k=1}^L$
 $(\mathbb{R}^{l_{2,k} \times l_{2,k-1}} \times \mathbb{R}^{l_{2,k}}))$, \dots , $\Phi_n = ((W_{n,1}, B_{n,1}), (W_{n,2}, B_{n,2}), \dots, (W_{n,L}, B_{n,L})) \in (\times_{k=1}^L$
 $(\mathbb{R}^{l_{n,k} \times l_{n,k-1}} \times \mathbb{R}^{l_{n,k}}))$ that

$$\mathbf{P}_n(\Phi_1, \Phi_2, \dots, \Phi_n) = \left(\left(\left(\begin{array}{ccccc} W_{1,1} & 0 & 0 & \cdots & 0 \\ 0 & W_{2,1} & 0 & \cdots & 0 \\ 0 & 0 & W_{3,1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & W_{n,1} \end{array} \right), \left(\begin{array}{c} B_{1,1} \\ B_{2,1} \\ B_{3,1} \\ \vdots \\ B_{n,1} \end{array} \right) \right), \right. \\ \left(\left(\begin{array}{ccccc} W_{1,2} & 0 & 0 & \cdots & 0 \\ 0 & W_{2,2} & 0 & \cdots & 0 \\ 0 & 0 & W_{3,2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & W_{n,2} \end{array} \right), \left(\begin{array}{c} B_{1,2} \\ B_{2,2} \\ B_{3,2} \\ \vdots \\ B_{n,2} \end{array} \right) \right), \dots, \\ \left(\left(\begin{array}{ccccc} W_{1,L} & 0 & 0 & \cdots & 0 \\ 0 & W_{2,L} & 0 & \cdots & 0 \\ 0 & 0 & W_{3,L} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & W_{n,L} \end{array} \right), \left(\begin{array}{c} B_{1,L} \\ B_{2,L} \\ B_{3,L} \\ \vdots \\ B_{n,L} \end{array} \right) \right) \right) \quad (101)$$

(cf. Definition 2.1).

Lemma 2.18. Let $n, L \in \mathbb{N}$, $(l_{1,0}, l_{1,1}, \dots, l_{1,L}), (l_{2,0}, l_{2,1}, \dots, l_{2,L}), \dots, (l_{n,0}, l_{n,1}, \dots, l_{n,L}) \in \mathbb{N}^{L+1}$, $\Phi_1 = ((W_{1,1}, B_{1,1}), (W_{1,2}, B_{1,2}), \dots, (W_{1,L}, B_{1,L})) \in (\times_{k=1}^L (\mathbb{R}^{l_{1,k} \times l_{1,k-1}} \times \mathbb{R}^{l_{1,k}}))$,
 $\Phi_2 = ((W_{2,1}, B_{2,1}), (W_{2,2}, B_{2,2}), \dots, (W_{2,L}, B_{2,L})) \in (\times_{k=1}^L (\mathbb{R}^{l_{2,k} \times l_{2,k-1}} \times \mathbb{R}^{l_{2,k}}))$, \dots ,
 $\Phi_n = ((W_{n,1}, B_{n,1}), (W_{n,2}, B_{n,2}), \dots, (W_{n,L}, B_{n,L})) \in (\times_{k=1}^L (\mathbb{R}^{l_{n,k} \times l_{n,k-1}} \times \mathbb{R}^{l_{n,k}}))$.
Then it holds that

$$\mathbf{P}_n(\Phi_1, \Phi_2, \dots, \Phi_n) \in \left(\times_{k=1}^L (\mathbb{R}^{(\sum_{j=1}^n l_{j,k}) \times (\sum_{j=1}^n l_{j,k-1})} \times \mathbb{R}^{\sum_{j=1}^n l_{j,k}}) \right) \quad (102)$$

(cf. Definition 2.17).

Proof of Lemma 2.18. Note that (101) establishes (102). The proof of Lemma 2.18 is thus completed. \square

Proposition 2.19. Let $a \in C(\mathbb{R}, \mathbb{R})$, $n \in \mathbb{N}$, $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbb{N}^n$ satisfy that $\mathcal{L}(\Phi_1) = \mathcal{L}(\Phi_2) = \dots = \mathcal{L}(\Phi_n)$ (cf. Definition 2.1). Then

(i) it holds that

$$\mathcal{R}_a(\mathbf{P}_n(\Phi)) \in C(\mathbb{R}^{[\sum_{j=1}^n \mathcal{I}(\Phi_j)]}, \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]}) \quad (103)$$

and

(ii) it holds for all $x_1 \in \mathbb{R}^{\mathcal{I}(\Phi_1)}$, $x_2 \in \mathbb{R}^{\mathcal{I}(\Phi_2)}$, \dots , $x_n \in \mathbb{R}^{\mathcal{I}(\Phi_n)}$ that

$$\begin{aligned} & (\mathcal{R}_a(\mathbf{P}_n(\Phi)))(x_1, x_2, \dots, x_n) \\ &= ((\mathcal{R}_a(\Phi_1))(x_1), (\mathcal{R}_a(\Phi_2))(x_2), \dots, (\mathcal{R}_a(\Phi_n))(x_n)) \in \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]} \end{aligned} \quad (104)$$

(cf. Definition 2.3 and Definition 2.17).

Proof of Proposition 2.19. Throughout this proof let $L \in \mathbb{N}$ satisfy that $L = \mathcal{L}(\Phi_1)$, let $l_{j,0}, l_{j,1}, \dots, l_{j,L} \in \mathbb{N}$, $j \in \{1, 2, \dots, n\}$, satisfy for all $j \in \{1, 2, \dots, n\}$ that $\mathcal{D}(\Phi_j) = (l_{j,0}, l_{j,1}, \dots, l_{j,L})$, let $((W_{j,1}, B_{j,1}), (W_{j,2}, B_{j,2}), \dots, (W_{j,L}, B_{j,L})) \in (\times_{k=1}^L (\mathbb{R}^{l_{j,k} \times l_{j,k-1}} \times \mathbb{R}^{l_{j,k}}))$, $j \in \{1, 2, \dots, n\}$, satisfy for all $j \in \{1, 2, \dots, n\}$ that

$$\Phi_j = ((W_{j,1}, B_{j,1}), (W_{j,2}, B_{j,2}), \dots, (W_{j,L}, B_{j,L})), \quad (105)$$

let $\alpha_k \in \mathbb{N}$, $k \in \{0, 1, \dots, L\}$, satisfy for all $k \in \{0, 1, \dots, L\}$ that $\alpha_k = \sum_{j=1}^n l_{j,k}$, let $((A_1, b_1), (A_2, b_2), \dots, (A_L, b_L)) \in (\times_{k=1}^L (\mathbb{R}^{\alpha_k \times \alpha_{k-1}} \times \mathbb{R}^{\alpha_k}))$ satisfy that

$$\mathbf{P}_n(\Phi) = ((A_1, b_1), (A_2, b_2), \dots, (A_L, b_L)) \quad (106)$$

(cf. Lemma 2.18), let $(x_{j,0}, x_{j,1}, \dots, x_{j,L-1}) \in (\mathbb{R}^{l_{j,0}} \times \mathbb{R}^{l_{j,1}} \times \dots \times \mathbb{R}^{l_{j,L-1}})$, $j \in \{1, 2, \dots, n\}$, satisfy for all $j \in \{1, 2, \dots, n\}$, $k \in \mathbb{N} \cap (0, L)$ that

$$x_{j,k} = \mathfrak{M}_{a,l_{j,k}}(W_{j,k}x_{j,k-1} + B_{j,k}) \quad (107)$$

(cf. Definition 2.2), and let $\mathfrak{r}_0 \in \mathbb{R}^{\alpha_0}$, $\mathfrak{r}_1 \in \mathbb{R}^{\alpha_1}$, \dots , $\mathfrak{r}_{L-1} \in \mathbb{R}^{\alpha_{L-1}}$ satisfy for all $k \in \{0, 1, \dots, L-1\}$ that $\mathfrak{r}_k = (x_{1,k}, x_{2,k}, \dots, x_{n,k})$. Observe that (106) demonstrates that $\mathcal{I}(\mathbf{P}_n(\Phi)) = \alpha_0$ and $\mathcal{O}(\mathbf{P}_n(\Phi)) = \alpha_L$. Combining this with item (ii) in Lemma 2.4, the fact that for all $k \in \{0, 1, \dots, L\}$ it holds that $\alpha_k = \sum_{j=1}^n l_{j,k}$, the fact that for all $j \in \{1, 2, \dots, n\}$ it holds that $\mathcal{I}(\Phi_j) = l_{j,0}$, and the fact that for all $j \in \{1, 2, \dots, n\}$ it holds that $\mathcal{O}(\Phi_j) = l_{j,L}$ ensures that

$$\begin{aligned} \mathcal{R}_a(\mathbf{P}_n(\Phi)) &\in C(\mathbb{R}^{\alpha_0}, \mathbb{R}^{\alpha_L}) = C(\mathbb{R}^{[\sum_{j=1}^n l_{j,0}]}, \mathbb{R}^{[\sum_{j=1}^n l_{j,L}]}) \\ &= C(\mathbb{R}^{[\sum_{j=1}^n \mathcal{I}(\Phi_j)]}, \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]}). \end{aligned} \quad (108)$$

This proves item (i). Moreover, observe that (101) and (106) demonstrate that for all $k \in \{1, 2, \dots, L\}$ it holds that

$$A_k = \begin{pmatrix} W_{1,k} & 0 & 0 & \cdots & 0 \\ 0 & W_{2,k} & 0 & \cdots & 0 \\ 0 & 0 & W_{3,k} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & W_{n,k} \end{pmatrix} \quad \text{and} \quad b_k = \begin{pmatrix} B_{1,k} \\ B_{2,k} \\ B_{3,k} \\ \vdots \\ B_{n,k} \end{pmatrix}. \quad (109)$$

Combining this with (6), (107), and the fact that for all $k \in \mathbb{N} \cap [0, L)$ it holds that $\mathbf{x}_k = (x_{1,k}, x_{2,k}, \dots, x_{n,k})$ implies that for all $k \in \mathbb{N} \cap (0, L)$ it holds that

$$\mathfrak{M}_{a,\alpha_k}(A_k \mathbf{x}_{k-1} + b_k) = \begin{pmatrix} \mathfrak{M}_{a,l_{1,k}}(W_{1,k}x_{1,k-1} + B_{1,k}) \\ \mathfrak{M}_{a,l_{2,k}}(W_{2,k}x_{2,k-1} + B_{2,k}) \\ \vdots \\ \mathfrak{M}_{a,l_{n,k}}(W_{n,k}x_{n,k-1} + B_{n,k}) \end{pmatrix} = \begin{pmatrix} x_{1,k} \\ x_{2,k} \\ \vdots \\ x_{n,k} \end{pmatrix} = \mathbf{x}_k. \quad (110)$$

This, (7), (105), (106), (107), (109), the fact that $\mathbf{x}_0 = (x_{1,0}, x_{2,0}, \dots, x_{n,0})$, and the fact that $\mathbf{x}_{L-1} = (x_{1,L-1}, x_{2,L-1}, \dots, x_{n,L-1})$ ensure that

$$\begin{aligned} (\mathcal{R}_a(\mathbf{P}_n(\Phi)))(x_{1,0}, x_{2,0}, \dots, x_{n,0}) &= (\mathcal{R}_a(\mathbf{P}_n(\Phi)))(\mathbf{x}_0) \\ &= A_L \mathbf{x}_{L-1} + b_L = \begin{pmatrix} W_{1,L}x_{1,L-1} + B_{1,L} \\ W_{2,L}x_{2,L-1} + B_{2,L} \\ \vdots \\ W_{n,L}x_{n,L-1} + B_{n,L} \end{pmatrix} = \begin{pmatrix} (\mathcal{R}_a(\Phi_1))(x_{1,0}) \\ (\mathcal{R}_a(\Phi_2))(x_{2,0}) \\ \vdots \\ (\mathcal{R}_a(\Phi_n))(x_{n,0}) \end{pmatrix}. \end{aligned} \quad (111)$$

This establishes item (ii). The proof of Proposition 2.19 is thus completed. \square

Proposition 2.20. *Let $n, L \in \mathbb{N}$, $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n$, $(l_{1,0}, l_{1,1}, \dots, l_{1,L})$, $(l_{2,0}, l_{2,1}, \dots, l_{2,L}), \dots, (l_{n,0}, l_{n,1}, \dots, l_{n,L}) \in \mathbb{N}^{L+1}$ satisfy for all $j \in \{1, 2, \dots, n\}$ that $\mathcal{D}(\Phi_j) = (l_{j,0}, l_{j,1}, \dots, l_{j,L})$ (cf. Definition 2.1). Then*

(i) *it holds that*

$$\mathcal{D}(\mathbf{P}_n(\Phi)) = (\sum_{j=1}^n l_{j,0}, \sum_{j=1}^n l_{j,1}, \dots, \sum_{j=1}^n l_{j,L}) \quad (112)$$

and

(ii) *it holds that*

$$\mathcal{P}(\mathbf{P}_n(\Phi)) \leq \frac{1}{2} [\sum_{j=1}^n \mathcal{P}(\Phi_j)]^2 \quad (113)$$

(cf. Definition 2.17).

Proof of Proposition 2.20. Note that the hypothesis that $\forall j \in \{1, 2, \dots, n\}: \mathcal{D}(\Phi_j) = (l_{j,0}, l_{j,1}, \dots, l_{j,L})$ and Lemma 2.18 assure that

$$\mathcal{D}(\mathbf{P}_n(\Phi)) = (\sum_{j=1}^n l_{j,0}, \sum_{j=1}^n l_{j,1}, \dots, \sum_{j=1}^n l_{j,L}). \quad (114)$$

This establishes item (i). Moreover, observe that (114) demonstrates that

$$\begin{aligned}
\mathcal{P}(\mathbf{P}_n(\Phi)) &= \sum_{k=1}^L \left[\sum_{i=1}^n l_{i,k} \right] \left[\left(\sum_{i=1}^n l_{i,k-1} \right) + 1 \right] \\
&= \sum_{k=1}^L \left[\sum_{i=1}^n l_{i,k} \right] \left[\left(\sum_{j=1}^n l_{j,k-1} \right) + 1 \right] \\
&\leq \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^L l_{i,k} (l_{j,k-1} + 1) \leq \sum_{i=1}^n \sum_{j=1}^n \sum_{k,\ell=1}^L l_{i,k} (l_{j,\ell-1} + 1) \\
&= \sum_{i=1}^n \sum_{j=1}^n \left[\sum_{k=1}^L l_{i,k} \right] \left[\sum_{\ell=1}^L (l_{j,\ell-1} + 1) \right] \\
&\leq \sum_{i=1}^n \sum_{j=1}^n \left[\sum_{k=1}^L \frac{1}{2} l_{i,k} (l_{i,k-1} + 1) \right] \left[\sum_{\ell=1}^L l_{j,\ell} (l_{j,\ell-1} + 1) \right] \\
&= \sum_{i=1}^n \sum_{j=1}^n \frac{1}{2} \mathcal{P}(\Phi_i) \mathcal{P}(\Phi_j) = \frac{1}{2} \left[\sum_{i=1}^n \mathcal{P}(\Phi_i) \right]^2.
\end{aligned} \tag{115}$$

The proof of Proposition 2.20 is thus completed. \square

Corollary 2.21. *Let $n \in \mathbb{N}$, $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n$ satisfy that $\mathcal{D}(\Phi_1) = \mathcal{D}(\Phi_2) = \dots = \mathcal{D}(\Phi_n)$ (cf. Definition 2.1). Then it holds that $\mathcal{P}(\mathbf{P}_n(\Phi)) \leq n^2 \mathcal{P}(\Phi_1)$ (cf. Definition 2.17).*

Proof of Corollary 2.21. Throughout this proof let $L \in \mathbb{N}$, $l_0, l_1, \dots, l_L \in \mathbb{N}$ satisfy that $\mathcal{D}(\Phi_1) = (l_0, l_1, \dots, l_L)$. Note that item (i) in Proposition 2.20 and the fact that $\forall j \in \{1, 2, \dots, n\}$: $\mathcal{D}(\Phi_j) = (l_0, l_1, \dots, l_L)$ demonstrate that

$$\begin{aligned}
\mathcal{P}(\mathbf{P}_n(\Phi_1, \Phi_2, \dots, \Phi_n)) &= \sum_{j=1}^L (nl_j) ((nl_{j-1}) + 1) \leq \sum_{j=1}^L (nl_j) ((nl_{j-1}) + n) \\
&= n^2 \left[\sum_{j=1}^L l_j (l_{j-1} + 1) \right] = n^2 \mathcal{P}(\Phi_1).
\end{aligned} \tag{116}$$

The proof of Corollary 2.21 is thus completed. \square

2.3.2 Parallelizations of ANNs with different lengths

Definition 2.22 (Parallelization of ANNs with different length). *Let $n \in \mathbb{N}$, $\Psi = (\Psi_1, \Psi_2, \dots, \Psi_n) \in \mathbf{N}^n$ satisfy for all $j \in \{1, 2, \dots, n\}$ that $\mathcal{H}(\Psi_j) = 1$ and $\mathcal{I}(\Psi_j) = \mathcal{O}(\Psi_j)$. Then we denote by*

$$\mathbf{P}_{n,\Psi}: \{(\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n: (\forall j \in \{1, 2, \dots, n\}: \mathcal{O}(\Phi_j) = \mathcal{I}(\Psi_j))\} \rightarrow \mathbf{N} \tag{117}$$

the function which satisfies for all $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n$ with $\forall j \in \{1, 2, \dots, n\}$: $\mathcal{O}(\Phi_j) = \mathcal{I}(\Psi_j)$ that

$$\mathbf{P}_{n, \Psi}(\Phi) = \mathbf{P}_n(\mathcal{E}_{\max_{k \in \{1, 2, \dots, n\}} \mathcal{L}(\Phi_k), \Psi_1}(\Phi_1), \dots, \mathcal{E}_{\max_{k \in \{1, 2, \dots, n\}} \mathcal{L}(\Phi_k), \Psi_n}(\Phi_n)) \quad (118)$$

(cf. Definition 2.1, Definition 2.12, Lemma 2.13, and Definition 2.17).

Corollary 2.23. Let $a \in C(\mathbb{R}, \mathbb{R})$, $n \in \mathbb{N}$, $\mathbb{I} = (\mathbb{I}_1, \mathbb{I}_2, \dots, \mathbb{I}_n)$, $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n$ satisfy for all $j \in \{1, 2, \dots, n\}$, $x \in \mathbb{R}^{\mathcal{O}(\Phi_j)}$ that $\mathcal{H}(\mathbb{I}_j) = 1$, $\mathcal{I}(\mathbb{I}_j) = \mathcal{O}(\mathbb{I}_j) = \mathcal{O}(\Phi_j)$, and $(\mathcal{R}_a(\mathbb{I}_j))(x) = x$ (cf. Definition 2.1 and Definition 2.3). Then

(i) it holds that

$$\mathcal{R}_a(\mathbf{P}_{n, \mathbb{I}}(\Phi)) \in C(\mathbb{R}^{[\sum_{j=1}^n \mathcal{I}(\Phi_j)]}, \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]}) \quad (119)$$

and

(ii) it holds for all $x_1 \in \mathbb{R}^{\mathcal{I}(\Phi_1)}$, $x_2 \in \mathbb{R}^{\mathcal{I}(\Phi_2)}$, \dots , $x_n \in \mathbb{R}^{\mathcal{I}(\Phi_n)}$ that

$$\begin{aligned} & (\mathcal{R}_a(\mathbf{P}_{n, \mathbb{I}}(\Phi)))(x_1, x_2, \dots, x_n) \\ &= ((\mathcal{R}_a(\Phi_1))(x_1), (\mathcal{R}_a(\Phi_2))(x_2), \dots, (\mathcal{R}_a(\Phi_n))(x_n)) \in \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]} \end{aligned} \quad (120)$$

(cf. Definition 2.22).

Proof of Corollary 2.23. Throughout this proof let $L \in \mathbb{N}$ satisfy that $L = \max_{j \in \{1, 2, \dots, n\}} \mathcal{L}(\Phi_j)$. Note that item (ii) in Lemma 2.13, the hypothesis that for all $j \in \{1, 2, \dots, n\}$ it holds that $\mathcal{H}(\mathbb{I}_j) = 1$, (69), (13), and item (ii) in Lemma 2.14 demonstrate

(I) that for all $j \in \{1, 2, \dots, n\}$ it holds that $\mathcal{L}(\mathcal{E}_{L, \mathbb{I}_j}(\Phi_j)) = L$ and $\mathcal{R}_a(\mathcal{E}_{L, \mathbb{I}_j}(\Phi_j)) \in C(\mathbb{R}^{\mathcal{I}(\Phi_j)}, \mathbb{R}^{\mathcal{O}(\Phi_j)})$ and

(II) that for all $j \in \{1, 2, \dots, n\}$, $x \in \mathbb{R}^{\mathcal{I}(\Phi_j)}$ it holds that

$$(\mathcal{R}_a(\mathcal{E}_{L, \mathbb{I}_j}(\Phi_j)))(x) = (\mathcal{R}_a(\Phi_j))(x) \quad (121)$$

(cf. Definition 2.12). Items (i)–(ii) in Proposition 2.19 therefore imply

(A) that

$$\mathcal{R}_a(\mathbf{P}_n(\mathcal{E}_{L, \mathbb{I}_1}(\Phi_1), \mathcal{E}_{L, \mathbb{I}_2}(\Phi_2), \dots, \mathcal{E}_{L, \mathbb{I}_n}(\Phi_n))) \in C(\mathbb{R}^{[\sum_{j=1}^n \mathcal{I}(\Phi_j)]}, \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]}) \quad (122)$$

and

(B) that for all $x_1 \in \mathbb{R}^{\mathcal{I}(\Phi_1)}, x_2 \in \mathbb{R}^{\mathcal{I}(\Phi_2)}, \dots, x_n \in \mathbb{R}^{\mathcal{I}(\Phi_n)}$ it holds that

$$\begin{aligned}
& (\mathcal{R}_a(\mathbf{P}_n(\mathcal{E}_{L, \mathbb{I}_1}(\Phi_1), \mathcal{E}_{L, \mathbb{I}_2}(\Phi_2), \dots, \mathcal{E}_{L, \mathbb{I}_n}(\Phi_n))))(x_1, x_2, \dots, x_n) \\
&= \left((\mathcal{R}_a(\mathcal{E}_{L, \mathbb{I}_1}(\Phi_1)))(x_1), (\mathcal{R}_a(\mathcal{E}_{L, \mathbb{I}_2}(\Phi_2)))(x_2), \dots, (\mathcal{R}_a(\mathcal{E}_{L, \mathbb{I}_n}(\Phi_n)))(x_n) \right) \\
&= \left((\mathcal{R}_a(\Phi_1))(x_1), (\mathcal{R}_a(\Phi_2))(x_2), \dots, (\mathcal{R}_a(\Phi_n))(x_n) \right)
\end{aligned} \tag{123}$$

(cf. Definition 2.17). Combining this with (118) and the fact that $L = \max_{j \in \{1, 2, \dots, n\}} \mathcal{L}(\Phi_j)$ ensures

(C) that

$$\mathcal{R}_a(\mathbf{P}_{n, \mathbb{I}}(\Phi)) \in C(\mathbb{R}^{[\sum_{j=1}^n \mathcal{I}(\Phi_j)]}, \mathbb{R}^{[\sum_{j=1}^n \mathcal{O}(\Phi_j)]}) \tag{124}$$

and

(D) that for all $x_1 \in \mathbb{R}^{\mathcal{I}(\Phi_1)}, x_2 \in \mathbb{R}^{\mathcal{I}(\Phi_2)}, \dots, x_n \in \mathbb{R}^{\mathcal{I}(\Phi_n)}$ it holds that

$$\begin{aligned}
& (\mathcal{R}_a(\mathbf{P}_{n, \mathbb{I}}(\Phi)))(x_1, x_2, \dots, x_n) \\
&= (\mathcal{R}_a(\mathbf{P}_n(\mathcal{E}_{L, \mathbb{I}_1}(\Phi_1), \mathcal{E}_{L, \mathbb{I}_2}(\Phi_2), \dots, \mathcal{E}_{L, \mathbb{I}_n}(\Phi_n))))(x_1, x_2, \dots, x_n) \\
&= \left((\mathcal{R}_a(\Phi_1))(x_1), (\mathcal{R}_a(\Phi_2))(x_2), \dots, (\mathcal{R}_a(\Phi_n))(x_n) \right).
\end{aligned} \tag{125}$$

This establishes items (i)–(ii). The proof of Corollary 2.23 is thus completed. \square

Corollary 2.24. *Let $n, L \in \mathbb{N}$, $\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_n \in \mathbb{N}$, $\Psi = (\Psi_1, \Psi_2, \dots, \Psi_n)$, $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n) \in \mathbf{N}^n$ satisfy for all $j \in \{1, 2, \dots, n\}$ that $\mathcal{D}(\Psi_j) = (\mathcal{O}(\Phi_j), \mathbf{i}_j, \mathcal{O}(\Phi_j))$ and $L = \max_{k \in \{1, 2, \dots, n\}} \mathcal{L}(\Phi_k)$ (cf. Definition 2.1). Then it holds that*

$$\begin{aligned}
& \mathcal{P}(\mathbf{P}_{n, \Psi}(\Phi)) \\
& \leq \frac{1}{2} \left(\left[\sum_{j=1}^n \left[\max \left\{ 1, \frac{\mathbf{i}_j}{\mathcal{O}(\Phi_j)} \right\} \right] \mathcal{P}(\Phi_j) \mathbf{1}_{(\mathcal{L}(\Phi_j), \infty)}(L) \right] \right. \\
& \quad \left. + \left[\sum_{j=1}^n \left((L - \mathcal{L}(\Phi_j) - 1) \mathbf{i}_j (\mathbf{i}_j + 1) + \mathcal{O}(\Phi_j) (\mathbf{i}_j + 1) \right) \mathbf{1}_{(\mathcal{L}(\Phi_j), \infty)}(L) \right] \right. \\
& \quad \left. + \left[\sum_{j=1}^n \mathcal{P}(\Phi_j) \mathbf{1}_{\{\mathcal{L}(\Phi_j)\}}(L) \right] \right)^2
\end{aligned} \tag{126}$$

(cf. Definition 2.22).

Proof of Corollary 2.24. Observe that (118), item (ii) in Proposition 2.20, and item (ii) in Lemma 2.13 assure that

$$\begin{aligned}
& \mathcal{P}(\mathbf{P}_{n,\Psi}(\Phi)) \\
&= \mathcal{P}(\mathbf{P}_n(\mathcal{E}_{L,\Psi_1}(\Phi_1), \mathcal{E}_{L,\Psi_2}(\Phi_2), \dots, \mathcal{E}_{L,\Psi_n}(\Phi_n))) \\
&\leq \frac{1}{2} \left[\sum_{j=1}^n \mathcal{P}(\mathcal{E}_{L,\Psi_j}(\Phi_j)) \right]^2 \\
&\leq \frac{1}{2} \left(\left[\sum_{j=1}^n \left[\max \left\{ 1, \frac{i_j}{\mathcal{O}(\Phi_j)} \right\} \right] \mathcal{P}(\Phi_j) \mathbf{1}_{(\mathcal{L}(\Phi_j), \infty)}(L) \right] \right. \\
&\quad \left. + \left[\sum_{j=1}^n \left((L - \mathcal{L}(\Phi_j) - 1) i_j (i_j + 1) + \mathcal{O}(\Phi_j) (i_j + 1) \right) \mathbf{1}_{(\mathcal{L}(\Phi_j), \infty)}(L) \right] \right. \\
&\quad \left. + \left[\sum_{j=1}^n \mathcal{P}(\Phi_j) \mathbf{1}_{\{\mathcal{L}(\Phi_j)\}}(L) \right] \right)^2
\end{aligned} \tag{127}$$

(cf. Definition 2.12 and Definition 2.17). The proof of Corollary 2.24 is thus completed. \square

2.4 Sums of ANNs

2.4.1 Sums of ANNs with the same length

Proposition 2.25. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $M \in \mathbb{N}$, $h_1, h_2, \dots, h_M \in \mathbb{R}$, $\Phi_1, \Phi_2, \dots, \Phi_M \in \mathbf{N}$ satisfy that $\mathcal{D}(\Phi_1) = \mathcal{D}(\Phi_2) = \dots = \mathcal{D}(\Phi_M)$ (cf. Definition 2.1). Then there exists $\Psi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{P}(\Psi) \leq M^2 \mathcal{P}(\Phi_1)$,*

(ii) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^{\mathcal{I}(\Phi_1)}, \mathbb{R}^{\mathcal{O}(\Phi_1)})$, and*

(iii) *it holds for all $x \in \mathbb{R}^{\mathcal{I}(\Phi_1)}$ that*

$$(\mathcal{R}_a(\Psi))(x) = \sum_{m=1}^M h_m (\mathcal{R}_a(\Phi_m))(x) \tag{128}$$

(cf. Definition 2.3).

Proof of Proposition 2.25. Throughout this proof let $d, \mathfrak{d} \in \mathbb{N}$ satisfy that $\mathcal{I}(\Phi_1) = d$ and $\mathcal{O}(\Phi_1) = \mathfrak{d}$, let $(A_1, b_1) \in \mathbb{R}^{\mathfrak{d} \times (M\mathfrak{d})} \times \mathbb{R}^{\mathfrak{d}}$, $(A_2, b_2) \in \mathbb{R}^{(M\mathfrak{d}) \times d} \times \mathbb{R}^{M\mathfrak{d}}$ satisfy that

$$A_1 = (h_1 \mathbf{I}_{\mathfrak{d}} \quad h_2 \mathbf{I}_{\mathfrak{d}} \quad \dots \quad h_M \mathbf{I}_{\mathfrak{d}}), \quad A_2 = \begin{pmatrix} \mathbf{I}_d \\ \mathbf{I}_d \\ \vdots \\ \mathbf{I}_d \end{pmatrix}, \quad b_1 = 0, \quad \text{and} \quad b_2 = 0 \tag{129}$$

(cf. Definition 2.10), let $\mathbb{A}_1, \mathbb{A}_2 \in \mathbf{N}$ satisfy that $\mathbb{A}_1 = (A_1, b_1)$ and $\mathbb{A}_2 = (A_2, b_2)$, and let $\Psi \in \mathbf{N}$ satisfy that

$$\Psi = \mathbb{A}_1 \bullet [\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)] \bullet \mathbb{A}_2 \quad (130)$$

(cf. Definition 2.5, Definition 2.17, Lemma 2.8, and Proposition 2.19). Note that (130) and items (i)–(ii) in Corollary 2.9 demonstrate that

$$\begin{aligned} \mathcal{P}(\Psi) &\leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{\mathcal{O}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))} \right\} \right] \mathcal{P}([\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)] \bullet \mathbb{A}_2) \\ &\leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{\mathcal{O}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))} \right\} \right] \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{\mathcal{I}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))+1} \right\} \right] \\ &\quad \cdot \mathcal{P}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)) \\ &= \left[\max \left\{ 1, \frac{\mathfrak{d}}{M\mathfrak{d}} \right\} \right] \left[\max \left\{ 1, \frac{d+1}{Md+1} \right\} \right] \mathcal{P}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)) \\ &= \mathcal{P}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)). \end{aligned} \quad (131)$$

Corollary 2.21 and the hypothesis that for all $m \in \{1, 2, \dots, M\}$ it holds that $\mathcal{D}(\Phi_m) = \mathcal{D}(\Phi_1)$ hence prove that

$$\mathcal{P}(\Psi) \leq \mathcal{P}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)) \leq M^2 \mathcal{P}(\Phi_1). \quad (132)$$

Next note that (129) and the fact that $\mathbb{A}_2 = (A_2, b_2)$ prove that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a(\mathbb{A}_2) \in C(\mathbb{R}^d, \mathbb{R}^{M^d})$ and $(\mathcal{R}_a(\mathbb{A}_2))(x) = (x, x, \dots, x) \in \mathbb{R}^{M^d}$. Proposition 2.19 and item (v) in Proposition 2.6 therefore ensure that

$$\begin{aligned} \mathcal{R}_a((\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)) \bullet \mathbb{A}_2) &= (\mathcal{R}_a(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))) \circ (\mathcal{R}_a(\mathbb{A}_2)) \\ &\in C(\mathbb{R}^{\mathcal{I}(\mathbb{A}_2)}, \mathbb{R}^{\mathcal{O}(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))}) \\ &= C(\mathbb{R}^d, \mathbb{R}^{\mathcal{O}(\Phi_1)+\mathcal{O}(\Phi_2)+\dots+\mathcal{O}(\Phi_M)}) \\ &= C(\mathbb{R}^d, \mathbb{R}^{M\mathfrak{d}}) \end{aligned} \quad (133)$$

and

$$\begin{aligned} \forall x \in \mathbb{R}^d: (\mathcal{R}_a((\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)) \bullet \mathbb{A}_2))(x) &= ([\mathcal{R}_a(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M))] \circ [\mathcal{R}_a(\mathbb{A}_2)])(x) \\ &= (\mathcal{R}_a(\mathbf{P}_M(\Phi_1, \Phi_2, \dots, \Phi_M)))(x, x, \dots, x) \\ &= ((\mathcal{R}_a(\Phi_1))(x), (\mathcal{R}_a(\Phi_2))(x), \dots, (\mathcal{R}_a(\Phi_M))(x)). \end{aligned} \quad (134)$$

Furthermore, observe that (129) and the fact that $\mathbb{A}_1 = (A_1, b_1)$ assure that for all $y_1, y_2, \dots, y_M \in \mathbb{R}^{\mathfrak{d}}$ it holds that $\mathcal{R}_a(\mathbb{A}_1) \in C(\mathbb{R}^{M\mathfrak{d}}, \mathbb{R}^{\mathfrak{d}})$ and

$$(\mathcal{R}_a(\mathbb{A}_1))(y_1, y_2, \dots, y_M) = \sum_{m=1}^M h_m y_m. \quad (135)$$

Combining this and item (v) in Proposition 2.6 with (130), (133), and (134) demonstrates that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^{\mathfrak{d}})$ and

$$(\mathcal{R}_a(\Psi))(x) = \sum_{m=1}^M h_m (\mathcal{R}_a(\Phi_m))(x). \quad (136)$$

This and (132) establish items (i)–(iii). The proof of Proposition 2.25 is thus completed. \square

2.4.2 Sums of ANNs with different lengths

Proposition 2.26. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $M, d, \mathfrak{d}, \mathbf{i}, L \in \mathbb{N}$, $h_1, h_2, \dots, h_M \in \mathbb{R}$, $\mathbb{I}, \Phi_1, \Phi_2, \dots, \Phi_M \in \mathbf{N}$ satisfy for all $m \in \{1, 2, \dots, M\}$, $x \in \mathbb{R}^{\mathfrak{d}}$ that $\mathcal{D}(\mathbb{I}) = (\mathfrak{d}, \mathbf{i}, \mathfrak{d})$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, $\mathcal{I}(\Phi_m) = d$, $\mathcal{O}(\Phi_m) = \mathfrak{d}$, and $L = \max_{m \in \{1, 2, \dots, M\}} \mathcal{L}(\Phi_m)$ (cf. Definition 2.1 and Definition 2.3). Then there exists $\Psi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^{\mathfrak{d}})$,*

(ii) *it holds for all $x \in \mathbb{R}^d$ that*

$$(\mathcal{R}_a(\Psi))(x) = \sum_{m=1}^M h_m (\mathcal{R}_a(\Phi_m))(x), \quad (137)$$

and

(iii) *it holds that*

$$\begin{aligned} \mathcal{P}(\Psi) \leq & \frac{1}{2} \left(\left[\sum_{m=1}^M \left[\max \left\{ 1, \frac{\mathbf{i}}{\mathfrak{d}} \right\} \right] \mathcal{P}(\Phi_m) \mathbb{1}_{(\mathcal{L}(\Phi_m), \infty)}(L) \right] \right. \\ & + \left[\sum_{m=1}^M \left((L - \mathcal{L}(\Phi_m) - 1) \mathbf{i}(\mathbf{i} + 1) + \mathfrak{d}(\mathbf{i} + 1) \right) \mathbb{1}_{(\mathcal{L}(\Phi_m), \infty)}(L) \right] \\ & \left. + \left[\sum_{m=1}^M \mathcal{P}(\Phi_m) \mathbb{1}_{\{\mathcal{L}(\Phi_m)\}}(L) \right] \right)^2. \end{aligned} \quad (138)$$

Proof of Proposition 2.26. Throughout this proof let $\mathfrak{J} = (\mathfrak{J}_1, \mathfrak{J}_2, \dots, \mathfrak{J}_M) \in \mathbf{N}^M$ satisfy for all $m \in \{1, 2, \dots, M\}$ that $\mathfrak{J}_m = \mathbb{I}$, let $(A_1, b_1) \in \mathbb{R}^{\mathfrak{d} \times (M\mathfrak{d})} \times \mathbb{R}^{\mathfrak{d}}$, $(A_2, b_2) \in \mathbb{R}^{(Md) \times d} \times \mathbb{R}^{Md}$ satisfy that

$$A_1 = (h_1 \mathbb{I}_{\mathfrak{d}} \quad h_2 \mathbb{I}_{\mathfrak{d}} \quad \dots \quad h_M \mathbb{I}_{\mathfrak{d}}), \quad A_2 = \begin{pmatrix} \mathbb{I}_d \\ \mathbb{I}_d \\ \vdots \\ \mathbb{I}_d \end{pmatrix}, \quad b_1 = 0, \quad \text{and} \quad b_2 = 0 \quad (139)$$

(cf. Definition 2.10), let $\mathbb{A}_1, \mathbb{A}_2 \in \mathbf{N}$ satisfy that $\mathbb{A}_1 = (A_1, b_1)$ and $\mathbb{A}_2 = (A_2, b_2)$, and let $\Psi \in \mathbf{N}$ satisfy that

$$\Psi = \mathbb{A}_1 \bullet (\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)) \bullet \mathbb{A}_2 \quad (140)$$

(cf. Definition 2.5, Definition 2.22, Lemma 2.8, and Corollary 2.23). Note that (140) and items (i)–(ii) in Corollary 2.9 demonstrate that

$$\begin{aligned} \mathcal{P}(\Psi) &\leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_1)}{\mathcal{O}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M))} \right\} \right] \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_2)+1}{\mathcal{I}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M))+1} \right\} \right] \\ &\quad \cdot \mathcal{P}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)) \\ &= \left[\max \left\{ 1, \frac{\mathfrak{d}}{M\mathfrak{d}} \right\} \right] \left[\max \left\{ 1, \frac{d+1}{Md+1} \right\} \right] \mathcal{P}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)) \\ &= \mathcal{P}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)). \end{aligned} \quad (141)$$

Corollary 2.24 hence proves that

$$\begin{aligned} \mathcal{P}(\Psi) &\leq \mathcal{P}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)) \\ &\leq \frac{1}{2} \left(\left[\sum_{m=1}^M \left[\max \left\{ 1, \frac{\mathfrak{i}}{\mathfrak{d}} \right\} \right] \mathcal{P}(\Phi_m) \mathbb{1}_{(\mathcal{L}(\Phi_m), \infty)}(L) \right] \right. \\ &\quad \left. + \left[\sum_{m=1}^M \left((L - \mathcal{L}(\Phi_m) - 1) \mathfrak{i}(\mathfrak{i} + 1) + \mathfrak{d}(\mathfrak{i} + 1) \right) \mathbb{1}_{(\mathcal{L}(\Phi_m), \infty)}(L) \right] \right. \\ &\quad \left. + \left[\sum_{m=1}^M \mathcal{P}(\Phi_m) \mathbb{1}_{\{\mathcal{L}(\Phi_m)\}}(L) \right] \right)^2. \end{aligned} \quad (142)$$

Next note that (139) and the fact that $\mathbb{A}_2 = (A_2, b_2)$ prove that $\mathcal{R}_a(\mathbb{A}_2) \in C(\mathbb{R}^d, \mathbb{R}^{Md})$ and

$$\forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{A}_2))(x) = (x, x, \dots, x) \in \mathbb{R}^{Md}. \quad (143)$$

Corollary 2.23 and item (v) in Proposition 2.6 therefore ensure that

$$\begin{aligned} \mathcal{R}_a(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M) \bullet \mathbb{A}_2) &\in C(\mathbb{R}^{\mathcal{I}(\mathbb{A}_2)}, \mathbb{R}^{\mathcal{O}(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M))}) \\ &= C(\mathbb{R}^d, \mathbb{R}^{Md}) \end{aligned} \quad (144)$$

and

$$\begin{aligned} \forall x \in \mathbb{R}^d: (\mathcal{R}_a(\mathbb{P}_{M, \mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M) \bullet \mathbb{A}_2))(x) \\ = ((\mathcal{R}_a(\Phi_1))(x), (\mathcal{R}_a(\Phi_2))(x), \dots, (\mathcal{R}_a(\Phi_M))(x)). \end{aligned} \quad (145)$$

In addition, observe that (139) and the fact that $\mathbb{A}_1 = (A_1, b_1)$ assure that $\mathcal{R}_a(\mathbb{A}_1) \in C(\mathbb{R}^{M\mathfrak{d}}, \mathbb{R}^{\mathfrak{d}})$ and

$$\forall y_1, y_2, \dots, y_M \in \mathbb{R}^{\mathfrak{d}}: (\mathcal{R}_a(\mathbb{A}_1))(y_1, y_2, \dots, y_M) = \sum_{m=1}^M h_m y_m. \quad (146)$$

Combining this, (144), (145), and (140) with item (v) in Proposition 2.6 demonstrates that

$$\mathcal{R}_a(\Psi) \in C(\mathbb{R}^{\mathcal{I}((P_{M,\mathcal{J}}(\Phi_1, \Phi_2, \dots, \Phi_M)) \bullet \mathbb{A}_2)}, \mathbb{R}^{\mathcal{O}(\mathbb{A}_1)}) = C(\mathbb{R}^d, \mathbb{R}^d) \quad (147)$$

and

$$\forall x \in \mathbb{R}^d: (\mathcal{R}_a(\Psi))(x) = \sum_{m=1}^M h_m(\mathcal{R}_a(\Phi_m))(x). \quad (148)$$

This and (142) establish items (i)–(iii). The proof of Proposition 2.26 is thus completed. \square

2.5 ANN representations for Euler approximations

2.5.1 ANN representations for one Euler step

Lemma 2.27. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $L_1 \in \mathbb{N} \cap [2, \infty)$, $L_2 \in \mathbb{N}$, $d, \mathbf{i}, l_{1,0}, l_{1,1}, \dots, l_{1,L_1}, l_{2,0}, l_{2,1}, \dots, l_{2,L_2} \in \mathbb{N}$, $\mathbb{I}, \Phi_1, \Phi_2 \in \mathbf{N}$ satisfy for all $k \in \{1, 2\}$, $x \in \mathbb{R}^d$ that $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, $\mathcal{I}(\Phi_k) = \mathcal{O}(\Phi_k) = d$, and $\mathcal{D}(\Phi_k) = (l_{k,0}, l_{k,1}, \dots, l_{k,L_k})$ (cf. Definition 2.1 and Definition 2.3). Then there exists $\Psi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,*

(ii) *it holds for all $x \in \mathbb{R}^d$ that*

$$(\mathcal{R}_a(\Psi))(x) = (\mathcal{R}_a(\Phi_2))(x) + ((\mathcal{R}_a(\Phi_1)) \circ (\mathcal{R}_a(\Phi_2)))(x), \quad (149)$$

(iii) *it holds that*

$$\mathcal{D}(\Psi) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, l_{1,L_1}), \quad (150)$$

and

(iv) *it holds that*

$$\begin{aligned} \mathcal{P}(\Psi) &= \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + (\mathbf{i} - d)(l_{2,L_2-1} + 1) + l_{1,1}(l_{2,L_2-1} - d) \\ &\quad + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) + \mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m} \right] + \mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m} \right]. \end{aligned} \quad (151)$$

Proof of Lemma 2.27. Throughout this proof let $A_1 \in \mathbb{R}^{d \times 2d}$, $A_2 \in \mathbb{R}^{2d \times d}$, $b_1 \in \mathbb{R}^d$, $b_2 \in \mathbb{R}^{2d}$ satisfy that

$$A_1 = \begin{pmatrix} \mathbf{I}_d & \mathbf{I}_d \end{pmatrix}, \quad A_2 = \begin{pmatrix} \mathbf{I}_d \\ \mathbf{I}_d \end{pmatrix}, \quad b_1 = 0, \quad \text{and} \quad b_2 = 0 \quad (152)$$

(cf. Definition 2.10) and let $\mathbb{A}_1 \in (\mathbb{R}^{d \times 2d} \times \mathbb{R}^d) \subseteq \mathbf{N}$, $\mathbb{A}_2 \in (\mathbb{R}^{2d \times d} \times \mathbb{R}^{2d}) \subseteq \mathbf{N}$, $\Psi \in \mathbf{N}$ satisfy that $\mathbb{A}_1 = (A_1, b_1)$, $\mathbb{A}_2 = (A_2, b_2)$, and

$$\Psi = \mathbb{A}_1 \bullet [\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2 \bullet \Phi_2 \quad (153)$$

(cf. Definition 2.5, Definition 2.11, Definition 2.17, Lemma 2.8, and item (i) in Lemma 2.13). Observe that (152) and the fact that $\mathbb{A}_2 = (A_2, b_2)$ ensure that for all $x \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_a(\mathbb{A}_2) \in C(\mathbb{R}^d, \mathbb{R}^{2d}) \quad \text{and} \quad (\mathcal{R}_a(\mathbb{A}_2))(x) = (x, x). \quad (154)$$

Item (v) in Proposition 2.6, item (i) in Lemma 2.14, and Proposition 2.19 hence imply that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a([\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2) \in C(\mathbb{R}^d, \mathbb{R}^{2d})$ and

$$\begin{aligned} (\mathcal{R}_a([\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2))(x) &= (\mathcal{R}_a(\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})))(x, x) \\ &= ((\mathcal{R}_a(\Phi_1))(x), (\mathcal{R}_a(\mathbb{I}^{\bullet(L_1-1)}))(x)) = ((\mathcal{R}_a(\Phi_1))(x), x). \end{aligned} \quad (155)$$

Item (v) in Proposition 2.6 therefore demonstrates that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a([\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2 \bullet \Phi_2) \in C(\mathbb{R}^d, \mathbb{R}^{2d})$ and

$$(\mathcal{R}_a([\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2 \bullet \Phi_2))(x) = \left((\mathcal{R}_a(\Phi_1))((\mathcal{R}_a(\Phi_2))(x)), (\mathcal{R}_a(\Phi_2))(x) \right). \quad (156)$$

In addition, note that (152) and the fact that $\mathbb{A}_1 = (A_1, b_1)$ ensure that for all $y = (y_1, y_2) \in \mathbb{R}^d \times \mathbb{R}^d$ it holds that

$$\mathcal{R}_a(\mathbb{A}_1) \in C(\mathbb{R}^{2d}, \mathbb{R}^d) \quad \text{and} \quad (\mathcal{R}_a(\mathbb{A}_1))(y) = y_1 + y_2. \quad (157)$$

Item (v) in Proposition 2.6, (153), and (156) hence prove that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$ and

$$(\mathcal{R}_a(\Psi))(x) = (\mathcal{R}_a(\Phi_1))((\mathcal{R}_a(\Phi_2))(x)) + (\mathcal{R}_a(\Phi_2))(x). \quad (158)$$

Next note that item (i) in Lemma 2.13 and item (i) in Proposition 2.20 demonstrate that

$$\mathcal{D}(\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})) = (2d, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, 2d). \quad (159)$$

Item (i) in Proposition 2.6 therefore ensures that

$$\mathcal{D}(\mathbb{A}_1 \bullet [\mathbf{P}_2(\Phi_1, \mathbb{I}^{\bullet(L_1-1)})] \bullet \mathbb{A}_2) = (d, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, d). \quad (160)$$

Combining this with item (i) in Proposition 2.6, (153), and the fact that $\mathcal{O}(\Phi_2) = l_{2,L_2} = d$ shows that

$$\mathcal{D}(\Psi) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, d). \quad (161)$$

The fact that $l_{1,L_1} = \mathcal{O}(\Phi_1) = d$ hence ensures that

$$\begin{aligned}
\mathcal{P}(\Psi) &= \left[\sum_{m=1}^{L_2-1} l_{2,m}(l_{2,m-1} + 1) \right] + (l_{1,1} + \mathbf{i})(l_{2,L_2-1} + 1) \\
&\quad + \left[\sum_{m=2}^{L_1-1} (l_{1,m} + \mathbf{i})(l_{1,m-1} + \mathbf{i} + 1) \right] + d(l_{1,L_1-1} + \mathbf{i} + 1) \\
&= \mathcal{P}(\Phi_2) - l_{2,L_2}(l_{2,L_2-1} + 1) + (l_{1,1} + \mathbf{i})(l_{2,L_2-1} + 1) \\
&\quad + \mathbf{i} \left[\sum_{m=2}^{L_1-1} l_{1,m} \right] + \mathbf{i} \left[\sum_{m=2}^{L_1-1} l_{1,m-1} \right] + \left[\sum_{m=2}^{L_1-1} l_{1,m}(l_{1,m-1} + 1) \right] \\
&\quad + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) + l_{1,L_1}(l_{1,L_1-1} + 1) + l_{1,L_1}\mathbf{i}.
\end{aligned} \tag{162}$$

This, the fact that $l_{2,L_2} = \mathcal{O}(\Phi_2) = d$, and the fact that $l_{1,0} = \mathcal{I}(\Phi_1) = d$ demonstrate that

$$\begin{aligned}
\mathcal{P}(\Psi) &= \mathcal{P}(\Phi_2) + (l_{1,1} - d + \mathbf{i})(l_{2,L_2-1} + 1) + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) \\
&\quad + \mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m} \right] + \mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m} \right] + \mathcal{P}(\Phi_1) - l_{1,1}(l_{1,0} + 1) \\
&= \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + (\mathbf{i} - d)(l_{2,L_2-1} + 1) + l_{1,1}(l_{2,L_2-1} - d) \\
&\quad + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) + \mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m} \right] + \mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m} \right].
\end{aligned} \tag{163}$$

Combining this with (158) and (161) establishes items (i)–(iv). The proof of Lemma 2.27 is thus completed. \square

Proposition 2.28. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $L_1 \in \mathbb{N} \cap [2, \infty)$, $L_2 \in \mathbb{N}$, $\mathbb{I}, \Phi_1, \Phi_2 \in \mathbf{N}$, $d, \mathbf{i}, l_{1,0}, l_{1,1}, \dots, l_{1,L_1}, l_{2,0}, l_{2,1}, \dots, l_{2,L_2} \in \mathbb{N}$ satisfy for all $k \in \{1, 2\}$, $x \in \mathbb{R}^d$ that $2 \leq \mathbf{i} \leq 2d$, $l_{2,L_2-1} \leq l_{1,L_1-1} + \mathbf{i}$, $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, $\mathcal{I}(\Phi_k) = \mathcal{O}(\Phi_k) = d$, and $\mathcal{D}(\Phi_k) = (l_{k,0}, l_{k,1}, \dots, l_{k,L_k})$ (cf. Definition 2.1 and Definition 2.3). Then there exists $\Psi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,*

(ii) *it holds for all $x \in \mathbb{R}^d$ that*

$$(\mathcal{R}_a(\Psi))(x) = (\mathcal{R}_a(\Phi_2))(x) + ((\mathcal{R}_a(\Phi_1)) \circ (\mathcal{R}_a(\Phi_2)))(x), \tag{164}$$

(iii) *it holds that*

$$\mathcal{D}(\Psi) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, l_{1,L_1}), \tag{165}$$

and

(iv) it holds that

$$\begin{aligned}\mathcal{P}(\Psi) &\leq \mathcal{P}(\Phi_2) + \mathcal{P}(\Phi_1) \left[\frac{1}{4} \mathcal{P}(\Phi_1) + \mathcal{P}(\mathbb{I}) - 1 \right] \\ &\leq \mathcal{P}(\Phi_2) + \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\Phi_1) \right]^2.\end{aligned}\tag{166}$$

Proof of Proposition 2.28. Throughout this proof let $\Psi \in \mathbf{N}$ satisfy that

(I) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(II) it holds for all $x \in \mathbb{R}^d$ that

$$(\mathcal{R}_a(\Psi))(x) = (\mathcal{R}_a(\Phi_2))(x) + ((\mathcal{R}_a(\Phi_1)) \circ (\mathcal{R}_a(\Phi_2)))(x),\tag{167}$$

(III) it holds that

$$\mathcal{D}(\Psi) = (l_{2,0}, l_{2,1}, \dots, l_{2,L_2-1}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1,L_1-1} + \mathbf{i}, l_{1,L_1}),\tag{168}$$

and

(IV) it holds that

$$\begin{aligned}\mathcal{P}(\Psi) &= \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + (\mathbf{i} - d)(l_{2,L_2-1} + 1) + l_{1,1}(l_{2,L_2-1} - d) \\ &\quad + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) + \mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m} \right] + \mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m} \right]\end{aligned}\tag{169}$$

(cf. Lemma 2.27). Note that the fact that $l_{1,0} = \mathcal{I}(\Phi_1) = d = \mathcal{O}(\Phi_1) = l_{1,L_1}$ implies that

$$\mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m} \right] \leq \frac{1}{2} \mathbf{i} \left[\sum_{m=2}^{L_1} l_{1,m}(l_{1,m-1} + 1) \right] = \frac{1}{2} \mathbf{i} [\mathcal{P}(\Phi_1) - l_{1,1}(d + 1)]\tag{170}$$

and

$$\begin{aligned}\mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m} \right] &\leq \frac{1}{2} \mathbf{i} \left[\sum_{m=1}^{L_1-2} l_{1,m}(l_{1,m-1} + 1) \right] \\ &= \frac{1}{2} \mathbf{i} [\mathcal{P}(\Phi_1) - d(l_{1,L_1-1} + 1) - l_{1,L_1-1}(l_{1,L_1-2} + 1)].\end{aligned}\tag{171}$$

Combining this with (IV) and the hypothesis that $l_{2,L_2-1} \leq l_{1,L_1-1} + \mathbf{i}$ ensures that

$$\begin{aligned}\mathcal{P}(\Psi) &\leq [1 + \mathbf{i}] \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + (\mathbf{i} - d)(l_{2,L_2-1} + 1) + l_{1,1}(l_{2,L_2-1} - d) \\ &\quad + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) - \frac{1}{2} \mathbf{i} l_{1,1}(d + 1) \\ &\quad - \frac{1}{2} \mathbf{i} d(l_{1,L_1-1} + 1) - \frac{1}{2} \mathbf{i} l_{1,L_1-1}(l_{1,L_1-2} + 1) \\ &\leq [1 + \mathbf{i}] \mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + [\max\{\mathbf{i} - d, 0\}](l_{1,L_1-1} + \mathbf{i} + 1) \\ &\quad + l_{1,1} [l_{1,L_1-1} + \mathbf{i} - d - \frac{1}{2} \mathbf{i} (d + 1)] + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) \\ &\quad - \frac{1}{2} \mathbf{i} d(l_{1,L_1-1} + 1) - \frac{1}{2} \mathbf{i} l_{1,L_1-1}(l_{1,L_1-2} + 1).\end{aligned}\tag{172}$$

Moreover, observe that the hypothesis that $2 \leq \mathbf{i} \leq 2d$ shows that

$$\begin{aligned} & l_{1,1}[\mathbf{i} - d - \tfrac{1}{2}\mathbf{i}(d+1)] - \tfrac{1}{2}\mathbf{i}d(l_{1,L_1-1} + 1) - \tfrac{1}{2}\mathbf{i}l_{1,L_1-1}(l_{1,L_1-2} + 1) \\ & \leq l_{1,1}[2d - d - (d+1)] - \tfrac{1}{2}\mathbf{i}l_{1,L_1-1} - \mathbf{i}l_{1,L_1-1} \leq -\tfrac{3}{2}\mathbf{i}l_{1,L_1-1}. \end{aligned} \quad (173)$$

This and (172) prove that

$$\begin{aligned} \mathcal{P}(\Psi) & \leq [1 + \mathbf{i}]\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + [\max\{\mathbf{i} - d, 0\}]l_{1,L_1-1} \\ & \quad + [\max\{\mathbf{i} - d, 0\}](\mathbf{i} + 1) + l_{1,1}l_{1,L_1-1} - \tfrac{3}{2}\mathbf{i}l_{1,L_1-1} + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) \\ & \leq [1 + \mathbf{i}]\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + [\max\{\mathbf{i} - d, 0\}]l_{1,L_1-1} \\ & \quad + \mathbf{i}(\mathbf{i} + 1) + l_{1,1}l_{1,L_1-1} + (L_1 - 2)\mathbf{i}(\mathbf{i} + 1) - \tfrac{3}{2}\mathbf{i}l_{1,L_1-1} \\ & \leq [1 + \mathbf{i}]\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + [\max\{\mathbf{i} - d, 0\}]l_{1,L_1-1} + l_{1,1}l_{1,L_1-1} \\ & \quad + (L_1 - 1)\mathbf{i}(\mathbf{i} + 1) - \tfrac{3}{2}\mathbf{i}l_{1,L_1-1} \\ & \leq [1 + \mathbf{i}]\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}l_{1,L_1-1} + (L_1 - 1)\mathbf{i}(\mathbf{i} + 1). \end{aligned} \quad (174)$$

Moreover, observe that

$$\begin{aligned} L_1 - 1 & \leq \left[\sum_{m=1}^{L_1} l_{1,m} \right] - 1 \leq \frac{1}{2} \left[\sum_{m=1}^{L_1} l_{1,m}(l_{1,m-1} + 1) \right] - 1 \\ & \leq \frac{1}{2}\mathcal{P}(\Phi_1) - 1 \leq \frac{1}{2}\mathcal{P}(\Phi_1). \end{aligned} \quad (175)$$

Combining this and (174) with the fact that $\forall k \in \mathbb{N} \cap [1, L_1]: l_{1,k} \leq \frac{1}{2}l_{1,k}(l_{1,k-1} + 1) \leq \frac{1}{2}\mathcal{P}(\Phi_1)$ demonstrates that

$$\begin{aligned} \mathcal{P}(\Psi) & \leq [1 + \mathbf{i}]\mathcal{P}(\Phi_1) + \mathcal{P}(\Phi_2) + l_{1,1}l_{1,L_1-1} + \tfrac{1}{2}\mathcal{P}(\Phi_1)\mathbf{i}(\mathbf{i} + 1) \\ & = \mathcal{P}(\Phi_2) + [1 + \mathbf{i} + \tfrac{1}{2}\mathbf{i}(\mathbf{i} + 1)]\mathcal{P}(\Phi_1) + l_{1,1}l_{1,L_1-1} \\ & \leq \mathcal{P}(\Phi_2) + [1 + \mathbf{i} + \tfrac{1}{2}\mathbf{i}(\mathbf{i} + 1)]\mathcal{P}(\Phi_1) + \tfrac{1}{4}[\mathcal{P}(\Phi_1)]^2. \end{aligned} \quad (176)$$

Furthermore, note that the hypothesis that $2 \leq \mathbf{i} \leq 2d$ and the hypothesis that $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$ prove that

$$\begin{aligned} \mathbf{i} + \tfrac{1}{2}\mathbf{i}(\mathbf{i} + 1) & = \mathbf{i}^2 + \mathbf{i} + \tfrac{1}{2}\mathbf{i} - \tfrac{1}{2}\mathbf{i}^2 \leq 2d\mathbf{i} + \mathbf{i} + d - \tfrac{1}{2}\mathbf{i}^2 \\ & = \mathbf{i}(d+1) + d(\mathbf{i} + 1) - \tfrac{1}{2}\mathbf{i}^2 = \mathcal{P}(\mathbb{I}) - \tfrac{1}{2}\mathbf{i}^2 \leq \mathcal{P}(\mathbb{I}) - 2. \end{aligned} \quad (177)$$

Combining this and (176) implies that

$$\begin{aligned} \mathcal{P}(\Psi) & \leq \mathcal{P}(\Phi_2) + [1 + \mathcal{P}(\mathbb{I}) - 2]\mathcal{P}(\Phi_1) + \tfrac{1}{4}[\mathcal{P}(\Phi_1)]^2 \\ & = \mathcal{P}(\Phi_2) + [\tfrac{1}{4}\mathcal{P}(\Phi_1) + \mathcal{P}(\mathbb{I}) - 1]\mathcal{P}(\Phi_1) \\ & \leq \mathcal{P}(\Phi_2) + \mathcal{P}(\mathbb{I})\mathcal{P}(\Phi_1) + \tfrac{1}{4}[\mathcal{P}(\mathbb{I})]^2 + [\mathcal{P}(\Phi_1)]^2 \\ & = \mathcal{P}(\Phi_2) + [\tfrac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\Phi_1)]^2. \end{aligned} \quad (178)$$

This, (I), (II), and (III) establish items (i)–(iv). The proof of Proposition 2.28 is thus completed. \square

2.5.2 ANN representations for multiple nested Euler steps

Corollary 2.29. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $d, \mathbf{i}, \mathfrak{L} \in \mathbb{N}$, $\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}} \in \mathbb{N}$, $(L_n)_{n \in \mathbb{N}_0} \subseteq \mathbb{N} \cap [2, \infty)$, $\mathbb{I}, \psi \in \mathbf{N}$, $(\phi_n)_{n \in \mathbb{N}_0} \subseteq \mathbf{N}$, let $l_{n,k} \in \mathbb{N}$, $k \in \{0, 1, \dots, L_n\}$, $n \in \mathbb{N}_0$, assume for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that $2 \leq \mathbf{i} \leq 2d$, $\ell_{\mathfrak{L}-1} \leq l_{0, L_0-1} + \mathbf{i}$, $l_{n, L_n-1} \leq l_{n+1, L_{n+1}-1}$, $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, $\mathcal{I}(\phi_n) = \mathcal{O}(\phi_n) = \mathcal{I}(\psi) = \mathcal{O}(\psi) = d$, $\mathcal{D}(\phi_n) = (l_{n,0}, l_{n,1}, \dots, l_{n, L_n})$, and $\mathcal{D}(\psi) = (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}})$, and let $f_n: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that*

$$f_0(x) = (\mathcal{R}_a(\psi))(x) \quad \text{and} \quad f_{n+1}(x) = f_n(x) + ([\mathcal{R}_a(\phi_n)] \circ f_n)(x) \quad (179)$$

(cf. Definition 2.1 and Definition 2.3). Then for every $n \in \mathbb{N}$ there exists $\Psi \in \mathbf{N}$ such that

- (i) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,
- (ii) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$,
- (iii) it holds that $\mathcal{H}(\Psi) = \mathcal{H}(\psi) + \sum_{k=0}^{n-1} \mathcal{H}(\phi_k)$,
- (iv) it holds that

$$\begin{aligned} \mathcal{D}(\Psi) = & (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}-1}, l_{0,1} + \mathbf{i}, l_{0,2} + \mathbf{i}, \dots, l_{0, L_0-1} + \mathbf{i}, \\ & l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \dots, l_{1, L_1-1} + \mathbf{i}, \dots, l_{n-1,1} + \mathbf{i}, l_{n-1,2} + \mathbf{i}, \dots, l_{n-1, L_{n-1}-1} + \mathbf{i}, d), \end{aligned} \quad (180)$$

and

- (v) it holds that $\mathcal{P}(\Psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{n-1} [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k)]^2$.

Proof of Corollary 2.29. We prove items (i)–(v) by induction on $n \in \mathbb{N}$. Note that the hypothesis that $\mathcal{D}(\psi) = (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}})$, the fact that $\ell_0 = \mathcal{I}(\psi) = \ell_{\mathfrak{L}} = \mathcal{O}(\psi) = d$, the hypothesis that $\mathcal{D}(\phi_0) = (l_{0,0}, l_{0,1}, \dots, l_{0, L_0})$, the hypothesis that $\ell_{\mathfrak{L}-1} \leq l_{0, L_0-1} + \mathbf{i}$, the hypothesis that $L_0 \in \mathbb{N} \cap [2, \infty)$, Proposition 2.28 (with $a = a$, $L_1 = L_0$, $L_2 = \mathfrak{L}$, $\mathbb{I} = \mathbb{I}$, $\Phi_1 = \phi_0$, $\Phi_2 = \psi$, $d = d$, $\mathbf{i} = \mathbf{i}$, $l_{1,v} = l_{0,v}$, $l_{2,w} = \ell_w$ for $v \in \{0, 1, \dots, L_0\}$, $w \in \{0, 1, \dots, \mathfrak{L}\}$ in the notation of Proposition 2.28), and (179) imply that there exists $\Upsilon \in \mathbf{N}$ which satisfies that

- (I) it holds that $\mathcal{R}_a(\Upsilon) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

- (II) it holds for all $x \in \mathbb{R}^d$ that

$$\begin{aligned} (\mathcal{R}_a(\Upsilon))(x) &= (\mathcal{R}_a(\psi))(x) + ([\mathcal{R}_a(\phi_0)] \circ [\mathcal{R}_a(\psi)])(x) \\ &= f_0(x) + ([\mathcal{R}_a(\phi_0)] \circ f_0)(x) = f_1(x), \end{aligned} \quad (181)$$

(III) it holds that

$$\mathcal{D}(\Upsilon) = (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}-1}, l_{0,1} + \mathbf{i}, l_{0,2} + \mathbf{i}, \dots, l_{0,L_0-1} + \mathbf{i}, l_{0,L_0}), \quad (182)$$

and

(IV) it holds that $\mathcal{P}(\Upsilon) \leq \mathcal{P}(\psi) + [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_0)]^2$.

Observe that (III) shows that $\mathcal{L}(\Upsilon) = \mathfrak{L} + L_0 - 1$. Hence, we obtain that

$$\mathcal{H}(\Upsilon) = \mathcal{L}(\Upsilon) - 1 = (\mathfrak{L} - 1) + (L_0 - 1) = \mathcal{H}(\psi) + \mathcal{H}(\phi_0). \quad (183)$$

Combining this with (I)–(IV) establishes items (i)–(v) in the base case $n = 1$. For the induction step $\mathbb{N} \ni n \rightarrow n+1 \in \mathbb{N} \cap [2, \infty)$ let $n \in \mathbb{N}$, $\Psi \in \mathbf{N}$, $l_0, l_1, \dots, l_{\mathfrak{L} + \sum_{k=0}^{n-1} (L_k - 1)} \in \mathbb{N}$ satisfy that

- (a) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,
- (b) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$,
- (c) it holds that $\mathcal{H}(\Psi) = \mathcal{H}(\psi) + \sum_{k=0}^{n-1} \mathcal{H}(\phi_k)$,
- (d) it holds that

$$\begin{aligned} \mathcal{D}(\Psi) &= (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}-1}, l_{0,1} + \mathbf{i}, l_{0,2} + \mathbf{i}, \dots, l_{0,L_0-1} + \mathbf{i}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \\ &\quad \dots, l_{1,L_1-1} + \mathbf{i}, \dots, l_{n-1,1} + \mathbf{i}, l_{n-1,2} + \mathbf{i}, \dots, l_{n-1,L_{n-1}-1} + \mathbf{i}, d) \\ &= (l_0, l_1, \dots, l_{\mathfrak{L} + \sum_{k=0}^{n-1} (L_k - 1)}), \end{aligned} \quad (184)$$

and

(e) it holds that $\mathcal{P}(\Psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{n-1} [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k)]^2$.

Observe that (d) and the hypothesis that $\forall k \in \mathbb{N}_0: l_{k,L_k-1} \leq l_{k+1,L_{k+1}-1}$ demonstrate that

$$l_{\mathcal{L}(\Psi)-1} = l_{\mathfrak{L}-1 + \sum_{k=0}^{n-1} (L_k - 1)} = l_{n-1, L_{n-1}-1} + \mathbf{i} \leq l_{n, L_n} + \mathbf{i}. \quad (185)$$

The hypothesis that $\mathcal{D}(\phi_n) = (l_{n,0}, l_{n,1}, \dots, l_{n,L_n})$, (d), the hypothesis that $L_n \in \mathbb{N} \cap [2, \infty)$, and Proposition 2.28 (with $a = a$, $L_1 = L_n$, $L_2 = \mathfrak{L} + \sum_{k=0}^{n-1} (L_k - 1)$, $\mathbb{I} = \mathbb{I}$, $\Phi_1 = \phi_n$, $\Phi_2 = \Psi$, $d = d$, $\mathbf{i} = \mathbf{i}$, $l_{1,v} = l_{n,v}$, $l_{2,w} = l_w$ for $v \in \{0, 1, \dots, L_n\}$, $w \in \{0, 1, \dots, \mathfrak{L} + \sum_{k=0}^{n-1} (L_k - 1)\}$ in the notation of Proposition 2.28) hence prove that there exists $\Phi \in \mathbf{N}$ which satisfies that

(A) it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(B) it holds for all $x \in \mathbb{R}^d$ that

$$(\mathcal{R}_a(\Phi))(x) = (\mathcal{R}_a(\Psi))(x) + ([\mathcal{R}_a(\phi_n)] \circ [\mathcal{R}_a(\Psi)])(x), \quad (186)$$

(C) it holds that

$$\begin{aligned} \mathcal{D}(\Phi) = & (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}-1}, l_{0,1} + \mathbf{i}, l_{0,2} + \mathbf{i}, \dots, l_{0,L_0-1} + \mathbf{i}, l_{1,1} + \mathbf{i}, l_{1,2} + \mathbf{i}, \\ & \dots, l_{1,L_1-1} + \mathbf{i}, \dots, l_{n,1} + \mathbf{i}, l_{n,2} + \mathbf{i}, \dots, l_{n,L_n-1} + \mathbf{i}, l_{n,L_n}), \end{aligned} \quad (187)$$

and

(D) it holds that $\mathcal{P}(\Phi) \leq \mathcal{P}(\Psi) + [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_n)]^2$.

Next note that (C) implies that $\mathcal{L}(\Phi) = \mathfrak{L} + \sum_{k=0}^n (L_k - 1)$. Hence, we obtain that

$$\mathcal{H}(\Phi) = \mathcal{L}(\Phi) - 1 = (\mathfrak{L} - 1) + \sum_{k=0}^n (L_k - 1) = \mathcal{H}(\psi) + \sum_{k=0}^n \mathcal{H}(\phi_k). \quad (188)$$

Moreover, observe that (B), (179), and (b) demonstrate that for all $x \in \mathbb{R}^d$ it holds that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(x) &= (\mathcal{R}_a(\Psi))(x) + ([\mathcal{R}_a(\phi_n)] \circ [\mathcal{R}_a(\Psi)])(x) \\ &= f_n(x) + ([\mathcal{R}_a(\phi_n)] \circ f_n)(x) = f_{n+1}(x). \end{aligned} \quad (189)$$

In addition, note that (D) and (e) ensure that

$$\begin{aligned} \mathcal{P}(\Phi) &\leq \mathcal{P}(\psi) + \left[\sum_{k=0}^{n-1} [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k)]^2 \right] + [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_n)]^2 \\ &= \mathcal{P}(\psi) + \sum_{k=0}^n [\frac{1}{2}\mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k)]^2. \end{aligned} \quad (190)$$

This, (A), (C), (188), and (189) prove items (i)–(v) in the case $n + 1$. Induction thus establishes items (i)–(v). The proof of Corollary 2.29 is thus completed. \square

Proposition 2.30. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $d, \mathfrak{L} \in \mathbb{N}$, $\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}} \in \mathbb{N}$, $\psi \in \mathbf{N}$, $(\phi_n)_{n \in \mathbb{N}_0} \subseteq \mathbf{N}$ satisfy for all $n \in \mathbb{N}_0$ that $\mathcal{I}(\phi_n) = \mathcal{O}(\phi_n) = \mathcal{I}(\psi) = \mathcal{O}(\psi) = d$, $\mathcal{L}(\phi_n) = 1$, and $\mathcal{D}(\psi) = (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}})$ and let $f_n: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that*

$$f_0(x) = (\mathcal{R}_a(\psi))(x) \quad \text{and} \quad f_{n+1}(x) = f_n(x) + ([\mathcal{R}_a(\phi_n)] \circ f_n)(x) \quad (191)$$

(cf. Definition 2.1 and Definition 2.3). Then for every $n \in \mathbb{N}_0$ there exists $\Psi \in \mathbf{N}$ such that

(i) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(ii) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$, and

(iii) it holds that $\mathcal{D}(\Psi) = \mathcal{D}(\psi)$.

Proof of Proposition 2.30. We prove items (i)–(iii) by induction on $n \in \mathbb{N}_0$. Note that (191) and the fact that $\mathcal{R}_a(\psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$ establish items (i)–(iii) in the base case $n = 0$. For the induction step $\mathbb{N}_0 \ni n \rightarrow n + 1 \in \mathbb{N}$ let $n \in \mathbb{N}_0$, $\Psi \in \mathbf{N}$ satisfy that

(I) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(II) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$, and

(III) it holds that $\mathcal{D}(\Psi) = \mathcal{D}(\psi)$,

and let $(A, b) \in (\mathbb{R}^{d \times d} \times \mathbb{R}^d) \subseteq \mathbf{N}$, $\mathbb{A} \in (\mathbb{R}^{d \times d} \times \mathbb{R}^d) \subseteq \mathbf{N}$, $\Phi \in \mathbf{N}$ satisfy that $\phi_n = (A, b)$, $\mathbb{A} = (A + \mathbb{I}_d, b)$, and $\Phi = \mathbb{A} \bullet \Psi$ (cf. Definition 2.5 and Definition 2.10). Observe that item (v) in Proposition 2.6 demonstrates that for all $x \in \mathbb{R}^d$ it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^d, \mathbb{R}^d)$ and

$$\begin{aligned} (\mathcal{R}_a(\Phi))(x) &= (\mathcal{R}_a(\mathbb{A}))((\mathcal{R}_a(\Psi))(x)) \\ &= (A + \mathbb{I}_d)((\mathcal{R}_a(\Psi))(x)) + b \\ &= A((\mathcal{R}_a(\Psi))(x)) + b + (\mathcal{R}_a(\Psi))(x) \\ &= (\mathcal{R}_a(\phi_n))((\mathcal{R}_a(\Psi))(x)) + (\mathcal{R}_a(\Psi))(x). \end{aligned} \tag{192}$$

Combining this with (191) and (II) proves that for all $x \in \mathbb{R}^d$ it holds that

$$(\mathcal{R}_a(\Phi))(x) = (\mathcal{R}_a(\phi_n))(f_n(x)) + f_n(x) = f_{n+1}(x). \tag{193}$$

In addition, note that (III), the fact that $\Phi = \mathbb{A} \bullet \Psi$, the fact that $\mathcal{L}(\mathbb{A}) = 1$, the fact that $\mathcal{I}(\mathbb{A}) = \mathcal{O}(\mathbb{A}) = \mathcal{O}(\Psi) = d$, and item (i) in Proposition 2.6 imply that $\mathcal{D}(\Phi) = \mathcal{D}(\Psi) = \mathcal{D}(\psi)$. Combining this and the fact that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^d, \mathbb{R}^d)$ with (193) proves items (i)–(iii) in the case $n + 1$. Induction thus establishes items (i)–(iii). The proof of Proposition 2.30 is thus completed. \square

Corollary 2.31. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $d, \mathbf{i}, L, \mathfrak{L} \in \mathbb{N}$, $\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}} \in \mathbb{N}$, $\mathbb{I}, \psi \in \mathbf{N}$, $(\phi_n)_{n \in \mathbb{N}_0} \subseteq \mathbf{N}$, let $l_{n,k} \in \mathbb{N}$, $k \in \{0, 1, \dots, L\}$, $n \in \mathbb{N}_0$, assume for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that $2 \leq \mathbf{i} \leq 2d$, $\ell_{\mathfrak{L}-1} \leq l_{0,L-1} + \mathbf{i}$, $l_{n,L-1} \leq l_{n+1,L-1}$, $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, $\mathcal{I}(\phi_n) = \mathcal{O}(\phi_n) = \mathcal{I}(\psi) = \mathcal{O}(\psi) = d$, $\mathcal{D}(\phi_n) = (l_{n,0}, l_{n,1}, \dots, l_{n,L})$, and $\mathcal{D}(\psi) = (\ell_0, \ell_1, \dots, \ell_{\mathfrak{L}})$, and let $f_n: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that*

$$f_0(x) = (\mathcal{R}_a(\psi))(x) \quad \text{and} \quad f_{n+1}(x) = f_n(x) + ([\mathcal{R}_a(\phi_n)] \circ f_n)(x) \tag{194}$$

(cf. Definition 2.1 and Definition 2.3). Then for every $n \in \mathbb{N}_0$ there exists $\Psi \in \mathbf{N}$ such that

(i) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(ii) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$,

(iii) it holds that $\mathcal{H}(\Psi) = \mathcal{H}(\psi) + \sum_{k=0}^{n-1} \mathcal{H}(\phi_k) = \mathcal{H}(\psi) + n \mathcal{H}(\phi_0)$, and

(iv) it holds that $\mathcal{P}(\Psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{n-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k) \right]^2$.

Proof of Corollary 2.31. To prove items (i)–(iv) we distinguish between the case $L = 1$ and the case $L \in \mathbb{N} \cap [2, \infty)$. We first prove items (i)–(iv) in the case $L = 1$. Observe that Proposition 2.30 ensures that there exist $\Psi_n \in \mathbf{N}$, $n \in \mathbb{N}_0$, which satisfy that

(I) it holds for all $n \in \mathbb{N}_0$ that $\mathcal{R}_a(\Psi_n) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(II) it holds for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi_n))(x) = f_n(x)$, and

(III) it holds for all $n \in \mathbb{N}_0$ that $\mathcal{D}(\Psi_n) = \mathcal{D}(\psi)$.

Next note that the hypothesis that $L = 1$ demonstrates that for all $n \in \mathbb{N}_0$ it holds that $\mathcal{H}(\phi_n) = 0$. Combining this with (III) implies that for all $n \in \mathbb{N}_0$ it holds that

$$\mathcal{H}(\Psi_n) = \mathcal{H}(\psi) = \mathcal{H}(\psi) + \sum_{k=0}^{n-1} \mathcal{H}(\phi_k) = \mathcal{H}(\psi) + n \mathcal{H}(\phi_0). \quad (195)$$

In addition, observe that (III) shows that for all $n \in \mathbb{N}_0$ it holds that

$$\mathcal{P}(\Psi_n) = \mathcal{P}(\psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{n-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k) \right]^2. \quad (196)$$

Combining this and (195) with (I)–(II) establishes items (i)–(iv) in the case $L = 1$. We now prove items (i)–(iv) in the case $L \in \mathbb{N} \cap [2, \infty)$. Note that (194), the fact that $\mathcal{R}_a(\psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$, the fact that $\mathcal{H}(\psi) = \mathcal{H}(\psi) + \sum_{k=0}^{-1} \mathcal{H}(\phi_k) = \mathcal{H}(\psi) + 0 \cdot \mathcal{H}(\phi_0)$, and the fact that $\mathcal{P}(\psi) = \mathcal{P}(\psi) + \sum_{k=0}^{-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k) \right]^2$ prove that there exists $\Psi \in \mathbf{N}$ such that

(a) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(b) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_0(x)$,

(c) it holds that $\mathcal{H}(\Psi) = \mathcal{H}(\psi) + \sum_{k=0}^{-1} \mathcal{H}(\phi_k) = \mathcal{H}(\psi) + 0 \cdot \mathcal{H}(\phi_0)$, and

(d) it holds that $\mathcal{P}(\Psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k) \right]^2$.

Moreover, observe that Corollary 2.29 and the fact that for all $k \in \mathbb{N}_0$ it holds that $\mathcal{H}(\phi_k) = L - 1 = \mathcal{H}(\phi_0)$ ensure that for every $n \in \mathbb{N}$ there exists $\Psi \in \mathbf{N}$ such that

(A) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(B) it holds for all $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi))(x) = f_n(x)$,

(C) it holds that $\mathcal{H}(\Psi) = \mathcal{H}(\psi) + \sum_{k=0}^{n-1} \mathcal{H}(\phi_k) = \mathcal{H}(\psi) + n \mathcal{H}(\phi_0)$, and

(D) it holds that $\mathcal{P}(\Psi) \leq \mathcal{P}(\psi) + \sum_{k=0}^{n-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\phi_k) \right]^2$.

Combining this with (a)–(d) proves items (i)–(iv) in the case $L \in \mathbb{N} \cap [2, \infty)$. The proof of Corollary 2.31 is thus completed. \square

2.5.3 ANN representations for multiple perturbed nested Euler steps

Proposition 2.32. *Let $a \in C(\mathbb{R}, \mathbb{R})$, $N, d, \mathbf{i} \in \mathbb{N}$, $\mathbb{I}, \Phi \in \mathbf{N}$, $A_1, A_2, \dots, A_N \in \mathbb{R}^{d \times d}$ satisfy for all $x \in \mathbb{R}^d$ that $2 \leq \mathbf{i} \leq 2d$, $\mathcal{D}(\mathbb{I}) = (d, \mathbf{i}, d)$, $(\mathcal{R}_a(\mathbb{I}))(x) = x$, and $\mathcal{I}(\Phi) = \mathcal{O}(\Phi) = d$ and let $Y_n = (Y_n^{x,y})_{(x,y) \in \mathbb{R}^d \times (\mathbb{R}^d)^N} : \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$, $n \in \{0, 1, \dots, N\}$, be the functions which satisfy for all $n \in \{0, 1, \dots, N-1\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_0^{x,y} = x$ and*

$$Y_{n+1}^{x,y} = Y_n^{x,y} + A_{n+1}((\mathcal{R}_a(\Phi))(Y_n^{x,y})) + y_{n+1} \quad (197)$$

(cf. Definition 2.1 and Definition 2.3). Then there exists $(\Psi_{n,y})_{(n,y) \in \{0,1,\dots,N\} \times (\mathbb{R}^d)^N} \subseteq \mathbf{N}$ such that

(i) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_{n,y}) \in C(\mathbb{R}^d, \mathbb{R}^d)$,

(ii) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi_{n,y}))(x) = Y_n^{x,y}$,

(iii) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{H}(\Psi_{n,y}) = \mathcal{H}(\mathbb{I}) + n \mathcal{H}(\Phi) = 1 + n \mathcal{H}(\Phi), \quad (198)$$

(iv) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{P}(\Psi_{n,y}) \leq \mathcal{P}(\mathbb{I}) + n \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\Phi) \right]^2, \quad (199)$$

(v) it holds for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_{n,y}))(x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (200)$$

and

(vi) it holds for all $n \in \{0, 1, \dots, N\}$, $m \in \mathbb{N}_0 \cap [0, n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Psi_{m,y}))(x) = (\mathcal{R}_a(\Psi_{m,z}))(x). \quad (201)$$

Proof of Proposition 2.32. Throughout this proof let $l_0, l_1, \dots, l_{\mathcal{L}(\Phi)} \in \mathbb{N}$ satisfy that $\mathcal{D}(\Phi) = (l_0, l_1, \dots, l_{\mathcal{L}(\Phi)})$, let $\mathbb{A}_{n,b} \in (\mathbb{R}^{d \times d} \times \mathbb{R}^d) \subseteq \mathbf{N}$, $n \in \{1, 2, \dots, N\}$, $b \in \mathbb{R}^d$, satisfy for all $n \in \{1, 2, \dots, N\}$, $b \in \mathbb{R}^d$ that

$$\mathbb{A}_{n,b} = (A_n, b) \in (\mathbb{R}^{d \times d} \times \mathbb{R}^d), \quad (202)$$

let $\rho_{n,y} \in \mathbf{N}$, $n \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$, satisfy for all $n \in \mathbb{N}$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that

$$\rho_{n,y} = \mathbb{A}_{\min\{n,N\}, y_{\min\{n,N\}}} \bullet \Phi \quad (203)$$

(cf. Definition 2.5), and let $\mathcal{Y}_n = (\mathcal{Y}_n^{x,y})_{(x,y) \in \mathbb{R}^d \times (\mathbb{R}^d)^N} : \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $\mathcal{Y}_0^{x,y} = x$ and

$$\mathcal{Y}_{n+1}^{x,y} = \mathcal{Y}_n^{x,y} + (\mathcal{R}_a(\rho_{n+1,y}))(\mathcal{Y}_n^{x,y}). \quad (204)$$

Observe that item (i) in Proposition 2.6 and the fact that for all $n \in \{1, 2, \dots, N\}$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that $\rho_{n,y} = \mathbb{A}_{n,y_n} \bullet \Phi$ prove that for all $n \in \{1, 2, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that $\mathcal{D}(\rho_{n,y}) = \mathcal{D}(\Phi) = (l_0, l_1, \dots, l_{\mathcal{L}(\Phi)})$. Corollary 2.31 (with $a = a$, $d = d$, $\mathbf{i} = \mathbf{i}$, $L = \mathcal{L}(\Phi)$, $\mathfrak{L} = 2$, $\ell_0 = d$, $\ell_1 = \mathbf{i}$, $\ell_2 = d$, $\mathbb{I} = \mathbb{I}$, $\psi = \mathbb{I}$, $(\mathbb{N}_0 \ni n \mapsto \phi_n \in \mathbf{N}) = (\mathbb{N}_0 \ni n \mapsto \rho_{n+1,y} \in \mathbf{N})$, $(\mathbb{N}_0 \times \{0, 1, \dots, \mathcal{L}(\Phi)\} \ni (n, k) \mapsto l_{n,k} \in \mathbb{N}) = (\mathbb{N}_0 \times \{0, 1, \dots, \mathcal{L}(\Phi)\} \ni (n, k) \mapsto l_k \in \mathbb{N})$, $(\mathbb{N}_0 \ni n \mapsto f_n \in C(\mathbb{R}^d, \mathbb{R}^d)) = (\mathbb{N}_0 \ni n \mapsto (\mathbb{R}^d \ni x \mapsto \mathcal{Y}_n^{x,y} \in \mathbb{R}^d) \in C(\mathbb{R}^d, \mathbb{R}^d))$ for $y \in (\mathbb{R}^d)^N$ in the notation of Corollary 2.31) and the fact that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that $(\mathcal{R}_a(\mathbb{I}))(x) = x = \mathcal{Y}_0^{x,y} = Y_0^{x,y}$ hence prove that there exist $\Psi_{n,y} \in \mathbf{N}$, $(n, y) \in \{0, 1, \dots, N\} \times (\mathbb{R}^d)^N$, which satisfy that

- (I) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_{n,y}) \in C(\mathbb{R}^d, \mathbb{R}^d)$,
- (II) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Psi_{n,y}))(x) = \mathcal{Y}_n^{x,y} = Y_n^{x,y}$,
- (III) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{H}(\Psi_{n,y}) = \mathcal{H}(\mathbb{I}) + \sum_{k=0}^{n-1} \mathcal{H}(\rho_{k+1,y}) = 1 + n\mathcal{H}(\Phi), \quad (205)$$

and

- (IV) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{P}(\Psi_{n,y}) \leq \mathcal{P}(\mathbb{I}) + \sum_{k=0}^{n-1} \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\rho_{k+1,y}) \right]^2 = \mathcal{P}(\mathbb{I}) + n \left[\frac{1}{2} \mathcal{P}(\mathbb{I}) + \mathcal{P}(\Phi) \right]^2. \quad (206)$$

Next we claim that for all $n \in \{0, 1, \dots, N\}$ it holds that

$$\forall x \in \mathbb{R}^d: \left[((\mathbb{R}^d)^N \ni y \mapsto Y_n^{x,y} \in \mathbb{R}^d) \in C((\mathbb{R}^d)^N, \mathbb{R}^d) \right]. \quad (207)$$

We now prove (207) by induction on $n \in \{0, 1, \dots, N\}$. Note that the fact that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that $Y_0^{x,y} = x$ proves (207) in the base case $n = 0$. For the induction step observe that (197) and the fact that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^d, \mathbb{R}^d)$ ensure that for all $n \in \{0, 1, \dots, N-1\}$ with

$$\forall x \in \mathbb{R}^d: \left[((\mathbb{R}^d)^N \ni y \mapsto Y_n^{x,y} \in \mathbb{R}^d) \in C((\mathbb{R}^d)^N, \mathbb{R}^d) \right] \quad (208)$$

it holds that

$$\forall x \in \mathbb{R}^d: \left[((\mathbb{R}^d)^N \ni y \mapsto Y_{n+1}^{x,y} \in \mathbb{R}^d) \in C((\mathbb{R}^d)^N, \mathbb{R}^d) \right]. \quad (209)$$

Induction thus proves (207). In addition, observe that (207) and (II) imply that for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$ it holds that

$$((\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_{n,y}))(x) \in \mathbb{R}^d) \in C((\mathbb{R}^d)^N, \mathbb{R}^d). \quad (210)$$

Next let $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ satisfy for all $k \in \mathbb{N} \cap [0, n]$ that $y_k = z_k$. We claim that for all $m \in \mathbb{N}_0 \cap [0, n]$ it holds that

$$Y_m^{x,y} = Y_m^{x,z}. \quad (211)$$

We now prove (211) by induction on $m \in \mathbb{N}_0 \cap [0, n]$. Note that the fact that $Y_0^{x,y} = x = Y_0^{x,z}$ implies (211) in the base case $m = 0$. For the induction step observe that (197) and the fact that for all $k \in \mathbb{N} \cap [0, n]$ it holds that $y_k = z_k$ ensure that for all $m \in \mathbb{N}_0 \cap (-\infty, n)$ with $Y_m^{x,y} = Y_m^{x,z}$ it holds that

$$\begin{aligned} Y_{m+1}^{x,y} &= Y_m^{x,y} + A_{m+1}((\mathcal{R}_a(\Phi))(Y_m^{x,y})) + y_{m+1} \\ &= Y_m^{x,z} + A_{m+1}((\mathcal{R}_a(\Phi))(Y_m^{x,z})) + z_{m+1} = Y_{m+1}^{x,z}. \end{aligned} \quad (212)$$

Induction thus proves (211). Note that (211) and (II) assure that for all $n \in \{0, 1, \dots, N\}$, $m \in \mathbb{N}_0 \cap [0, n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ it holds that

$$(\mathcal{R}_a(\Psi_{m,y}))(x) = (\mathcal{R}_a(\Psi_{m,z}))(x). \quad (213)$$

Combining this with (210) and (I)–(IV) establishes items (i)–(vi). The proof of Proposition 2.32 is thus completed. \square

3 ANN approximation results

This section establishes in Theorem 3.12 in Subsection 3.3 below the main result of this article. Some of the material presented in Subsection 3.1 and Subsection 3.2 are well-known concepts and results in the scientific literature. In particular, the material in Subsection 3.1.1 and Subsection 3.1.2 consists mainly of reformulations of concepts and results in Elbrächter et al. [11, Appendix A.3 and Appendix A.4]. Moreover, our proof of Proposition 3.5 in Subsection 3.2.1 below is inspired by Elbrächter et al. [11, Section 6] and Yarotsky [37, Section 3.2]. Furthermore, Lemma 3.8 and Lemma 3.9 are elementary and essentially well-known in the scientific literature. In addition, our proof of Lemma 3.11 is based on a well-known Gronwall argument.

3.1 ANN approximations for the square function

3.1.1 Explicit approximations for the square function on $[0, 1]$

Lemma 3.1. *Let $g_n: \mathbb{R} \rightarrow [0, 1]$, $n \in \mathbb{N}$, be the functions which satisfy for all $n \in \mathbb{N}$, $x \in \mathbb{R}$ that*

$$g_1(x) = \begin{cases} 2x & : x \in [0, \frac{1}{2}) \\ 2 - 2x & : x \in [\frac{1}{2}, 1] \\ 0 & : x \in \mathbb{R} \setminus [0, 1] \end{cases} \quad (214)$$

and $g_{n+1}(x) = g_1(g_n(x))$. Then

(i) *it holds for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}]$ that*

$$g_n(x) = \begin{cases} 2^n(x - \frac{2k}{2^n}) & : x \in [\frac{2k}{2^n}, \frac{2k+1}{2^n}] \\ 2^n(\frac{2k+2}{2^n} - x) & : x \in [\frac{2k+1}{2^n}, \frac{2k+2}{2^n}] \end{cases} \quad (215)$$

and

(ii) *it holds for all $n \in \mathbb{N}$, $x \in \mathbb{R} \setminus [0, 1]$ that $g_n(x) = 0$.*

Proof of Lemma 3.1. First, we claim that for all $n \in \mathbb{N}$ it holds that

$$\left(\forall k \in \{0, 1, \dots, 2^{n-1} - 1\}, x \in [\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}] : \right. \\ \left. g_n(x) = \begin{cases} 2^n(x - \frac{2k}{2^n}) & : x \in [\frac{2k}{2^n}, \frac{2k+1}{2^n}] \\ 2^n(\frac{2k+2}{2^n} - x) & : x \in [\frac{2k+1}{2^n}, \frac{2k+2}{2^n}] \end{cases} \right). \quad (216)$$

We now prove (216) by induction on $n \in \mathbb{N}$. Note that (214) establishes (216) in the base case $n = 1$. For the induction step $\mathbb{N} \ni n \rightarrow n + 1 \in \mathbb{N} \cap [2, \infty)$ assume that there exists $n \in \mathbb{N}$ such that for all $k \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}]$ it holds that

$$g_n(x) = \begin{cases} 2^n(x - \frac{2k}{2^n}) & : x \in [\frac{2k}{2^n}, \frac{2k+1}{2^n}] \\ 2^n(\frac{2k+2}{2^n} - x) & : x \in [\frac{2k+1}{2^n}, \frac{2k+2}{2^n}] \end{cases}. \quad (217)$$

Observe that (214) and (217) imply that for all $l \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{2l}{2^n}, \frac{2l+(1/2)}{2^n}]$ it holds that

$$g_{n+1}(x) = g_1(g_n(x)) = g_1(2^n(x - \frac{2l}{2^n})) = 2[2^n(x - \frac{2l}{2^n})] = 2^{n+1}(x - \frac{2l}{2^n}). \quad (218)$$

In addition, note that (214) and (217) ensure that for all $l \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{2l+(1/2)}{2^n}, \frac{2l+1}{2^n}]$ it holds that

$$\begin{aligned} g_{n+1}(x) &= g_1(g_n(x)) = g_1(2^n(x - \frac{2l}{2^n})) = 2 - 2[2^n(x - \frac{2l}{2^n})] \\ &= 2 - 2^{n+1}x + 4l = 2^{n+1}(\frac{4l+2}{2^{n+1}} - x). \end{aligned} \quad (219)$$

Moreover, observe that (214) and (217) demonstrate that for all $l \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{2l+1}{2^n}, \frac{2l+(3/2)}{2^n}]$ it holds that

$$\begin{aligned} g_{n+1}(x) &= g_1(g_n(x)) = g_1(2^n(\frac{2l+2}{2^n} - x)) = 2 - 2[2^n(\frac{2l+2}{2^n} - x)] \\ &= 2 - 2(2l + 2) + 2^{n+1}x = 2^{n+1}x - 4l - 2 \\ &= 2^{n+1}(x - \frac{4l+2}{2^{n+1}}). \end{aligned} \quad (220)$$

Next note that (214) and (217) prove that for all $l \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in [\frac{2l+(3/2)}{2^n}, \frac{2l+2}{2^n}]$ it holds that

$$g_{n+1}(x) = g_1(g_n(x)) = g_1(2^n(\frac{2l+2}{2^n} - x)) = 2[2^n(\frac{2l+2}{2^n} - x)] = 2^{n+1}(\frac{2l+2}{2^n} - x). \quad (221)$$

Moreover, observe that for all $k \in \{0, 2, 4, 6, \dots\} \cap [0, 2^n - 2]$ it holds that

$$[\frac{2k}{2^{n+1}}, \frac{2k+1}{2^{n+1}}] = [\frac{2(k/2)}{2^n}, \frac{2(k/2)+(1/2)}{2^n}], \quad [\frac{2k+1}{2^{n+1}}, \frac{2k+2}{2^{n+1}}] = [\frac{2(k/2)+(1/2)}{2^n}, \frac{2(k/2)+1}{2^n}], \quad (222)$$

and $k/2 \in \{0, 1, \dots, 2^{n-1} - 1\}$. This, (218), and (219) demonstrate that for all $k \in \{0, 2, 4, 6, \dots\} \cap [0, 2^n - 2]$, $x \in [\frac{k}{2^n}, \frac{k+1}{2^n}]$ it holds that

$$\begin{aligned} g_{n+1}(x) &= \begin{cases} 2^{n+1}(x - \frac{2(k/2)}{2^n}) & : x \in [\frac{2(k/2)}{2^n}, \frac{2(k/2)+(1/2)}{2^n}] \\ 2^{n+1}(\frac{4(k/2)+2}{2^{n+1}} - x) & : x \in [\frac{2(k/2)+(1/2)}{2^n}, \frac{2(k/2)+1}{2^n}] \end{cases} \\ &= \begin{cases} 2^{n+1}(x - \frac{2k}{2^{n+1}}) & : x \in [\frac{2k}{2^{n+1}}, \frac{2k+1}{2^{n+1}}] \\ 2^{n+1}(\frac{2k+2}{2^{n+1}} - x) & : x \in [\frac{2k+1}{2^{n+1}}, \frac{2k+2}{2^{n+1}}] \end{cases}. \end{aligned} \quad (223)$$

In addition, observe that for all $k \in \{1, 3, 5, 7, \dots\} \cap [1, 2^n - 1]$ it holds that

$$\begin{aligned} \left[\frac{2k}{2^{n+1}}, \frac{2k+1}{2^{n+1}} \right] &= \left[\frac{2((k-1)/2)+1}{2^n}, \frac{2((k-1)/2)+(3/2)}{2^n} \right], \\ \left[\frac{2k+1}{2^{n+1}}, \frac{2k+2}{2^{n+1}} \right] &= \left[\frac{2((k-1)/2)+(3/2)}{2^n}, \frac{2((k-1)/2)+2}{2^n} \right], \end{aligned} \quad (224)$$

and $(k-1)/2 \in \{0, 1, \dots, 2^{n-1} - 1\}$. This, (220), and (221) demonstrate that for all $k \in \{1, 3, 5, 7, \dots\} \cap [1, 2^n - 1]$, $x \in \left[\frac{k}{2^n}, \frac{k+1}{2^n} \right]$ it holds that

$$\begin{aligned} g_{n+1}(x) &= \begin{cases} 2^{n+1} \left(x - \frac{4((k-1)/2)+2}{2^{n+1}} \right) & : x \in \left[\frac{2((k-1)/2)+1}{2^n}, \frac{2((k-1)/2)+(3/2)}{2^n} \right] \\ 2^{n+1} \left(\frac{2((k-1)/2)+2}{2^n} - x \right) & : x \in \left[\frac{2((k-1)/2)+(3/2)}{2^n}, \frac{2((k-1)/2)+2}{2^n} \right] \end{cases} \\ &= \begin{cases} 2^{n+1} \left(x - \frac{2k}{2^{n+1}} \right) & : x \in \left[\frac{2k}{2^{n+1}}, \frac{2k+1}{2^{n+1}} \right] \\ 2^{n+1} \left(\frac{2k+2}{2^{n+1}} - x \right) & : x \in \left[\frac{2k+1}{2^{n+1}}, \frac{2k+2}{2^{n+1}} \right] \end{cases}. \end{aligned} \quad (225)$$

Combining this with (223) ensures that for all $k \in \{0, 1, \dots, 2^n - 1\}$, $x \in \left[\frac{k}{2^n}, \frac{k+1}{2^n} \right]$ it holds that

$$g_{n+1}(x) = \begin{cases} 2^{n+1} \left(x - \frac{2k}{2^{n+1}} \right) & : x \in \left[\frac{2k}{2^{n+1}}, \frac{2k+1}{2^{n+1}} \right] \\ 2^{n+1} \left(\frac{2k+2}{2^{n+1}} - x \right) & : x \in \left[\frac{2k+1}{2^{n+1}}, \frac{2k+2}{2^{n+1}} \right] \end{cases}. \quad (226)$$

Induction thus proves (216). Observe that (216) establishes item (i). Next we claim that for all $n \in \mathbb{N}$ it holds that

$$\forall x \in \mathbb{R} \setminus [0, 1]: g_n(x) = 0. \quad (227)$$

We now prove (227) by induction on $n \in \mathbb{N}$. Note that (214) establishes (227) in the base case $n = 1$. For the induction step observe that (214) ensures that for all $n \in \mathbb{N}$ with $(\forall x \in \mathbb{R} \setminus [0, 1]: g_n(x) = 0)$ it holds that

$$(\forall x \in \mathbb{R} \setminus [0, 1]: g_{n+1}(x) = g_1(g_n(x)) = g_1(0) = 0). \quad (228)$$

Induction thus proves (227). Note that (227) establishes item (ii). The proof of Lemma 3.1 is thus completed. \square

Lemma 3.2. *Let $g_n: [0, 1] \rightarrow [0, 1]$, $n \in \mathbb{N}$, be the functions which satisfy for all $n \in \mathbb{N}$, $x \in [0, 1]$ that*

$$g_1(x) = \begin{cases} 2x & : x \in [0, \frac{1}{2}) \\ 2 - 2x & : x \in [\frac{1}{2}, 1] \end{cases} \quad (229)$$

and $g_{n+1}(x) = g_1(g_n(x))$, and let $f_n: [0, 1] \rightarrow [0, 1]$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $k \in \{0, 1, \dots, 2^n - 1\}$, $x \in \left[\frac{k}{2^n}, \frac{k+1}{2^n} \right)$ that $f_n(1) = 1$ and

$$f_n(x) = \left[\frac{2k+1}{2^n} \right] x - \frac{(k^2+k)}{2^{2n}}. \quad (230)$$

Then it holds for all $n \in \mathbb{N}_0$, $x \in [0, 1]$ that

$$f_n(x) = x - \left[\sum_{m=1}^n (2^{-2m} g_m(x)) \right] \quad \text{and} \quad |x^2 - f_n(x)| \leq 2^{-2n-2}. \quad (231)$$

Proof of Lemma 3.2. Note that (230) proves that for all $n \in \mathbb{N}_0$, $l \in \{0, 1, \dots, 2^n - 1\}$ it holds that

$$f_n\left(\frac{l}{2^n}\right) = \left[\frac{2l+1}{2^n} \right] \frac{l}{2^n} - \frac{(l^2+l)}{2^{2n}} = \frac{(2l+1)l - (l^2+l)}{2^{2n}} = \frac{l^2}{2^{2n}} = \left[\frac{l}{2^n} \right]^2. \quad (232)$$

The hypothesis that for all $n \in \mathbb{N}_0$ it holds that $f_n(1) = 1$ hence ensures that for all $n \in \mathbb{N}_0$, $l \in \{0, 1, \dots, 2^n\}$ it holds that

$$f_n\left(\frac{l}{2^n}\right) = \left[\frac{l}{2^n} \right]^2. \quad (233)$$

This and Lemma 3.1 demonstrate that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1}\}$ it holds that

$$\begin{aligned} f_{n-1}\left(\frac{2k}{2^n}\right) - f_n\left(\frac{2k}{2^n}\right) &= f_{n-1}\left(\frac{k}{2^{n-1}}\right) - f_n\left(\frac{2k}{2^n}\right) = \left[\frac{k}{2^{n-1}} \right]^2 - \left[\frac{2k}{2^n} \right]^2 \\ &= 0 = 2^{-2n} g_n\left(\frac{2k}{2^n}\right). \end{aligned} \quad (234)$$

In addition, note that (230) and (233) imply that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$ it holds that

$$\begin{aligned} f_{n-1}\left(\frac{2k+1}{2^n}\right) &= f_{n-1}\left(\frac{k+(1/2)}{2^{n-1}}\right) = \left[\frac{2k+1}{2^{n-1}} \right] \left[\frac{2k+1}{2^n} \right] - \frac{(k^2+k)}{2^{2(n-1)}} = \frac{(4k^2+4k+1)}{2^{2n-1}} - \frac{(2k^2+2k)}{2^{2n-1}} \\ &= \frac{2k^2+2k+1}{2^{2n-1}} = \frac{4k^2+4k+2}{2^{2n}} \end{aligned} \quad (235)$$

and

$$f_n\left(\frac{2k+1}{2^n}\right) = \left[\frac{2k+1}{2^n} \right]^2 = \frac{4k^2+4k+1}{2^{2n}}. \quad (236)$$

Lemma 3.1 hence assures that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$ it holds that

$$f_{n-1}\left(\frac{2k+1}{2^n}\right) - f_n\left(\frac{2k+1}{2^n}\right) = \frac{(4k^2+4k+2)}{2^{2n}} - \frac{(4k^2+4k+1)}{2^{2n}} = 2^{-2n} = 2^{-2n} g_n\left(\frac{2k+1}{2^n}\right). \quad (237)$$

Combining this with (234) shows that for all $n \in \mathbb{N}$, $l \in \{0, 1, \dots, 2^n\}$ it holds that

$$f_{n-1}\left(\frac{l}{2^n}\right) - f_n\left(\frac{l}{2^n}\right) = 2^{-2n} g_n\left(\frac{l}{2^n}\right). \quad (238)$$

Furthermore, observe that (233) demonstrates that for all $n \in \mathbb{N}_0$, $l \in \{0, 1, \dots, 2^n - 1\}$ it holds that

$$\left[\frac{2l+1}{2^n} \right] \left[\frac{l+1}{2^n} \right] - \frac{(l^2+l)}{2^{2n}} = \frac{(2l+1)(l+1) - l(l+1)}{2^{2n}} = \frac{(l+1)^2}{2^{2n}} = \left[\frac{l+1}{2^n} \right]^2 = f_n\left(\frac{l+1}{2^n}\right). \quad (239)$$

Combining this with (230) implies that for all $n \in \mathbb{N}_0$ it holds that $f_n \in C([0, 1], \mathbb{R})$ and

$$\forall l \in \{0, 1, \dots, 2^n - 1\}, x \in \left[\frac{l}{2^n}, \frac{l+1}{2^n}\right]: f_n(x) = \left[\frac{2l+1}{2^n}\right] x - \frac{(l^2+l)}{2^{2n}}. \quad (240)$$

The fact that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$ it holds that $\left[\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}\right] = \left[\frac{2k}{2^n}, \frac{2k+1}{2^n}\right] \cup \left[\frac{2k+1}{2^n}, \frac{2k+2}{2^n}\right]$ hence ensures that there exist $(a_{n,k}, b_{n,k}, c_{n,k}) \in \mathbb{R}^3$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$, $n \in \mathbb{N}$, such that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in \left[\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}\right]$ it holds that

$$f_{n-1}(x) - f_n(x) = \begin{cases} a_{n,k} \left(x - \frac{(2k+1)}{2^n}\right) + b_{n,k} & : x \in \left[\frac{2k}{2^n}, \frac{2k+1}{2^n}\right] \\ c_{n,k} \left(x - \frac{(2k+1)}{2^n}\right) + b_{n,k} & : x \in \left[\frac{2k+1}{2^n}, \frac{2k+2}{2^n}\right] \end{cases}. \quad (241)$$

Lemma 3.1 and (238) therefore prove that for all $n \in \mathbb{N}$, $k \in \{0, 1, \dots, 2^{n-1} - 1\}$, $x \in \left[\frac{k}{2^{n-1}}, \frac{k+1}{2^{n-1}}\right]$ it holds that

$$f_{n-1}(x) - f_n(x) = 2^{-2n} g_n(x). \quad (242)$$

Hence, we obtain that for all $n \in \mathbb{N}$, $x \in [0, 1]$ it holds that

$$f_{n-1}(x) - f_n(x) = 2^{-2n} g_n(x). \quad (243)$$

Next note that (230) ensures that for all $x \in [0, 1]$ it holds that $f_0(x) = x$. Combining this with (243) implies that for all $m \in \mathbb{N}_0$, $x \in [0, 1]$ it holds that

$$\begin{aligned} f_m(x) &= f_0(x) + \left[\sum_{n=1}^m (f_n(x) - f_{n-1}(x)) \right] \\ &= f_0(x) - \left[\sum_{n=1}^m (f_{n-1}(x) - f_n(x)) \right] = x - \left[\sum_{n=1}^m 2^{-2n} g_n(x) \right]. \end{aligned} \quad (244)$$

Moreover, observe that (240) demonstrates that for all $m \in \mathbb{N}_0$, $l \in \{0, 1, \dots, 2^m - 1\}$, $x \in \left[\frac{l}{2^m}, \frac{l+1}{2^m}\right]$ it holds that

$$\begin{aligned} f_m(x) - x^2 &= \left[\frac{2l+1}{2^m}\right] x - \frac{(l^2+l)}{2^{2m}} - x^2 = \left[\frac{l+1}{2^m}\right] x + \left[\frac{l}{2^m}\right] x - \left[\frac{l+1}{2^m}\right] \left[\frac{l}{2^m}\right] - x^2 \\ &= \left(x - \frac{l}{2^m}\right) \left(\frac{l+1}{2^m} - x\right) \geq 0. \end{aligned} \quad (245)$$

The fact that for all $a \in \mathbb{R}$, $b \in (a, \infty)$, $r \in [a, b]$ it holds that $(r-a)(b-r) \leq \frac{1}{4}(b-a)^2$ hence proves that for all $m \in \mathbb{N}_0$, $l \in \{0, 1, \dots, 2^m - 1\}$, $x \in \left[\frac{l}{2^m}, \frac{l+1}{2^m}\right]$ it holds that

$$\begin{aligned} |f_m(x) - x^2| &= f_m(x) - x^2 = \left(x - \frac{l}{2^m}\right) \left(\frac{l+1}{2^m} - x\right) \\ &\leq \frac{1}{4} \left(\frac{l+1}{2^m} - \frac{l}{2^m}\right)^2 = \frac{1}{4} \left(\frac{1}{2^m}\right)^2 = \frac{1}{2^2} \left(\frac{1}{2^{2m}}\right) = \frac{1}{2^{2m+2}} = 2^{-2m-2}. \end{aligned} \quad (246)$$

Therefore, we obtain that for all $m \in \mathbb{N}_0$, $x \in [0, 1]$ it holds that

$$|f_m(x) - x^2| \leq 2^{-2m-2}. \quad (247)$$

Combining this with (244) establishes (231). The proof of Lemma 3.2 is thus completed. \square

3.1.2 ANN approximations for the square function on $[0, 1]$

Proposition 3.3. *Let $\varepsilon \in (0, 1]$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$. Then there exists $\Phi \in \mathbf{N}$ such that*

- (i) *it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$,*
- (ii) *it holds for all $x \in \mathbb{R} \setminus [0, 1]$ that $(\mathcal{R}_a(\Phi))(x) = a(x)$,*
- (iii) *it holds for all $x \in [0, 1]$ that $|x^2 - (\mathcal{R}_a(\Phi))(x)| \leq \varepsilon$,*
- (iv) *it holds that $\mathcal{P}(\Phi) \leq \max\{10 \log_2(\varepsilon^{-1}) - 7, 13\}$, and*
- (v) *it holds that $\mathcal{L}(\Phi) \leq \max\{\frac{1}{2} \log_2(\varepsilon^{-1}) + 1, 2\}$*

(cf. Definition 2.1 and Definition 2.3).

Proof of Proposition 3.3. Throughout this proof let $M \in \mathbb{N}$ satisfy that

$$M = \min\left(\mathbb{N} \cap [2, \infty) \cap \left[\frac{1}{2} \log_2(\varepsilon^{-1}), \infty\right)\right), \quad (248)$$

let $g_n: \mathbb{R} \rightarrow [0, 1]$, $n \in \mathbb{N}$, be the functions which satisfy for all $n \in \mathbb{N}$, $x \in \mathbb{R}$ that

$$g_1(x) = \begin{cases} 2x & : x \in [0, \frac{1}{2}) \\ 2 - 2x & : x \in [\frac{1}{2}, 1] \\ 0 & : x \in \mathbb{R} \setminus [0, 1] \end{cases} \quad (249)$$

and $g_{n+1}(x) = g_1(g_n(x))$, let $f_n: [0, 1] \rightarrow [0, 1]$, $n \in \mathbb{N}_0$, be the functions which satisfy for all $n \in \mathbb{N}_0$, $k \in \{0, 1, \dots, 2^n - 1\}$, $x \in [\frac{k}{2^n}, \frac{k+1}{2^n})$ that $f_n(1) = 1$ and

$$f_n(x) = \left[\frac{2k+1}{2^n}\right] x - \frac{(k^2+k)}{2^{2n}}, \quad (250)$$

let $(A_k, b_k) \in \mathbb{R}^{4 \times 4} \times \mathbb{R}^4$, $k \in \mathbb{N} \cap [2, \infty)$, satisfy for all $k \in \mathbb{N} \cap [2, \infty)$ that

$$A_k = \begin{pmatrix} 2 & -4 & 2 & 0 \\ 2 & -4 & 2 & 0 \\ 2 & -4 & 2 & 0 \\ (-2)^{3-2k} & 2^{4-2k} & (-2)^{3-2k} & 1 \end{pmatrix} \quad \text{and} \quad b_k = \begin{pmatrix} 0 \\ -\frac{1}{2} \\ -1 \\ 0 \end{pmatrix}, \quad (251)$$

let $\mathbb{A}_k \in \mathbb{R}^{1 \times 4} \times \mathbb{R}$, $k \in \mathbb{N} \cap [2, \infty)$, satisfy for all $k \in \mathbb{N} \cap [2, \infty)$ that

$$\mathbb{A}_k = \left(\left((-2)^{3-2k} \quad 2^{4-2k} \quad (-2)^{3-2k} \quad 1 \right), 0 \right), \quad (252)$$

let $\phi_k \in \mathbf{N}$, $k \in \mathbb{N} \cap [2, \infty)$, satisfy that

$$\phi_2 = \left(\left(\left(\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ -\frac{1}{2} \\ -1 \\ 0 \end{pmatrix} \right), \mathbb{A}_2 \right) \right) \quad (253)$$

and

$$\forall k \in \mathbb{N} \cap [3, \infty): \phi_k = \left(\left(\left(\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ -\frac{1}{2} \\ -1 \\ 0 \end{pmatrix} \right), (A_2, b_2), \dots, (A_{k-1}, b_{k-1}), \mathbb{A}_k \right), \quad (254)$$

and let $r_k = (r_{k,1}, r_{k,2}, r_{k,3}, r_{k,4}): \mathbb{R} \rightarrow \mathbb{R}^4$, $k \in \mathbb{N}$, be the functions which satisfy for all $x \in \mathbb{R}$, $k \in \mathbb{N}$ that

$$r_1(x) = (r_{1,1}(x), r_{1,2}(x), r_{1,3}(x), r_{1,4}(x)) = \mathfrak{M}_{a,4}(x, x - \frac{1}{2}, x - 1, x) \quad (255)$$

and

$$r_{k+1}(x) = (r_{k+1,1}(x), r_{k+1,2}(x), r_{k+1,3}(x), r_{k+1,4}(x)) = \mathfrak{M}_{a,4}(A_{k+1}r_k(x) + b_{k+1}) \quad (256)$$

(cf. Definition 2.2). Note that (255), (6), (249), and the hypothesis that for all $x \in \mathbb{R}$ it holds that $a(x) = \max\{x, 0\}$ show that for all $x \in \mathbb{R}$ it holds that

$$\begin{aligned} 2r_{1,1}(x) - 4r_{1,2}(x) + 2r_{1,3}(x) &= 2a(x) - 4a(x - \frac{1}{2}) + 2a(x - 1) \\ &= 2 \max\{x, 0\} - 4 \max\{x - \frac{1}{2}, 0\} + 2 \max\{x - 1, 0\} = g_1(x). \end{aligned} \quad (257)$$

Furthermore, observe that (255), (6), the hypothesis that for all $x \in \mathbb{R}$ it holds that $a(x) = \max\{x, 0\}$, and the fact that for all $x \in [0, 1]$ it holds that $f_0(x) = x = \max\{x, 0\}$ imply that for all $x \in \mathbb{R}$ it holds that

$$r_{1,4}(x) = \max\{x, 0\} = \begin{cases} f_0(x) & : x \in [0, 1] \\ \max\{x, 0\} & : x \in \mathbb{R} \setminus [0, 1] \end{cases}. \quad (258)$$

Next we claim that for all $k \in \mathbb{N}$ it holds that

$$(\forall x \in \mathbb{R}: 2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x) = g_k(x)) \quad (259)$$

and

$$\left(\forall x \in \mathbb{R}: r_{k,4}(x) = \begin{cases} f_{k-1}(x) & : x \in [0, 1] \\ \max\{x, 0\} & : x \in \mathbb{R} \setminus [0, 1] \end{cases} \right). \quad (260)$$

We now prove (259)–(260) by induction on $k \in \mathbb{N}$. Note that (257) and (258) prove (259)–(260) in the base case $k = 1$. For the induction step $\mathbb{N} \ni k \rightarrow k + 1 \in \mathbb{N} \cap [2, \infty)$ assume that there exists $k \in \mathbb{N}$ such that for all $x \in \mathbb{R}$ it holds that

$$2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x) = g_k(x) \quad (261)$$

$$\text{and} \quad r_{k,4}(x) = \begin{cases} f_{k-1}(x) & : x \in [0, 1] \\ \max\{x, 0\} & : x \in \mathbb{R} \setminus [0, 1] \end{cases}. \quad (262)$$

Observe that (261), (257), (251), (6), and (256) ensure that for all $x \in \mathbb{R}$ it holds that

$$\begin{aligned} g_{k+1}(x) &= g_1(g_k(x)) = g_1(2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x)) \\ &= 2a(2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x)) \\ &\quad - 4a(2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x) - \frac{1}{2}) \\ &\quad + 2a(2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x) - 1) \\ &= 2r_{k+1,1}(x) - 4r_{k+1,2}(x) + 2r_{k+1,3}(x). \end{aligned} \quad (263)$$

In addition, observe that (6), (251), (256), and (261) demonstrate that for all $x \in \mathbb{R}$ it holds that

$$\begin{aligned} r_{k+1,4}(x) &= a((-2)^{3-2(k+1)}r_{k,1}(x) + 2^{4-2(k+1)}r_{k,2}(x) + (-2)^{3-2(k+1)}r_{k,3}(x) + r_{k,4}(x)) \\ &= a((-2)^{1-2k}r_{k,1}(x) + 2^{2-2k}r_{k,2}(x) + (-2)^{1-2k}r_{k,3}(x) + r_{k,4}(x)) \\ &= a(2^{-2k}[-2r_{k,1}(x) + 2^2r_{k,2}(x) - 2r_{k,3}(x)] + r_{k,4}(x)) \\ &= a(-[2^{-2k}][2r_{k,1}(x) - 4r_{k,2}(x) + 2r_{k,3}(x)] + r_{k,4}(x)) \\ &= a(-[2^{-2k}]g_k(x) + r_{k,4}(x)). \end{aligned} \quad (264)$$

Combining this with (262), Lemma 3.2, the hypothesis that for all $x \in \mathbb{R}$ it holds that $a(x) = \max\{x, 0\}$, and the fact that for all $x \in [0, 1]$ it holds that $f_k(x) \geq 0$ shows that for all $x \in [0, 1]$ it holds that

$$\begin{aligned} r_{k+1,4}(x) &= a(-[2^{-2k}g_k(x)] + f_{k-1}(x)) \\ &= a\left(- (2^{-2k}g_k(x)) + x - \left[\sum_{j=1}^{k-1} (2^{-2j}g_j(x))\right]\right) \\ &= a\left(x - \left[\sum_{j=1}^k 2^{-2j}g_j(x)\right]\right) = a(f_k(x)) = f_k(x). \end{aligned} \quad (265)$$

Next note that (264), (262), item (ii) in Lemma 3.1, and the hypothesis that for all $x \in \mathbb{R}$ it holds that $a(x) = \max\{x, 0\}$ prove that for all $x \in \mathbb{R} \setminus [0, 1]$ it holds that

$$r_{k+1,4}(x) = a\left(- (2^{-2k}g_k(x)) + r_{k,4}(x)\right) = a(\max\{x, 0\}) = \max\{x, 0\}. \quad (266)$$

Combining (263) and (265) hence proves (259)–(260) in the case $k + 1$. Induction thus establishes (259)–(260). Next note that (7), (251), (252), (259), (253), (254), (255), and (256) assure that for all $m \in \mathbb{N} \cap [2, \infty)$, $x \in \mathbb{R}$ it holds that $\mathcal{R}_a(\phi_m) \in C(\mathbb{R}, \mathbb{R})$ and

$$\begin{aligned}
& (\mathcal{R}_a(\phi_m))(x) \\
&= (-2)^{3-2m}r_{m-1,1}(x) + 2^{4-2m}r_{m-1,2}(x) + (-2)^{3-2m}r_{m-1,3}(x) + r_{m-1,4}(x) \\
&= (-2)^{4-2m} \left(\left[\frac{r_{m-1,1}(x)+r_{m-1,3}(x)}{(-2)} \right] + r_{m-1,2}(x) \right) + r_{m-1,4}(x) \\
&= 2^{4-2m} \left(\left[\frac{r_{m-1,1}(x)+r_{m-1,3}(x)}{(-2)} \right] + r_{m-1,2}(x) \right) + r_{m-1,4}(x) \tag{267} \\
&= 2^{2-2m} (4r_{m-1,2}(x) - 2r_{m-1,1}(x) - 2r_{m-1,3}(x)) + r_{m-1,4}(x) \\
&= -[2^{-2(m-1)}] [2r_{m-1,1}(x) - 4r_{m-1,2}(x) + 2r_{m-1,3}(x)] + r_{m-1,4}(x) \\
&= -[2^{-2(m-1)}] g_{m-1}(x) + r_{m-1,4}(x).
\end{aligned}$$

Combining this with (260) and Lemma 3.2 shows that for all $m \in \mathbb{N} \cap [2, \infty)$, $x \in [0, 1]$ it holds that

$$\begin{aligned}
(\mathcal{R}_a(\phi_m))(x) &= -(2^{-2(m-1)}g_{m-1}(x)) + f_{m-2}(x) \\
&= -(2^{-2(m-1)}g_{m-1}(x)) + x - \left[\sum_{j=1}^{m-2} 2^{-2j} g_j(x) \right] \tag{268} \\
&= x - \left[\sum_{j=1}^{m-1} 2^{-2j} g_j(x) \right] = f_{m-1}(x).
\end{aligned}$$

Lemma 3.2 therefore implies that for all $m \in \mathbb{N} \cap [2, \infty)$, $x \in [0, 1]$ it holds that

$$|x^2 - (\mathcal{R}_a(\phi_m))(x)| \leq 2^{-2(m-1)-2} = 2^{-2m}. \tag{269}$$

Next note that (248) assures that

$$\begin{aligned}
M &= \min \left(\mathbb{N} \cap \left[\max \left\{ 2, \frac{1}{2} \log_2(\varepsilon^{-1}) \right\}, \infty \right) \right) \\
&\geq \min \left(\left[\max \left\{ 2, \frac{1}{2} \log_2(\varepsilon^{-1}) \right\}, \infty \right) \right) \tag{270} \\
&= \max \left\{ 2, \frac{1}{2} \log_2(\varepsilon^{-1}) \right\} \geq \frac{1}{2} \log_2(\varepsilon^{-1}).
\end{aligned}$$

This and (269) demonstrate that for all $x \in [0, 1]$ it holds that

$$|x^2 - (\mathcal{R}_a(\phi_M))(x)| \leq 2^{-2M} \leq 2^{-\log_2(\varepsilon^{-1})} = \varepsilon. \tag{271}$$

Moreover, observe that item (ii) in Lemma 3.1, (260), and (267) ensure that for all $m \in \mathbb{N} \cap [2, \infty)$, $x \in \mathbb{R} \setminus [0, 1]$ it holds that

$$\begin{aligned}
(\mathcal{R}_a(\phi_m))(x) &= -2^{-2(m-1)}g_{m-1}(x) + r_{m-1,4}(x) \\
&= r_{m-1,4}(x) = \max\{x, 0\} = a(x). \tag{272}
\end{aligned}$$

Furthermore, observe that (248), (253), and (254) assure that

$$\mathcal{L}(\phi_M) = M \leq \max\{\frac{1}{2}\log_2(\varepsilon^{-1}) + 1, 2\}. \quad (273)$$

This, (248), (253), and (254) show that

$$\begin{aligned} \mathcal{P}(\phi_M) &= 4(1+1) + \left[\sum_{j=2}^{M-1} 4(4+1) \right] + (4+1) \\ &= 8 + 20(M-2) + 5 \leq 20 \max\{\frac{1}{2}\log_2(\varepsilon^{-1}) - 1, 0\} + 13 \\ &= \max\{10\log_2(\varepsilon^{-1}) - 20, 0\} + 13 = \max\{10\log_2(\varepsilon^{-1}) - 7, 13\}. \end{aligned} \quad (274)$$

Combining (271), (273), (272), and the fact that $\mathcal{R}_a(\phi_M) \in C(\mathbb{R}, \mathbb{R})$ hence establishes items (i)–(v). The proof of Proposition 3.3 is thus completed. \square

3.1.3 ANN approximations for the square function on \mathbb{R}

Proposition 3.4. *Let $\varepsilon \in (0, 1]$, $q \in (2, \infty)$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$. Then there exists $\Phi \in \mathbf{N}$ such that*

- (i) *it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$,*
- (ii) *it holds that $(\mathcal{R}_a(\Phi))(0) = 0$,*
- (iii) *it holds for all $x \in \mathbb{R}$ that $0 \leq (\mathcal{R}_a(\Phi))(x) \leq \varepsilon + |x|^2$,*
- (iv) *it holds for all $x \in \mathbb{R}$ that $|x^2 - (\mathcal{R}_a(\Phi))(x)| \leq \varepsilon \max\{1, |x|^q\}$,*
- (v) *it holds that $\mathcal{P}(\Phi) \leq \max\{\lfloor \frac{40q}{(q-2)} \rfloor \log_2(\varepsilon^{-1}) + \frac{80}{(q-2)} - 28, 52\}$, and*
- (vi) *it holds that $\mathcal{L}(\Phi) \leq \max\{\frac{q}{2(q-2)} \log_2(\varepsilon^{-1}) + \frac{1}{(q-2)} + 1, 2\}$*

(cf. Definition 2.1 and Definition 2.3).

Proof of Proposition 3.4. Throughout this proof let $\delta \in (0, 1]$ satisfy that $\delta = 2^{-2/(q-2)}\varepsilon^{q/(q-2)}$, let $\mathbb{A}_1 \in (\mathbb{R}^{2 \times 1} \times \mathbb{R}^2) \subseteq \mathbf{N}$, $\mathbb{A}_2 \in (\mathbb{R}^{1 \times 2} \times \mathbb{R}) \subseteq \mathbf{N}$ satisfy that

$$\mathbb{A}_1 = \left(\begin{pmatrix} (\frac{\varepsilon}{2})^{1/(q-2)} \\ -(\frac{\varepsilon}{2})^{1/(q-2)} \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \quad \text{and} \quad \mathbb{A}_2 = \left(\left((\frac{\varepsilon}{2})^{-2/(q-2)} \quad (\frac{\varepsilon}{2})^{-2/(q-2)} \right), 0 \right), \quad (275)$$

let $\Psi \in \mathbf{N}$ satisfy that

- (I) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}, \mathbb{R})$,*
- (II) *it holds for all $x \in \mathbb{R} \setminus [0, 1]$ that $(\mathcal{R}_a(\Psi))(x) = a(x)$,*

(III) it holds for all $x \in [0, 1]$ that $|x^2 - (\mathcal{R}_a(\Psi))(x)| \leq \delta$,

(IV) it holds that $\mathcal{P}(\Psi) \leq \max\{10 \log_2(\delta^{-1}) - 7, 13\}$, and

(V) it holds that $\mathcal{L}(\Psi) \leq \max\{\frac{1}{2} \log_2(\delta^{-1}) + 1, 2\}$

(cf. Proposition 3.3), and let $\Phi \in \mathbf{N}$ satisfy that

$$\Phi = \mathbb{A}_2 \bullet [\mathbf{P}_2(\Psi, \Psi)] \bullet \mathbb{A}_1 \quad (276)$$

(cf. Definition 2.5, Definition 2.17, and Lemma 2.8). Note that Proposition 2.19 and item (v) in Proposition 2.6 ensure that for all $x \in \mathbb{R}$ it holds that

$$\begin{aligned} (\mathcal{R}_a((\mathbf{P}_2(\Psi, \Psi)) \bullet \mathbb{A}_1))(x) &= (\mathcal{R}_a(\mathbf{P}_2(\Psi, \Psi)))(\mathcal{R}_a(\mathbb{A}_1))(x) \\ &= (\mathcal{R}_a(\mathbf{P}_2(\Psi, \Psi)))(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x, -\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x) \\ &= \begin{pmatrix} (\mathcal{R}_a(\Psi))\left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \\ (\mathcal{R}_a(\Psi))\left(-\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \end{pmatrix}. \end{aligned} \quad (277)$$

Item (v) in Proposition 2.6 and (276) therefore demonstrate that for all $x \in \mathbb{R}$ it holds that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(x) &= (\mathcal{R}_a(\mathbb{A}_2))(\mathcal{R}_a([\mathbf{P}_2(\Psi, \Psi)] \bullet \mathbb{A}_1))(x) \\ &= \begin{pmatrix} \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} & \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \end{pmatrix} \begin{pmatrix} [\mathcal{R}_a(\Psi)]\left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \\ [\mathcal{R}_a(\Psi)]\left(-\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \end{pmatrix} \\ &= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left([\mathcal{R}_a(\Psi)]\left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) + [\mathcal{R}_a(\Psi)]\left(-\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \right). \end{aligned} \quad (278)$$

This, (I), (II), and the hypothesis that for all $x \in \mathbb{R}$ it holds that $a(x) = \max\{x, 0\}$ imply that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(0) &= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left([\mathcal{R}_a(\Psi)](0) + [\mathcal{R}_a(\Psi)](0) \right) \\ &= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} (a(0) + a(0)) = 0. \end{aligned} \quad (279)$$

Moreover, observe that (I) and (II) ensure that for all $x \in \mathbb{R} \setminus [-1, 1]$ it holds that

$$\begin{aligned} [\mathcal{R}_a(\Psi)](x) + [\mathcal{R}_a(\Psi)](-x) &= a(x) + a(-x) = \max\{x, 0\} + \max\{-x, 0\} \\ &= \max\{x, 0\} - \min\{x, 0\} = |x|. \end{aligned} \quad (280)$$

Furthermore, note that (II) and (III) show that

$$\begin{aligned}
& \sup_{x \in [-1,1]} |x^2 - ([\mathcal{R}_a(\Psi)](x) + [\mathcal{R}_a(\Psi)](-x))| \\
&= \max \left\{ \sup_{x \in [-1,0]} |x^2 - (a(x) + [\mathcal{R}_a(\Psi)](-x))|, \sup_{x \in [0,1]} |x^2 - ([\mathcal{R}_a(\Psi)](x) + a(-x))| \right\} \\
&= \max \left\{ \sup_{x \in [-1,0]} |(-x)^2 - (\mathcal{R}_a(\Psi))(-x)|, \sup_{x \in [0,1]} |x^2 - (\mathcal{R}_a(\Psi))(x)| \right\} \\
&= \sup_{x \in [0,1]} |x^2 - (\mathcal{R}_a(\Psi))(x)| \leq \delta.
\end{aligned} \tag{281}$$

Next observe that (278) and (280) prove that for all $x \in \mathbb{R} \setminus [-(\varepsilon/2)^{-1/(q-2)}, (\varepsilon/2)^{-1/(q-2)}]$ it holds that

$$\begin{aligned}
0 &\leq [\mathcal{R}_a(\Phi)](x) \\
&= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left([\mathcal{R}_a(\Psi)]\left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) + [\mathcal{R}_a(\Psi)]\left(-\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \right) \\
&= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left| \left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x \right| = \left(\frac{\varepsilon}{2}\right)^{-1/(q-2)} |x| \leq |x|^2.
\end{aligned} \tag{282}$$

The triangle inequality therefore ensures that for all $x \in \mathbb{R} \setminus [-(\varepsilon/2)^{-1/(q-2)}, (\varepsilon/2)^{-1/(q-2)}]$ it holds that

$$\begin{aligned}
|x^2 - (\mathcal{R}_a(\Phi))(x)| &= |x^2 - \left(\frac{\varepsilon}{2}\right)^{-1/(q-2)} |x|| \leq (|x|^2 + \left(\frac{\varepsilon}{2}\right)^{-1/(q-2)} |x|) \\
&= (|x|^q |x|^{-(q-2)} + \left(\frac{\varepsilon}{2}\right)^{-1/(q-2)} |x|^q |x|^{-(q-1)}) \\
&\leq (|x|^q \left(\frac{\varepsilon}{2}\right)^{(q-2)/(q-2)} + \left(\frac{\varepsilon}{2}\right)^{-1/(q-2)} |x|^q \left(\frac{\varepsilon}{2}\right)^{(q-1)/(q-2)}) \\
&= \left(\frac{\varepsilon}{2} + \frac{\varepsilon}{2}\right) |x|^q = \varepsilon |x|^q \leq \varepsilon \max\{1, |x|^q\}.
\end{aligned} \tag{283}$$

Next note that (278), (281), and the fact that $\delta = 2^{-2/(q-2)} \varepsilon^{q/(q-2)}$ demonstrate that for all $x \in [-(\varepsilon/2)^{-1/(q-2)}, (\varepsilon/2)^{-1/(q-2)}]$ it holds that

$$\begin{aligned}
& |x^2 - (\mathcal{R}_a(\Phi))(x)| \\
&= \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left| \left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right)^2 - \left([\mathcal{R}_a(\Psi)]\left(\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) + [\mathcal{R}_a(\Psi)]\left(-\left(\frac{\varepsilon}{2}\right)^{1/(q-2)}x\right) \right) \right| \\
&\leq \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \left[\sup_{y \in [-1,1]} |y^2 - ([\mathcal{R}_a(\Psi)](y) + [\mathcal{R}_a(\Psi)](-y))| \right] \\
&\leq \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} \delta = \left(\frac{\varepsilon}{2}\right)^{-2/(q-2)} 2^{-2/(q-2)} \varepsilon^{q/(q-2)} = \varepsilon \leq \varepsilon \max\{1, |x|^q\}.
\end{aligned} \tag{284}$$

Combining this and (283) implies that for all $x \in \mathbb{R}$ it holds that

$$|x^2 - (\mathcal{R}_a(\Phi))(x)| \leq \varepsilon \max\{1, |x|^q\} \leq \varepsilon(1 + |x|^q). \tag{285}$$

In addition, note that (284) ensures that for all $x \in [-(\varepsilon/2)^{-1/(q-2)}, (\varepsilon/2)^{-1/(q-2)}]$ it holds that

$$|(\mathcal{R}_a(\Phi))(x)| \leq |x^2 - (\mathcal{R}_a(\Phi))(x)| + |x|^2 \leq \varepsilon + |x|^2. \quad (286)$$

This and (282) show for all $x \in \mathbb{R}$ that

$$|(\mathcal{R}_a(\Phi))(x)| \leq \varepsilon + |x|^2. \quad (287)$$

Furthermore, observe that the fact that $\delta = 2^{-2/(q-2)}\varepsilon^{q/(q-2)}$ ensures that

$$\log_2(\delta^{-1}) = \log_2(2^{2/(q-2)}\varepsilon^{-q/(q-2)}) = \frac{2}{(q-2)} + \left\lceil \frac{q}{(q-2)} \right\rceil \log_2(\varepsilon^{-1}). \quad (288)$$

Next note that Corollary 2.21 implies that $\mathcal{P}(\mathbf{P}_2(\Psi, \Psi)) \leq 4\mathcal{P}(\Psi)$. Corollary 2.9, (276), (IV), and (288) hence ensure that

$$\begin{aligned} \mathcal{P}(\Phi) &\leq \left[\max \left\{ 1, \frac{\mathcal{O}(\mathbb{A}_2)}{\mathcal{O}(\mathbf{P}_2(\Psi, \Psi))} \right\} \right] \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A}_1)+1}{\mathcal{I}(\mathbf{P}_2(\Psi, \Psi))+1} \right\} \right] \mathcal{P}(\mathbf{P}_2(\Psi, \Psi)) \\ &= \left[\max \left\{ 1, \frac{1}{2} \right\} \right] \left[\max \left\{ 1, \frac{2}{3} \right\} \right] \mathcal{P}(\mathbf{P}_2(\Psi, \Psi)) \\ &= \mathcal{P}(\mathbf{P}_2(\Psi, \Psi)) \leq 4\mathcal{P}(\Psi) \leq 4 \max \{ 10 \log_2(\delta^{-1}) - 7, 13 \} \\ &= \max \left\{ 40 \left\lceil \frac{2}{(q-2)} \right\rceil + 40 \left\lceil \frac{q}{(q-2)} \right\rceil \log_2(\varepsilon^{-1}) - 28, 52 \right\} \\ &= \max \left\{ \left\lceil \frac{40q}{(q-2)} \right\rceil \log_2(\varepsilon^{-1}) + \frac{80}{(q-2)} - 28, 52 \right\}. \end{aligned} \quad (289)$$

In addition, observe that item (ii) in Proposition 2.6, (276), (V), and (288) demonstrate that

$$\begin{aligned} \mathcal{L}(\Phi) &= \mathcal{L}(\mathbf{P}_2(\Psi, \Psi)) = \mathcal{L}(\Psi) \leq \max \left\{ \frac{1}{2} \log_2(\delta^{-1}) + 1, 2 \right\} \\ &= \max \left\{ \left\lceil \frac{q}{2(q-2)} \right\rceil \log_2(\varepsilon^{-1}) + \frac{1}{(q-2)} + 1, 2 \right\}. \end{aligned} \quad (290)$$

Combining this with (279), (282), (287), (285), (289) establishes items (i)–(vi). The proof of Proposition 3.4 is thus completed. \square

3.2 ANN approximations for products

3.2.1 ANN approximations for one-dimensional products

Proposition 3.5. *Let $\varepsilon \in (0, 1]$, $q \in (2, \infty)$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$. Then there exists $\Phi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^2, \mathbb{R})$,*

(ii) *it holds for all $x \in \mathbb{R}$ that $(\mathcal{R}_a(\Phi))(x, 0) = (\mathcal{R}_a(\Phi))(0, x) = 0$,*

(iii) *it holds for all $x, y \in \mathbb{R}$ that*

$$|xy - (\mathcal{R}_a(\Phi))(x, y)| \leq \varepsilon \max\{1, |x|^q, |y|^q\}, \quad (291)$$

(iv) it holds for all $x, y \in \mathbb{R}$ that

$$|(\mathcal{R}_a(\Phi))(x, y)| \leq \frac{3}{2} \left(\frac{\varepsilon}{3} + x^2 + y^2 \right) \leq 1 + 2x^2 + 2y^2, \quad (292)$$

(v) it holds that

$$\begin{aligned} \mathcal{P}(\Phi) &\leq \frac{360q}{(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1) \right] + \frac{1}{(q-2)} - 252 \\ &\leq \frac{360q}{(q-2)} \left[\log_2(\varepsilon^{-1}) + q + 1 \right] - 252, \end{aligned} \quad (293)$$

and

(vi) it holds that

$$\begin{aligned} \mathcal{L}(\Phi) &\leq \frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1) \right] + \frac{(q-1)}{(q-2)} \\ &\leq \frac{q}{(q-2)} \left[\log_2(\varepsilon^{-1}) + q \right] \end{aligned} \quad (294)$$

(cf. Definition 2.1 and Definition 2.3).

Proof of Proposition 3.5. Throughout this proof let $\delta \in (0, 1]$ satisfy that $\delta = \varepsilon(2^{q-1} + 1)^{-1}$, let $\mathbb{A}_1 \in (\mathbb{R}^{3 \times 2} \times \mathbb{R}^3) \subseteq \mathbf{N}$, $\mathbb{A}_2 \in (\mathbb{R}^{1 \times 3} \times \mathbb{R}) \subseteq \mathbf{N}$ satisfy that

$$\mathbb{A}_1 = \left(\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right) \quad \text{and} \quad \mathbb{A}_2 = \left(\left(\frac{1}{2} \quad -\frac{1}{2} \quad -\frac{1}{2} \right), 0 \right), \quad (295)$$

let $\Psi \in \mathbf{N}$ satisfy that

- (I) it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}, \mathbb{R})$,
- (II) it holds that $[\mathcal{R}_a(\Psi)](0) = 0$,
- (III) it holds for all $x \in \mathbb{R}$ that $0 \leq [\mathcal{R}_a(\Psi)](x) \leq \delta + |x|^2$,
- (IV) it holds for all $x \in \mathbb{R}$ that $|x^2 - [\mathcal{R}_a(\Psi)](x)| \leq \delta \max\{1, |x|^q\}$,
- (V) it holds that $\mathcal{P}(\Psi) \leq \max\left\{ \left[\frac{40q}{(q-2)} \right] \log_2(\delta^{-1}) + \frac{80}{(q-2)} - 28, 52 \right\}$, and
- (VI) it holds that $\mathcal{L}(\Psi) \leq \max\left\{ \left[\frac{q}{2(q-2)} \right] \log_2(\delta^{-1}) + \frac{1}{(q-2)} + 1, 2 \right\}$

(cf. Proposition 3.4), and let $\Phi \in \mathbf{N}$ satisfy that

$$\Phi = \mathbb{A}_2 \bullet [\mathbf{P}_3(\Psi, \Psi, \Psi)] \bullet \mathbb{A}_1 \quad (296)$$

(cf. Definition 2.5, Definition 2.17, and Lemma 2.8). Note that item (v) in Proposition 2.6 and Proposition 2.19 ensure that for all $x, y \in \mathbb{R}$ it holds that $\mathcal{R}_a([\mathbf{P}_3(\Psi, \Psi, \Psi)] \bullet \mathbb{A}_1) \in C(\mathbb{R}^2, \mathbb{R}^3)$ and

$$\begin{aligned} [\mathcal{R}_a([\mathbf{P}_3(\Psi, \Psi, \Psi)] \bullet \mathbb{A}_1)](x, y) &= [\mathcal{R}_a(\mathbf{P}_3(\Psi, \Psi, \Psi))]([\mathcal{R}_a(\mathbb{A}_1)](x, y)) \\ &= [\mathcal{R}_a(\mathbf{P}_3(\Psi, \Psi, \Psi))](x + y, x, y) = \begin{pmatrix} [\mathcal{R}_a(\Psi)](x + y) \\ [\mathcal{R}_a(\Psi)](x) \\ [\mathcal{R}_a(\Psi)](y) \end{pmatrix}. \end{aligned} \quad (297)$$

Item (v) in Proposition 2.6 and (296) therefore demonstrate that for all $x, y \in \mathbb{R}$ it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^2, \mathbb{R})$ and

$$\begin{aligned} [\mathcal{R}_a(\Phi)](x, y) &= (\mathcal{R}_a(\mathbb{A}_2 \bullet [\mathbf{P}_3(\Psi, \Psi, \Psi)] \bullet \mathbb{A}_1))(x, y) \\ &= [\mathcal{R}_a(\mathbb{A}_2)](\mathcal{R}_a([\mathbf{P}_3(\Psi, \Psi, \Psi)] \bullet \mathbb{A}_1)(x, y)) \\ &= \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} [\mathcal{R}_a(\Psi)](x + y) \\ [\mathcal{R}_a(\Psi)](x) \\ [\mathcal{R}_a(\Psi)](y) \end{pmatrix} \\ &= \frac{1}{2}[\mathcal{R}_a(\Psi)](x + y) - \frac{1}{2}[\mathcal{R}_a(\Psi)](x) - \frac{1}{2}[\mathcal{R}_a(\Psi)](y). \end{aligned} \quad (298)$$

The fact that for all $\alpha, \beta \in \mathbb{R}$ it holds that $\alpha\beta = \frac{1}{2}|\alpha + \beta|^2 - \frac{1}{2}|\alpha|^2 - \frac{1}{2}|\beta|^2$, the triangle inequality, and (IV) hence ensure that for all $x, y \in \mathbb{R}$ it holds that

$$\begin{aligned} &|[\mathcal{R}_a(\Phi)](x, y) - xy| \\ &= \left| \frac{1}{2}[[\mathcal{R}_a(\Psi)](x + y) - |x + y|^2] - \frac{1}{2}[[\mathcal{R}_a(\Psi)](x) - |x|^2] - \frac{1}{2}[[\mathcal{R}_a(\Psi)](y) - |y|^2] \right| \\ &\leq \frac{1}{2} |[\mathcal{R}_a(\Psi)](x + y) - |x + y|^2| + \frac{1}{2} |[\mathcal{R}_a(\Psi)](x) - |x|^2| + \frac{1}{2} |[\mathcal{R}_a(\Psi)](y) - |y|^2| \\ &\leq \frac{\delta}{2} [\max\{1, |x + y|^q\} + \max\{1, |x|^q\} + \max\{1, |y|^q\}]. \end{aligned} \quad (299)$$

This, the fact that for all $\alpha, \beta \in \mathbb{R}$, $p \in [1, \infty)$ it holds that $|\alpha + \beta|^p \leq 2^{p-1}(|\alpha|^p + |\beta|^p)$, and the fact that $\delta = \varepsilon(2^{q-1} + 1)^{-1}$ establish that for all $x, y \in \mathbb{R}$ it holds that

$$\begin{aligned} &|[\mathcal{R}_a(\Phi)](x, y) - xy| \\ &\leq \frac{\delta}{2} [\max\{1, 2^{q-1}|x|^q + 2^{q-1}|y|^q\} + \max\{1, |x|^q\} + \max\{1, |y|^q\}] \\ &\leq \frac{\delta}{2} [\max\{1, 2^{q-1}|x|^q\} + 2^{q-1}|y|^q + \max\{1, |x|^q\} + \max\{1, |y|^q\}] \\ &\leq \frac{\delta}{2} [2^q + 2] \max\{1, |x|^q, |y|^q\} = \varepsilon \max\{1, |x|^q, |y|^q\}. \end{aligned} \quad (300)$$

Moreover, observe that (III), (298), the triangle inequality, the fact that for all $\alpha, \beta \in \mathbb{R}$ it holds that $|\alpha + \beta|^2 \leq 2(|\alpha|^2 + |\beta|^2)$, and the fact that $\delta = \varepsilon(2^{q-1} + 1)^{-1}$

prove that for all $x, y \in \mathbb{R}$ it holds that

$$\begin{aligned}
|[\mathcal{R}_a(\Phi)](x, y)| &\leq \frac{1}{2}|[\mathcal{R}_a(\Psi)](x + y)| + \frac{1}{2}|[\mathcal{R}_a(\Psi)](x)| + \frac{1}{2}|[\mathcal{R}_a(\Psi)](y)| \\
&\leq \frac{1}{2}(\delta + |x + y|^2) + \frac{1}{2}(\delta + |x|^2) + \frac{1}{2}(\delta + |y|^2) \\
&\leq \frac{3\delta}{2} + \frac{3}{2}(|x|^2 + |y|^2) = \left[\frac{3\varepsilon}{2}\right][2^{q-1} + 1]^{-1} + \frac{3}{2}(|x|^2 + |y|^2) \\
&= \frac{3}{2}\left[\frac{\varepsilon}{(2^{q-1}+1)} + |x|^2 + |y|^2\right] \leq \frac{3}{2}\left[\frac{\varepsilon}{3} + |x|^2 + |y|^2\right].
\end{aligned} \tag{301}$$

Next note that (I) and (298) prove that for all $x, y \in \mathbb{R}$ it holds that

$$\begin{aligned}
[\mathcal{R}_a(\Phi)](x, 0) &= \frac{1}{2}[\mathcal{R}_a(\Psi)](x) - \frac{1}{2}[\mathcal{R}_a(\Psi)](x) - \frac{1}{2}[\mathcal{R}_a(\Psi)](0) = 0 \\
&= \frac{1}{2}[\mathcal{R}_a(\Psi)](y) - \frac{1}{2}[\mathcal{R}_a(\Psi)](0) - \frac{1}{2}[\mathcal{R}_a(\Psi)](y) = [\mathcal{R}_a(\Phi)](0, y).
\end{aligned} \tag{302}$$

Furthermore, observe that the fact that $\delta = \varepsilon(2^{q-1} + 1)^{-1}$ shows that

$$\begin{aligned}
\left[\frac{q}{2(q-2)}\right] \log_2(\delta^{-1}) + \frac{1}{(q-2)} &= \left[\frac{q}{2(q-2)}\right] \log_2(\varepsilon^{-1}(2^{q-1} + 1)) + \frac{1}{(q-2)} \\
&= \frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1) \right] + \frac{1}{(q-2)} \\
&= \left[\frac{q}{2(q-2)}\right] \log_2(\varepsilon^{-1}) + \left[\frac{q}{2(q-2)}\right] \log_2(2^{q-1} + 1) + \frac{1}{(q-2)}.
\end{aligned} \tag{303}$$

Moreover, observe that Corollary 2.21 implies that $\mathcal{P}(\mathbf{P}_3(\Psi, \Psi, \Psi)) \leq 9\mathcal{P}(\Psi)$. Items (i)–(ii) in Corollary 2.9, (V), (296), and (303) hence ensure that

$$\begin{aligned}
\mathcal{P}(\Phi) &\leq \left[\max\left\{1, \frac{\mathcal{O}(\mathbb{A}_2)}{\mathcal{O}(\mathbf{P}_3(\Psi, \Psi, \Psi))}\right\} \right] \left[\max\left\{1, \frac{\mathcal{I}(\mathbb{A}_1)+1}{\mathcal{I}(\mathbf{P}_3(\Psi, \Psi, \Psi))+1}\right\} \right] \mathcal{P}(\mathbf{P}_3(\Psi, \Psi, \Psi)) \\
&= \left[\max\left\{1, \frac{1}{3}\right\} \right] \left[\max\left\{1, \frac{3}{4}\right\} \right] \mathcal{P}(\mathbf{P}_3(\Psi, \Psi, \Psi)) = \mathcal{P}(\mathbf{P}_3(\Psi, \Psi, \Psi)) \\
&\leq 9\mathcal{P}(\Psi) \leq 9 \max\left\{\left[\frac{40q}{(q-2)}\right] \log_2(\delta^{-1}) + \frac{80}{(q-2)} - 28, 52\right\} \\
&= \max\left\{720\left(\left[\frac{q}{2(q-2)}\right] \log_2(\delta^{-1}) + \frac{1}{(q-2)}\right) - 252, 468\right\} \\
&= \max\left\{720\left(\left[\frac{q}{2(q-2)}\right] \log_2(\varepsilon^{-1}) + \left[\frac{q}{2(q-2)}\right] \log_2(2^{q-1} + 1) + \frac{1}{(q-2)}\right) - 252, 468\right\} \\
&= \max\left\{\frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)) + \frac{720}{(q-2)} - 252, 468\right\}.
\end{aligned} \tag{304}$$

Next note that the fact that for all $r \in (-\infty, 4]$ it holds that $r \geq 2r - 4 = 2(r - 2)$ ensures that for all $r \in (2, 4]$ it holds that $\frac{r(r-1)}{(r-2)} \geq \frac{r}{(r-2)} \geq 2$. This and the fact that for all $r \in [3, \infty)$ it holds that $\frac{r(r-1)}{(r-2)} \geq r - 1 \geq 2$ imply that for all $r \in (2, \infty)$ it holds that $\frac{r(r-1)}{(r-2)} \geq 2$. Hence, we obtain that for all $r \in (2, \infty)$ it holds that

$$\begin{aligned}
\left[\frac{360r}{(r-2)}\right] \log_2(2^{r-1} + 1) - 252 &\geq \left[\frac{360r}{(r-2)}\right] \log_2(2^{r-1}) - 252 \\
&= \frac{360r(r-1)}{(r-2)} - 252 \geq 720 - 252 = 468.
\end{aligned} \tag{305}$$

Combining this with (304) shows that

$$\mathcal{P}(\Phi) \leq \frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)) + \frac{720}{(q-2)} - 252. \quad (306)$$

The fact that

$$\begin{aligned} \log_2(2^{q-1} + 1) &= \log_2(2^{q-1} + 1) - \log_2(2^q) + q = \log_2\left(\frac{2^{q-1}+1}{2^q}\right) + q \\ &= \log_2(2^{-1} + 2^{-q}) + q \leq \log_2(2^{-1} + 2^{-2}) + q \\ &= \log_2\left(\frac{3}{4}\right) + q = \log_2(3) - 2 + q \end{aligned} \quad (307)$$

hence proves that

$$\begin{aligned} \mathcal{P}(\Phi) &\leq \frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)) + \frac{720}{(q-2)} - 252 \\ &\leq \frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + q + \log_2(3) - 2) + \frac{720}{(q-2)} - 252 \\ &= \frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + q + \log_2(3) - 2 + \frac{2}{q}) - 252 \\ &\leq \frac{360q}{(q-2)} (\log_2(\varepsilon^{-1}) + q + \log_2(3) - 1) - 252. \end{aligned} \quad (308)$$

In addition, observe that item (ii) in Proposition 2.6, (296), (VI), the fact that $\delta = \varepsilon(2^{q-1} + 1)^{-1}$, and (303) demonstrate that

$$\begin{aligned} \mathcal{L}(\Phi) &= \mathcal{L}(\mathbf{P}_3(\Psi, \Psi, \Psi)) = \mathcal{L}(\Psi) \\ &\leq \max\left\{\left\lfloor \frac{q}{2(q-2)} \right\rfloor \log_2(\delta^{-1}) + \frac{1}{(q-2)} + 1, 2\right\} \\ &\leq \max\left\{\frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)\right] + \frac{(q-1)}{(q-2)}, 2\right\}. \end{aligned} \quad (309)$$

Furthermore, note that the fact for all $r \in (2, \infty)$ it holds that $\frac{r(r-1)}{(r-2)} \geq 2$ assures that

$$\begin{aligned} &\frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)\right] + \frac{(q-1)}{(q-2)} \\ &\geq \left\lfloor \frac{q}{2(q-2)} \right\rfloor \log_2(2^{q-1}) + 1 = \frac{q(q-1)}{2(q-2)} + 1 \geq 2. \end{aligned} \quad (310)$$

Combining this with (309) proves that

$$\begin{aligned} \mathcal{L}(\Phi) &\leq \frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 1)\right] + \frac{(q-1)}{(q-2)} \\ &\leq \frac{q}{2(q-2)} \left[\log_2(\varepsilon^{-1}) + \log_2(2^{q-1} + 2^{q-1})\right] + \frac{q}{(q-2)} \\ &= \frac{q}{(q-2)} \left[\frac{\log_2(\varepsilon^{-1})}{2} + \frac{q}{2} + 1\right] \leq \frac{q}{(q-2)} \left[\log_2(\varepsilon^{-1}) + \frac{q}{2} + \frac{q}{2}\right] \\ &= \frac{q}{(q-2)} \left[\log_2(\varepsilon^{-1}) + q\right]. \end{aligned} \quad (311)$$

This, the fact that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^2, \mathbb{R})$, (300), (301), (302), and (308) establish items (i)–(vi). The proof of Proposition 3.5 is thus completed. \square

3.2.2 ANN approximations for multi-dimensional products

Definition 3.6 (The Euclidean norm). *We denote by $\|\cdot\| : (\cup_{d \in \mathbb{N}} \mathbb{R}^d) \rightarrow [0, \infty)$ the function which satisfies for all $d \in \mathbb{N}$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ that*

$$\|x\| = \left[\sum_{j=1}^d |x_j|^2 \right]^{1/2}. \quad (312)$$

Proposition 3.7. *Let $\varepsilon \in (0, 1]$, $q \in (2, \infty)$, $d \in \mathbb{N}$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$. Then there exists $\Phi \in \mathbf{N}$ such that*

(i) *it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,*

(ii) *it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Phi))(t, 0) = (\mathcal{R}_a(\Phi))(0, x) = 0$,*

(iii) *it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that*

$$\|tx - (\mathcal{R}_a(\Phi))(t, x)\| \leq \varepsilon(\sqrt{d} [\max\{1, |t|^q\}] + \|x\|^q), \quad (313)$$

(iv) *it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that*

$$\|(\mathcal{R}_a(\Phi))(t, x)\| \leq \sqrt{d}(1 + 2t^2) + 2\|x\|^2, \quad (314)$$

(v) *it holds that $\mathcal{P}(\Phi) \leq d^2 \left[\frac{360q}{(q-2)} \right] [\log_2(\varepsilon^{-1}) + q + 1] - 252d^2$, and*

(vi) *it holds that $\mathcal{L}(\Phi) \leq \frac{q}{(q-2)} [\log_2(\varepsilon^{-1}) + q]$*

(cf. Definition 2.1, Definition 2.3, and Definition 3.6).

Proof of Proposition 3.7. Throughout this proof let $v, w \in \mathbb{R}^{2 \times 1}$, $b \in \mathbb{R}^{2d}$, $A \in \mathbb{R}^{(2d) \times (d+1)}$ satisfy that

$$v = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad w = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad b = 0, \quad (315)$$

and

$$A = \begin{pmatrix} w & v & 0 & 0 & \cdots & 0 \\ w & 0 & v & 0 & \cdots & 0 \\ w & 0 & 0 & v & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ w & 0 & 0 & 0 & \cdots & v \end{pmatrix}, \quad (316)$$

let $\Psi \in \mathbf{N}$ satisfy that

(I) *it holds that $\mathcal{R}_a(\Psi) \in C(\mathbb{R}^2, \mathbb{R})$,*

- (II) it holds for all $x \in \mathbb{R}$ that $[\mathcal{R}_a(\Psi)](x, 0) = [\mathcal{R}_a(\Psi)](0, x) = 0$,
- (III) it holds for all $x, y \in \mathbb{R}$ that $|xy - [\mathcal{R}_a(\Psi)](x, y)| \leq \varepsilon \max\{1, |x|^q, |y|^q\}$,
- (IV) it holds for all $x, y \in \mathbb{R}$ that $|[\mathcal{R}_a(\Psi)](x, y)| \leq 1 + 2x^2 + 2y^2$,
- (V) it holds that $\mathcal{P}(\Psi) \leq \frac{360q}{(q-2)}[\log_2(\varepsilon^{-1}) + q + 1] - 252$, and
- (VI) it holds that $\mathcal{L}(\Psi) \leq \frac{q}{(q-2)}[\log_2(\varepsilon^{-1}) + q]$

(cf. Proposition 3.5), and let $\mathbb{A} \in (\mathbb{R}^{2d \times (d+1)} \times \mathbb{R}^{2d}) \subseteq \mathbf{N}$, $\Phi \in \mathbf{N}$ satisfy that

$$\mathbb{A} = (A, b) \quad \text{and} \quad \Phi = [\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)] \bullet \mathbb{A} \quad (317)$$

(cf. Definition 2.5 and Definition 2.17). Observe that (315) and (316) ensure that for all $y = (y_1, y_2, \dots, y_{d+1}) \in \mathbb{R}^{d+1}$ it holds that

$$Ay = \begin{pmatrix} y_1 w + y_2 v \\ y_1 w + y_3 v \\ \vdots \\ y_1 w + y_{d+1} v \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_1 \\ y_3 \\ \vdots \\ y_1 \\ y_{d+1} \end{pmatrix}. \quad (318)$$

Combining this with (317) proves that for all $t \in \mathbb{R}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_a(\mathbb{A}) \in C(\mathbb{R}^{d+1}, \mathbb{R}^{2d}) \quad \text{and} \quad (\mathcal{R}_a(\mathbb{A}))(t, x) = (t, x_1, t, x_2, \dots, t, x_d). \quad (319)$$

Proposition 2.19, (317), and item (v) in Proposition 2.6 hence demonstrate that for all $t \in \mathbb{R}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$ and

$$\begin{aligned} (\mathcal{R}_a(\Phi))(t, x) &= ([\mathcal{R}_a(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi))] \circ [\mathcal{R}_a(\mathbb{A})])(t, x) \\ &= [\mathcal{R}_a(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi))](t, x_1, t, x_2, \dots, t, x_d) \\ &= ((\mathcal{R}_a(\Psi))(t, x_1), (\mathcal{R}_a(\Psi))(t, x_2), \dots, (\mathcal{R}_a(\Psi))(t, x_d)). \end{aligned} \quad (320)$$

Combining this with (II) proves that for all $t \in \mathbb{R}$ it holds that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(t, 0, 0, \dots, 0) &= ((\mathcal{R}_a(\Psi))(t, 0), (\mathcal{R}_a(\Psi))(t, 0), \dots, (\mathcal{R}_a(\Psi))(t, 0)) \\ &= (0, 0, \dots, 0) = 0. \end{aligned} \quad (321)$$

Next note that (II) and (320) imply that for all $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(0, x) &= ((\mathcal{R}_a(\Psi))(0, x_1), (\mathcal{R}_a(\Psi))(0, x_2), \dots, (\mathcal{R}_a(\Psi))(0, x_d)) \\ &= (0, 0, \dots, 0) = 0. \end{aligned} \quad (322)$$

In addition, observe that the triangle inequality and the fact that for all $r \in [1, \infty)$, $(x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that

$$\left[\sum_{j=1}^d |x_j|^{2r} \right]^{1/2} \leq \left[\sum_{j=1}^d |x_j|^2 \right]^{r/2} \quad (323)$$

prove that for all $b \in \mathbb{R}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$, $r \in [1, \infty)$ it holds that

$$\begin{aligned} \left[\sum_{j=1}^d (|b| + |x_j|^r)^2 \right]^{1/2} &\leq \left[\sum_{j=1}^d b^2 \right]^{1/2} + \left[\sum_{j=1}^d |x_j|^{2r} \right]^{1/2} \\ &\leq |b|\sqrt{d} + \left[\sum_{j=1}^d |x_j|^2 \right]^{r/2} = |b|\sqrt{d} + \|x\|^r. \end{aligned} \quad (324)$$

This, (III), and (320) assure that for all $t \in \mathbb{R}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that

$$\begin{aligned} \|tx - (\mathcal{R}_a(\Phi))(t, x)\| &= \left[\sum_{j=1}^d |tx_j - (\mathcal{R}_a(\Psi))(t, x_j)|^2 \right]^{1/2} \\ &\leq \left[\sum_{j=1}^d [\varepsilon \max\{1, |t|^q, |x_j|^q\}]^2 \right]^{1/2} \leq \varepsilon \left[\sum_{j=1}^d [\max\{1, |t|^q\} + |x_j|^q]^2 \right]^{1/2} \\ &\leq \varepsilon(\sqrt{d} [\max\{1, |t|^q\}] + \|x\|^q). \end{aligned} \quad (325)$$

Furthermore, observe that (IV), (320), and (324) show that for all $t \in \mathbb{R}$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ it holds that

$$\begin{aligned} \|(\mathcal{R}_a(\Phi))(t, x)\| &= \left[\sum_{j=1}^d |(\mathcal{R}_a(\Psi))(t, x_j)|^2 \right]^{1/2} \\ &\leq \left[\sum_{j=1}^d (1 + 2|t|^2 + 2|x_j|^2)^2 \right]^{1/2} \\ &= \left[\sum_{j=1}^d (1 + 2|t|^2 + |\sqrt{2}x_j|^2)^2 \right]^{1/2} \\ &\leq \sqrt{d}(1 + 2|t|^2) + \|\sqrt{2}x\|^2 = \sqrt{d}(1 + 2|t|^2) + 2\|x\|^2. \end{aligned} \quad (326)$$

In addition, note that Corollary 2.21 implies that

$$\mathcal{P}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)) \leq d^2 \mathcal{P}(\Psi). \quad (327)$$

Item (ii) in Corollary 2.9, (V), and (317) hence ensure that

$$\begin{aligned} \mathcal{P}(\Phi) &\leq \left[\max \left\{ 1, \frac{\mathcal{I}(\mathbb{A})+1}{\mathcal{I}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi))+1} \right\} \right] \mathcal{P}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)) \\ &= \left[\max \left\{ 1, \frac{d+2}{2d+1} \right\} \right] \mathcal{P}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)) = \mathcal{P}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)) \quad (328) \\ &\leq d^2 \mathcal{P}(\Psi) \leq d^2 \left[\frac{360q}{(q-2)} \right] [\log_2(\varepsilon^{-1}) + q + 1] - 252d^2. \end{aligned}$$

Next note that item (ii) in Proposition 2.6, (VI), and (317) demonstrate that

$$\mathcal{L}(\Phi) = \mathcal{L}(\mathbf{P}_d(\Psi, \Psi, \dots, \Psi)) = \mathcal{L}(\Psi) \leq \frac{q}{(q-2)} [\log_2(\varepsilon^{-1}) + q]. \quad (329)$$

This, the fact that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$, (321), (322), (325), (326), and (328) establish items (i)–(vi). The proof of Proposition 3.7 is thus completed. \square

3.3 Space-time ANN approximations for Euler approximations

3.3.1 Space-time representations for Euler approximations

Lemma 3.8. *Let $N, d \in \mathbb{N}$, $\mu \in C(\mathbb{R}^d, \mathbb{R}^d)$, $T \in (0, \infty)$, $(t_n)_{n \in \{-1, 0, 1, \dots, N+1\}} \subseteq \mathbb{R}$ satisfy that $t_{-1} < 0 = t_0 < t_1 < \dots < t_N = T < t_{N+1}$, let $f_n: \mathbb{R} \rightarrow \mathbb{R}$, $n \in \{0, 1, \dots, N\}$, be the functions which satisfy for all $n \in \{0, 1, \dots, N\}$, $t \in \mathbb{R}$ that*

$$f_n(t) = \left[\frac{(t-t_{n-1})}{(t_n-t_{n-1})} \right] \mathbb{1}_{(t_{n-1}, t_n)}(t) + \left[\frac{(t_{n+1}-t)}{(t_{n+1}-t_n)} \right] \mathbb{1}_{(t_n, t_{n+1})}(t), \quad (330)$$

and let $Y = (Y_t^{x,y})_{(t,x,y) \in [0,T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N} : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$ be the function which satisfies for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_0^{x,y} = x$ and

$$Y_t^{x,y} = Y_{t_n}^{x,y} + \frac{(t-t_n)}{(t_{n+1}-t_n)} \left[(t_{n+1} - t_n) \mu(Y_{t_n}^{x,y}) + y_{n+1} \right] \quad (331)$$

(cf. Definition 2.1 and Definition 2.3). Then

(i) it holds that

$$\begin{aligned} ([0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \ni (t, x, y) \mapsto Y_t^{x,y} \in \mathbb{R}^d) \\ \in C([0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N, \mathbb{R}^d) \quad (332) \end{aligned}$$

and

(ii) it holds for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$Y_t^{x,y} = \sum_{n=0}^N f_n(t) Y_{t_n}^{x,y}. \quad (333)$$

Proof of Lemma 3.8. Observe that (331) ensures that for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
& Y_{t_n}^{x,y} \left(\frac{t_{n+1}-t}{t_{n+1}-t_n} \right) + Y_{t_{n+1}}^{x,y} \left(\frac{t-t_n}{t_{n+1}-t_n} \right) \\
&= Y_{t_n}^{x,y} \left(1 - \frac{t-t_n}{t_{n+1}-t_n} \right) + Y_{t_{n+1}}^{x,y} \left(\frac{t-t_n}{t_{n+1}-t_n} \right) \\
&= Y_{t_n}^{x,y} \left(1 - \frac{t-t_n}{t_{n+1}-t_n} \right) \\
&\quad + \left(Y_{t_n}^{x,y} + \frac{t_{n+1}-t_n}{t_{n+1}-t_n} [(t_{n+1} - t_n) \mu(Y_{t_n}^{x,y}) + y_{n+1}] \right) \left(\frac{t-t_n}{t_{n+1}-t_n} \right) \\
&= Y_{t_n}^{x,y} + [(t_{n+1} - t_n) \mu(Y_{t_n}^{x,y}) + y_{n+1}] \left(\frac{t-t_n}{t_{n+1}-t_n} \right) = Y_t^{x,y}.
\end{aligned} \tag{334}$$

Hence, we obtain that for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
Y_t^{x,y} &= Y_{t_0}^{x,y} \mathbb{1}_{\{t_0\}}(t) + \sum_{n=0}^{N-1} (Y_t^{x,y} \mathbb{1}_{(t_n, t_{n+1}]}(t)) \\
&= Y_{t_0}^{x,y} \mathbb{1}_{\{t_0\}}(t) + \sum_{n=0}^{N-1} \left(\left[Y_{t_n}^{x,y} \left(\frac{t_{n+1}-t}{t_{n+1}-t_n} \right) + Y_{t_{n+1}}^{x,y} \left(\frac{t-t_n}{t_{n+1}-t_n} \right) \right] \mathbb{1}_{(t_n, t_{n+1}]}(t) \right) \\
&= Y_{t_0}^{x,y} \mathbb{1}_{\{t_0\}}(t) + \left[\sum_{n=0}^{N-1} Y_{t_n}^{x,y} \left(\frac{t-t_{n+1}}{t_n-t_{n+1}} \right) \mathbb{1}_{(t_n, t_{n+1}]}(t) \right] \\
&\quad + \left[\sum_{n=1}^N Y_{t_n}^{x,y} \left(\frac{t-t_{n-1}}{t_n-t_{n-1}} \right) \mathbb{1}_{(t_{n-1}, t_n]}(t) \right].
\end{aligned} \tag{335}$$

Combining this with (330) implies that for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
Y_t^{x,y} &= Y_{t_0}^{x,y} \mathbb{1}_{\{t_0\}}(t) + Y_{t_0}^{x,y} \left(\frac{t-t_1}{t_0-t_1} \right) \mathbb{1}_{(t_0, t_1]}(t) + Y_{t_N}^{x,y} \left(\frac{t-t_{N-1}}{t_N-t_{N-1}} \right) \mathbb{1}_{(t_{N-1}, t_N]}(t) \\
&\quad + \sum_{n=1}^{N-1} Y_{t_n}^{x,y} \left[\left(\frac{t-t_{n+1}}{t_n-t_{n+1}} \right) \mathbb{1}_{(t_n, t_{n+1}]}(t) + \left(\frac{t-t_{n-1}}{t_n-t_{n-1}} \right) \mathbb{1}_{(t_{n-1}, t_n]}(t) \right] \\
&= Y_{t_0}^{x,y} \left(\frac{t_1-t}{t_1-t_0} \right) \mathbb{1}_{[t_0, t_1]}(t) + Y_{t_N}^{x,y} f_N(t) + \sum_{n=1}^{N-1} f_n(t) Y_{t_n}^{x,y} \\
&= \sum_{n=0}^N f_n(t) Y_{t_n}^{x,y}.
\end{aligned} \tag{336}$$

Next we claim that for all $n \in \{0, 1, \dots, N\}$ it holds that

$$(\mathbb{R}^d \times (\mathbb{R}^d)^N \ni (x, y) \mapsto Y_{t_n}^{x,y} \in \mathbb{R}^d) \in C(\mathbb{R}^d \times (\mathbb{R}^d)^N, \mathbb{R}^d). \tag{337}$$

We now prove (337) by induction on $n \in \{0, 1, \dots, N\}$. Note that the fact that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that $Y_{t_0}^{x,y} = Y_0^{x,y} = x$ proves (337) in the base case $n = 0$. For the induction step assume there exists $n \in \{0, 1, \dots, N-1\}$ which satisfies that

$$(\mathbb{R}^d \times (\mathbb{R}^d)^N \ni (x, y) \mapsto Y_{t_n}^{x,y} \in \mathbb{R}^d) \in C(\mathbb{R}^d \times (\mathbb{R}^d)^N, \mathbb{R}^d). \tag{338}$$

Observe that (331) ensures that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$Y_{t_{n+1}}^{x,y} = Y_{t_n}^{x,y} + (t_{n+1} - t_n) \mu(Y_{t_n}^{x,y}) + y_{n+1}. \quad (339)$$

Combining this with (338) and the hypothesis that $\mu \in C(\mathbb{R}^d, \mathbb{R}^d)$ demonstrates that

$$(\mathbb{R}^d \times (\mathbb{R}^d)^N \ni (x, y) \mapsto Y_{t_{n+1}}^{x,y} \in \mathbb{R}^d) \in C(\mathbb{R}^d \times (\mathbb{R}^d)^N, \mathbb{R}^d). \quad (340)$$

Induction thus proves (337). Next observe that (336), (337), and the fact that for all $n \in \{0, 1, \dots, N\}$ it holds that $f_n \in C(\mathbb{R}, \mathbb{R})$ show that

$$\begin{aligned} ([0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \ni (t, x, y) \mapsto Y_t^{x,y} \in \mathbb{R}^d) \\ \in C([0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N, \mathbb{R}^d). \end{aligned} \quad (341)$$

Combining this with (336) establishes items (i)–(ii). The proof of Lemma 3.8 is thus completed. \square

3.3.2 ANN representations for hat functions

Lemma 3.9. *Let $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$, let $\alpha, \beta, \gamma, h \in \mathbb{R}$ satisfy that $\alpha < \beta < \gamma$, let $W_1 \in \mathbb{R}^{4 \times 1}$, $B_1 \in \mathbb{R}^4$, $W_2 \in \mathbb{R}^{1 \times 4}$, $B_2 \in \mathbb{R}$ satisfy that*

$$W_1 = \begin{pmatrix} \frac{1}{(\beta-\alpha)} \\ \frac{1}{(\beta-\alpha)} \\ \frac{1}{(\gamma-\beta)} \\ \frac{1}{(\gamma-\beta)} \end{pmatrix}, \quad B_1 = \begin{pmatrix} -\frac{\alpha}{(\beta-\alpha)} \\ -\frac{\beta}{(\beta-\alpha)} \\ -\frac{\beta}{(\gamma-\beta)} \\ -\frac{\gamma}{(\gamma-\beta)} \end{pmatrix}, \quad (342)$$

$$W_2 = (h \quad -h \quad -h \quad h), \quad B_2 = 0, \quad (343)$$

and let $\Phi \in (\mathbb{R}^{4 \times 1} \times \mathbb{R}^4) \times (\mathbb{R}^{1 \times 4} \times \mathbb{R}) \subseteq \mathbf{N}$ satisfy that $\Phi = ((W_1, B_1), (W_2, B_2))$ (cf. Definition 2.1). Then

(i) it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$ and

(ii) it holds for all $t \in \mathbb{R}$ that

$$\begin{aligned} (\mathcal{R}_a(\Phi))(t) &= \left[\frac{(t-\alpha)h}{(\beta-\alpha)} \right] \mathbb{1}_{(\alpha, \beta]}(t) + \left[\frac{(\gamma-t)h}{(\gamma-\beta)} \right] \mathbb{1}_{(\beta, \gamma)}(t) \\ &= \begin{cases} 0 & : t \in (-\infty, \alpha] \cup [\gamma, \infty) \\ \frac{(t-\alpha)h}{(\beta-\alpha)} & : t \in (\alpha, \beta] \\ \frac{(\gamma-t)h}{(\gamma-\beta)} & : t \in (\beta, \gamma) \end{cases} \end{aligned} \quad (344)$$

(cf. Definition 2.3).

Proof of Lemma 3.9. Observe that for all $t \in \mathbb{R}$ it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$ and

$$\begin{aligned}
(\mathcal{R}_a(\Phi))(t) &= W_2(\mathfrak{M}_{a,4}(W_1 t + B_1)) + B_2 \\
&= h \max\left\{\frac{(t-\alpha)}{(\beta-\alpha)}, 0\right\} - h \max\left\{\frac{(t-\beta)}{(\beta-\alpha)}, 0\right\} - h \max\left\{\frac{(t-\beta)}{(\gamma-\beta)}, 0\right\} + h \max\left\{\frac{(t-\gamma)}{(\gamma-\beta)}, 0\right\} \\
&= h[0 - 0 - 0 + 0] \mathbf{1}_{(-\infty, \alpha]}(t) + h \left[\frac{(t-\alpha)}{(\beta-\alpha)} - 0 - 0 + 0 \right] \mathbf{1}_{(\alpha, \beta]}(t) \\
&\quad + h \left[\frac{(t-\alpha)}{(\beta-\alpha)} - \frac{(t-\beta)}{(\beta-\alpha)} - \frac{(t-\beta)}{(\gamma-\beta)} + 0 \right] \mathbf{1}_{(\beta, \gamma)}(t) \\
&\quad + h \left[\frac{(t-\alpha)}{(\beta-\alpha)} - \frac{(t-\beta)}{(\beta-\alpha)} - \frac{(t-\beta)}{(\gamma-\beta)} + \frac{(t-\gamma)}{(\gamma-\beta)} \right] \mathbf{1}_{[\gamma, \infty)}(t) \\
&= h \left[\frac{(t-\alpha)}{(\beta-\alpha)} \right] \mathbf{1}_{(\alpha, \beta]}(t) + h \left[1 - \frac{(t-\beta)}{(\gamma-\beta)} \right] \mathbf{1}_{(\beta, \gamma)}(t) \\
&= \left[\frac{(t-\alpha)h}{(\beta-\alpha)} \right] \mathbf{1}_{(\alpha, \beta]}(t) + \left[\frac{(\gamma-t)h}{(\gamma-\beta)} \right] \mathbf{1}_{(\beta, \gamma)}(t)
\end{aligned} \tag{345}$$

(cf. Definition 2.2). The proof of Lemma 3.9 is thus completed. \square

3.3.3 A posteriori error estimates for space-time ANN approximations

Proposition 3.10. *Let $N, d \in \mathbb{N}$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$, let $T \in (0, \infty)$, $(t_n)_{n \in \{0, 1, \dots, N\}} \subseteq \mathbb{R}$ satisfy for all $n \in \{0, 1, \dots, N\}$ that $t_n = \frac{nT}{N}$, let $\mathfrak{D} \in [1, \infty)$, $\varepsilon \in (0, 1]$, $q \in (2, \infty)$ satisfy that*

$$\mathfrak{D} = \left\lceil \frac{720q}{(q-2)} \right\rceil \left[\log_2(\varepsilon^{-1}) + q + 1 \right] - 504, \tag{346}$$

let $\Phi \in \mathbf{N}$ satisfy that $\mathcal{I}(\Phi) = \mathcal{O}(\Phi) = d$, and let $Y = (Y_t^{x,y})_{(t,x,y) \in [0,T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N} : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$ be the function which satisfies for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_0^{x,y} = x$ and

$$Y_t^{x,y} = Y_{t_n}^{x,y} + \left(\frac{tN}{T} - n \right) \left[\frac{T}{N} (\mathcal{R}_a(\Phi))(Y_{t_n}^{x,y}) + y_{n+1} \right] \tag{347}$$

(cf. Definition 2.1 and Definition 2.3). Then there exist $\Psi_y \in \mathbf{N}$, $y \in (\mathbb{R}^d)^N$, such that

(i) it holds for all $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_y) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(ii) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|Y_t^{x,y} - (\mathcal{R}_a(\Psi_y))(t, x)\| \leq \varepsilon (2\sqrt{d} + \|Y_{t_n}^{x,y}\|^q + \|Y_{t_{n+1}}^{x,y}\|^q), \tag{348}$$

(iii) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|(\mathcal{R}_a(\Psi_y))(t, x)\| \leq 6\sqrt{d} + 2(\|Y_{t_n}^{x,y}\|^2 + \|Y_{t_{n+1}}^{x,y}\|^2), \quad (349)$$

(iv) it holds for all $y \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} \mathcal{P}(\Psi_y) &\leq \frac{1}{2} \left[6d^2 N^2 \mathcal{H}(\Phi) \right. \\ &\quad \left. + 3N \left[d^2 \mathcal{D} + (23 + 6N \mathcal{H}(\Phi) + 7d^2 + N[4d^2 + \mathcal{P}(\Phi)]^2)^2 \right] \right]^2, \end{aligned} \quad (350)$$

(v) it holds for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_y))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (351)$$

and

(vi) it holds for all $n \in \{0, 1, \dots, N\}$, $t \in [0, t_n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Psi_y))(t, x) = (\mathcal{R}_a(\Psi_z))(t, x) \quad (352)$$

(cf. Definition 3.6).

Proof of Proposition 3.10. Throughout this proof let $t_n \in \mathbb{R}$, $n \in \{-1, N+1\}$, satisfy for all $n \in \{-1, N+1\}$ that $t_n = \frac{nT}{N}$, let $(\mathbb{I}_\mathfrak{d})_{\mathfrak{d} \in \mathbb{N}} \subseteq \mathbf{N}$ satisfy for all $\mathfrak{d} \in \mathbb{N}$, $x \in \mathbb{R}^\mathfrak{d}$ that $\mathcal{R}_a(\mathbb{I}_\mathfrak{d}) \in C(\mathbb{R}^\mathfrak{d}, \mathbb{R}^\mathfrak{d})$, $\mathcal{D}(\mathbb{I}_\mathfrak{d}) = (\mathfrak{d}, 2\mathfrak{d}, \mathfrak{d})$, and

$$(\mathcal{R}_a(\mathbb{I}_\mathfrak{d}))(x) = x \quad (353)$$

(cf., e.g., [25, Lemma 5.4]), let $(\Pi_n)_{n \in \{0, 1, \dots, N\}} \subseteq \mathbf{N}$ satisfy for all $n \in \{0, 1, \dots, N\}$, $t \in \mathbb{R}$ that $\mathcal{I}(\Pi_n) = \mathcal{O}(\Pi_n) = 1$, $\mathcal{H}(\Pi_n) = 1$, $\mathcal{P}(\Pi_n) = 13$, and

$$(\mathcal{R}_a(\Pi_n))(t) = \left[\frac{(t-t_{n-1})}{(t_n-t_{n-1})} \right] \mathbb{1}_{(t_{n-1}, t_n]}(t) + \left[\frac{(t_{n+1}-t)}{(t_{n+1}-t_n)} \right] \mathbb{1}_{(t_n, t_{n+1})}(t) \quad (354)$$

(cf. Lemma 3.9), let $(\Xi_{n,y})_{(n,y) \in \{0, 1, \dots, N\} \times (\mathbb{R}^d)^N} \subseteq \mathbf{N}$ satisfy that

- (I) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Xi_{n,y}) \in C(\mathbb{R}^d, \mathbb{R}^d)$,
- (II) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Xi_{n,y}))(x) = Y_{t_n}^{x,y}$,
- (III) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{H}(\Xi_{n,y}) = 1 + n\mathcal{H}(\Phi)$,

(IV) it holds for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{P}(\Xi_{n,y}) \leq \mathcal{P}(\mathbb{I}_d) + n \left[\frac{1}{2} \mathcal{P}(\mathbb{I}_d) + \mathcal{P}(\Phi) \right]^2, \quad (355)$$

(V) it holds for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Xi_{n,y}))(x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (356)$$

and

(VI) it holds for all $n \in \{0, 1, \dots, N\}$, $m \in \mathbb{N}_0 \cap [0, n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Xi_{m,y}))(x) = (\mathcal{R}_a(\Xi_{m,z}))(x) \quad (357)$$

(cf. Proposition 2.32), let $\Gamma \in \mathbf{N}$ satisfy that

(a) it holds that $\mathcal{R}_a(\Gamma) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(b) it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\Gamma))(t, 0) = (\mathcal{R}_a(\Gamma))(0, x) = 0$,

(c) it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that

$$\|tx - (\mathcal{R}_a(\Gamma))(t, x)\| \leq \varepsilon (\sqrt{d} [\max\{1, |t|^q\}] + \|x\|^q), \quad (358)$$

(d) it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that

$$\|(\mathcal{R}_a(\Gamma))(t, x)\| \leq \sqrt{d}(1 + 2t^2) + 2\|x\|^2, \quad (359)$$

(e) it holds that $\mathcal{P}(\Gamma) \leq d^2 \left[\frac{360q}{(q-2)} \right] [\log_2(\varepsilon^{-1}) + q + 1] - 252d^2$, and

(f) it holds that $\mathcal{L}(\Gamma) \leq \frac{q}{(q-2)} [\log_2(\varepsilon^{-1}) + q]$

(cf. Proposition 3.7), let $(\Psi_{n,y})_{(n,y) \in \{0,1,\dots,N\} \times (\mathbb{R}^d)^N} \subseteq \mathbf{N}$ satisfy for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{I}(\Psi_{n,y}) = d + 1$, $\mathcal{O}(\Psi_{n,y}) = d$, and

$$\Psi_{n,y} = \Gamma \odot_{\mathbb{I}_{d+1}} [\mathbb{P}_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y})] \quad (360)$$

(cf. Definition 2.15, Definition 2.22, Proposition 2.16, and Corollary 2.23), let $L_y \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$, satisfy for all $y \in (\mathbb{R}^d)^N$ that $L_y = \max_{n \in \{0,1,\dots,N\}} \mathcal{L}(\Psi_{n,y})$, and let $(\Phi_y)_{y \in (\mathbb{R}^d)^N} \subseteq \mathbf{N}$ satisfy that

(A) it holds for all $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Phi_y) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(B) it holds for all $y \in (\mathbb{R}^d)^N$, $z \in \mathbb{R}^{d+1}$ that

$$(\mathcal{R}_a(\Phi_y))(z) = \sum_{n=0}^N (\mathcal{R}_a(\Psi_{n,y}))(z), \quad (361)$$

and

(C) it holds for all $y \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} \mathcal{P}(\Phi_y) &\leq \frac{1}{2} \left[\left[\sum_{n=0}^N 2 \mathcal{P}(\Psi_{n,y}) \mathbf{1}_{(\mathcal{L}(\Psi_{n,y}), \infty)}(L_y) \right] \right. \\ &\quad \left. + \left[\sum_{n=0}^N ((L_y - \mathcal{L}(\Psi_{n,y}) - 1) 2d(2d+1) + d(2d+1)) \mathbf{1}_{(\mathcal{L}(\Psi_{n,y}), \infty)}(L_y) \right] \right. \\ &\quad \left. + \left[\sum_{n=0}^N \mathcal{P}(\Psi_{n,y}) \mathbf{1}_{\{\mathcal{L}(\Psi_{n,y})\}}(L_y) \right] \right]^2 \end{aligned} \quad (362)$$

(cf. Proposition 2.26). Note that (III) and the fact that for all $n \in \{0, 1, \dots, N\}$ it holds that $\mathcal{H}(\Pi_n) = 1$ ensure that for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that $\mathcal{L}(\Xi_{n,y}) = 2 + n\mathcal{H}(\Phi) \geq 2$, $\mathcal{L}(\Pi_n) = 2$, and

$$\max\{\mathcal{L}(\Pi_n), \mathcal{L}(\Xi_{n,y})\} = \max\{2, 2 + n\mathcal{H}(\Phi)\} = 2 + n\mathcal{H}(\Phi) = \mathcal{L}(\Xi_{n,y}). \quad (363)$$

Corollary 2.24 (with $a = a$, $n = 2$, $L = \max\{\mathcal{L}(\Pi_n), \mathcal{L}(\Xi_{n,y})\}$, $i_1 = 2$, $i_2 = 2d$, $\Psi = (\mathbb{I}_1, \mathbb{I}_d)$, $\Phi = (\Pi_n, \Xi_{n,y})$ for $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ in the notation of Corollary 2.24), (IV), and the fact that for all $n \in \{0, 1, \dots, N\}$ it holds that $\mathcal{P}(\Pi_n) = 13$ hence prove that for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \mathcal{P}(\mathbb{P}_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y})) &\leq \frac{1}{2} (2\mathcal{P}(\Pi_n) + 6(\mathcal{L}(\Xi_{n,y}) - 3) + 3 + \mathcal{P}(\Xi_{n,y}))^2 \\ &= \frac{1}{2} (11 + 6\mathcal{L}(\Xi_{n,y}) + \mathcal{P}(\Xi_{n,y}))^2 \\ &\leq \frac{1}{2} (11 + 6(2 + n\mathcal{H}(\Phi)) + \mathcal{P}(\mathbb{I}_d) + n[\frac{1}{2}\mathcal{P}(\mathbb{I}_d) + \mathcal{P}(\Phi)]^2)^2 \\ &= \frac{1}{2} (23 + 6n\mathcal{H}(\Phi) + \mathcal{P}(\mathbb{I}_d) + n[\frac{1}{2}\mathcal{P}(\mathbb{I}_d) + \mathcal{P}(\Phi)]^2)^2. \end{aligned} \quad (364)$$

Moreover, observe that (364) and (e) imply that $2\mathcal{P}(\Gamma) \leq d^2\mathfrak{D}$. Combining this with Proposition 2.16, (364), and the fact that $\mathcal{P}(\mathbb{I}_d) = 4d^2 + 3d \leq 4(d^2 + d)$ ensures that for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \mathcal{P}(\Psi_{n,y}) &= \mathcal{P}(\Gamma \odot_{\mathbb{I}_{d+1}} [\mathbb{P}_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y})]) \\ &\leq \max\left\{1, \frac{2(d+1)}{(d+1)}\right\} (\mathcal{P}(\Gamma) + \mathcal{P}(\mathbb{P}_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y}))) \\ &\leq d^2\mathfrak{D} + (23 + 6n\mathcal{H}(\Phi) + \mathcal{P}(\mathbb{I}_d) + n[\frac{1}{2}\mathcal{P}(\mathbb{I}_d) + \mathcal{P}(\Phi)]^2)^2 \\ &\leq d^2\mathfrak{D} + (23 + 6n\mathcal{H}(\Phi) + 4d^2 + 3d + n[2(d^2 + d) + \mathcal{P}(\Phi)]^2)^2. \end{aligned} \quad (365)$$

Next note that (III), (363), (69), (118), (101), item (ii) in Proposition 2.16, and item (i) in Lemma 2.13 demonstrate that for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
\mathcal{L}(\Psi_{n,y}) &= \mathcal{L}(\Gamma) + \mathcal{L}(\mathbf{P}_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y})) \\
&= \mathcal{L}(\Gamma) + \mathcal{L}(\mathbf{P}_2(\mathcal{E}_{\max\{\mathcal{L}(\Pi_n), \mathcal{L}(\Xi_{n,y})\}, \mathbb{I}_1}(\Pi_n), \mathcal{E}_{\max\{\mathcal{L}(\Pi_n), \mathcal{L}(\Xi_{n,y})\}, \mathbb{I}_d}(\Xi_{n,y}))) \\
&= \mathcal{L}(\Gamma) + \mathcal{L}(\mathbf{P}_2(\mathcal{E}_{\mathcal{L}(\Xi_{n,y}), \mathbb{I}_1}(\Pi_n), \mathcal{E}_{\mathcal{L}(\Xi_{n,y}), \mathbb{I}_d}(\Xi_{n,y}))) \\
&= \mathcal{L}(\Gamma) + \mathcal{L}(\mathcal{E}_{\mathcal{L}(\Xi_{n,y}), \mathbb{I}_d}(\Xi_{n,y})) \\
&= \mathcal{L}(\Gamma) + \mathcal{L}((\mathbb{I}_d)^{\bullet 0} \bullet \Xi_{n,y}) \\
&= \mathcal{L}(\Gamma) + \mathcal{L}((\mathbb{I}_d)^{\bullet 0}) + \mathcal{L}(\Xi_{n,y}) - 1 \\
&= \mathcal{L}(\Gamma) + \mathcal{L}(\Xi_{n,y}) = \mathcal{L}(\Gamma) + \mathcal{H}(\Xi_{n,y}) + 1 \\
&= \mathcal{L}(\Gamma) + 2 + n\mathcal{H}(\Phi).
\end{aligned} \tag{366}$$

Therefore, we obtain that for all $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
\mathcal{L}(\Psi_{N,y}) - \mathcal{L}(\Psi_{n,y}) - 1 &= (\mathcal{L}(\Gamma) + 2 + N\mathcal{H}(\Phi)) - (\mathcal{L}(\Gamma) + 2 + n\mathcal{H}(\Phi)) - 1 \\
&= (N - n)\mathcal{H}(\Phi) - 1.
\end{aligned} \tag{367}$$

In addition, note that (366) proves that for all $y \in (\mathbb{R}^d)^N$ it holds that $L_y = \mathcal{L}(\Psi_{N,y}) = \mathcal{L}(\Psi_{N,0}) = L_0$. The fact that $\sum_{n=0}^{N-1} (N - n) = \sum_{m=1}^N m = \frac{1}{2}N(N + 1)$, (362), and (367) hence assure that for all $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
&\mathcal{P}(\Phi_y) \\
&\leq \frac{1}{2} \left[\sum_{n=0}^{N-1} (2\mathcal{P}(\Psi_{n,y}) \right. \\
&\quad \left. + \max\{(\mathcal{L}(\Psi_{N,y}) - \mathcal{L}(\Psi_{n,y}) - 1)2d(2d + 1) + d(2d + 1), 0\}) \right] + \mathcal{P}(\Psi_{N,y}) \Big]^2 \\
&= \frac{1}{2} \left[\sum_{n=0}^{N-1} (2\mathcal{P}(\Psi_{n,y}) + \max\{(N - n)\mathcal{H}(\Phi)2d(2d + 1) - d(2d + 1), 0\}) \right] \\
&\quad \left. + \mathcal{P}(\Psi_{N,y}) \right]^2 \\
&\leq \frac{1}{2} \left[(2N + 1)\mathcal{P}(\Psi_{N,y}) \right. \\
&\quad \left. + \max\{\mathcal{H}(\Phi)2d(2d + 1) \left[\sum_{n=0}^{N-1} (N - n) \right] - Nd(2d + 1), 0\} \right]^2 \\
&= \frac{1}{2} \left[(2N + 1)\mathcal{P}(\Psi_{N,y}) + \max\{\mathcal{H}(\Phi)d(2d + 1)N(N + 1) - Nd(2d + 1), 0\} \right]^2.
\end{aligned} \tag{368}$$

This and (365) imply that for all $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \mathcal{P}(\Phi_y) &\leq \frac{1}{2} \left[(2N+1) \left[d^2 \mathfrak{D} + (23 + 6N\mathcal{H}(\Phi) + 4d^2 + 3d + N[2(d^2 + d) + \mathcal{P}(\Phi)]^2)^2 \right] \right. \\ &\quad \left. + \max\{\mathcal{H}(\Phi)d(2d+1)N(N+1) - Nd(2d+1), 0\} \right]^2. \end{aligned} \quad (369)$$

Therefore, we obtain that for all $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} &\mathcal{P}(\Phi_y) \\ &\leq \frac{1}{2} \left[(2N+1) \left[d^2 \mathfrak{D} + (23 + 6N\mathcal{H}(\Phi) + 7d^2 + N[4d^2 + \mathcal{P}(\Phi)]^2)^2 \right] \right. \\ &\quad \left. + \max\{\mathcal{H}(\Phi)d(2d+1)N(N+1) - Nd(2d+1), 0\} \right]^2 \\ &\leq \frac{1}{2} \left[3N \left[d^2 \mathfrak{D} + (23 + 6N\mathcal{H}(\Phi) + 7d^2 + N[4d^2 + \mathcal{P}(\Phi)]^2)^2 \right] + 6d^2 N^2 \mathcal{H}(\Phi) \right]^2. \end{aligned} \quad (370)$$

In addition, note that (354), (II), and Lemma 3.8 (with $N = N$, $d = d$, $\mu = \mathcal{R}_a(\Phi)$, $T = T$, $(\{-1, 0, 1, \dots, N+1\} \ni n \mapsto t_n \in \mathbb{R}) = (\{-1, 0, 1, \dots, N+1\} \ni n \mapsto t_n \in \mathbb{R})$, $(\{0, 1, \dots, N\} \ni n \mapsto f_n \in C(\mathbb{R}, \mathbb{R})) = (\{0, 1, \dots, N\} \ni n \mapsto \mathcal{R}_a(\Pi_n) \in C(\mathbb{R}, \mathbb{R}))$, $Y = Y$ in the notation of Lemma 3.8) ensure that for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$Y_t^{x,y} = \sum_{n=0}^N [(\mathcal{R}_a(\Pi_n))(t)] Y_{t_n}^{x,y} = \sum_{n=0}^N [(\mathcal{R}_a(\Pi_n))(t)] [(\mathcal{R}_a(\Xi_{n,y}))(x)]. \quad (371)$$

Moreover, observe that (360), (361), item (iv) in Proposition 2.16 (with $\Psi = \mathbb{I}_{d+1}$, $\Phi_1 = \Gamma$, $\Phi_2 = P_{2,(\mathbb{I}_1, \mathbb{I}_d)}(\Pi_n, \Xi_{n,y})$, $\mathbf{i} = 2(d+1)$ for $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ in the notation of Proposition 2.16), and Corollary 2.23 (with $a = a$, $n = 2$, $\mathbb{I} = (\mathbb{I}_1, \mathbb{I}_d)$, $\Phi = (\Pi_n, \Xi_{n,y})$ for $n \in \{0, 1, \dots, N\}$, $y \in (\mathbb{R}^d)^N$ in the notation of Corollary 2.23) demonstrate that for all $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$(\mathcal{R}_a(\Phi_y))(t, x) = \sum_{n=0}^N (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_n))(t), (\mathcal{R}_a(\Xi_{n,y}))(x)). \quad (372)$$

Next note that (354) shows that for all $k \in \{0, 1, \dots, N\}$, $t \in \mathbb{R} \setminus (t_{k-1}, t_{k+1})$ it holds that

$$(\mathcal{R}_a(\Pi_k))(t) = 0. \quad (373)$$

Combining this, (371), and (372) with (b) proves that for all $k \in \{0, 1, \dots, N-1\}$, $t \in [t_k, t_{k+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} Y_t^{x,y} &= \sum_{n=0}^N [(\mathcal{R}_a(\Pi_n))(t)] [(\mathcal{R}_a(\Xi_{n,y}))(x)] \\ &= [(\mathcal{R}_a(\Pi_k))(t)] [(\mathcal{R}_a(\Xi_{k,y}))(x)] + [(\mathcal{R}_a(\Pi_{k+1}))(t)] [(\mathcal{R}_a(\Xi_{k+1,y}))(x)] \end{aligned} \quad (374)$$

and

$$\begin{aligned} (\mathcal{R}_a(\Phi_y))(t, x) &= \sum_{n=0}^N (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_n))(t), (\mathcal{R}_a(\Xi_{n,y}))(x)) \\ &= (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_k))(t), (\mathcal{R}_a(\Xi_{k,y}))(x)) \\ &\quad + (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_{k+1}))(t), (\mathcal{R}_a(\Xi_{k+1,y}))(x)). \end{aligned} \quad (375)$$

The triangle inequality, (c), and (d) hence establish that for all $k \in \{0, 1, \dots, N-1\}$, $t \in [t_k, t_{k+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} &\|Y_t^{x,y} - (\mathcal{R}_a(\Phi_y))(t, x)\| \\ &\leq \sum_{n=k}^{k+1} \left\| [(\mathcal{R}_a(\Pi_n))(t)] [(\mathcal{R}_a(\Xi_{n,y}))(x)] - (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_n))(t), (\mathcal{R}_a(\Xi_{n,y}))(x)) \right\| \\ &\leq \sum_{n=k}^{k+1} \varepsilon (\sqrt{d} [\max\{1, |(\mathcal{R}_a(\Pi_n))(t)|^q\}] + \|(\mathcal{R}_a(\Xi_{n,y}))(x)\|^q) \end{aligned} \quad (376)$$

and

$$\begin{aligned} \|(\mathcal{R}_a(\Phi_y))(t, x)\| &\leq \sum_{n=k}^{k+1} \|(\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_n))(t), (\mathcal{R}_a(\Xi_{n,y}))(x))\| \\ &\leq \sum_{n=k}^{k+1} (\sqrt{d}(1 + 2|(\mathcal{R}_a(\Pi_n))(t)|^2) + 2\|(\mathcal{R}_a(\Xi_{n,y}))(x)\|^2). \end{aligned} \quad (377)$$

Next note that (354) ensures that for all $n \in \{0, 1, \dots, N\}$, $t \in \mathbb{R}$ it holds that $0 \leq (\mathcal{R}_a(\Pi_n))(t) \leq 1$. Combining this with (376), (377), and (II) demonstrates that for all $k \in \{0, 1, \dots, N-1\}$, $t \in [t_k, t_{k+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \|Y_t^{x,y} - (\mathcal{R}_a(\Phi_y))(t, x)\| &\leq \sum_{n=k}^{k+1} \varepsilon (\sqrt{d} + \|(\mathcal{R}_a(\Xi_{n,y}))(x)\|^q) \\ &= \varepsilon (\sqrt{d} + \|Y_{t_k}^{x,y}\|^q) + \varepsilon (\sqrt{d} + \|Y_{t_{k+1}}^{x,y}\|^q) \\ &= \varepsilon (2\sqrt{d} + \|Y_{t_k}^{x,y}\|^q + \|Y_{t_{k+1}}^{x,y}\|^q) \end{aligned} \quad (378)$$

and

$$\begin{aligned}
\|(\mathcal{R}_a(\Phi_y))(t, x)\| &\leq \sum_{n=k}^{k+1} (3\sqrt{d} + 2\|(\mathcal{R}_a(\Xi_{n,y}))(x)\|^2) \\
&= 3\sqrt{d} + 2\|Y_{t_k}^{x,y}\|^2 + 3\sqrt{d} + 2\|Y_{t_{k+1}}^{x,y}\|^2 \\
&= 6\sqrt{d} + 2(\|Y_{t_k}^{x,y}\|^2 + \|Y_{t_{k+1}}^{x,y}\|^2).
\end{aligned} \tag{379}$$

Furthermore, observe that (372), (V), and (a) ensure that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Phi_y))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d). \tag{380}$$

In addition, observe that (b), (372) and (373) demonstrate that for all $n \in \{0, 1, \dots, N\}$, $t \in [0, t_n]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$(\mathcal{R}_a(\Phi_y))(t, x) = \sum_{k=0}^n (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_k))(t), (\mathcal{R}_a(\Xi_{k,y}))(x)). \tag{381}$$

This and (VI) show that for all $n \in \{0, 1, \dots, N\}$, $t \in [0, t_n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ it holds that

$$\begin{aligned}
(\mathcal{R}_a(\Phi_y))(t, x) &= \sum_{m=0}^n (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_m))(t), (\mathcal{R}_a(\Xi_{m,y}))(x)) \\
&= \sum_{m=0}^n (\mathcal{R}_a(\Gamma))((\mathcal{R}_a(\Pi_m))(t), (\mathcal{R}_a(\Xi_{m,z}))(x)) \\
&= (\mathcal{R}_a(\Phi_z))(t, x).
\end{aligned} \tag{382}$$

Combining this with (A), (370), (378), (379), and (380) establishes items (i)–(vi). The proof of Proposition 3.10 is thus completed. \square

3.3.4 A priori estimates for Euler approximations

Lemma 3.11. *Let $N, d \in \mathbb{N}$, $c, C \in [0, \infty)$, $A_1, A_2, \dots, A_N \in \mathbb{R}^{d \times d}$, let $\|\cdot\|: \mathbb{R}^d \rightarrow [0, \infty)$ be a norm on \mathbb{R}^d , let $\|\|\cdot\|\|: \mathbb{R}^{d \times d} \rightarrow [0, \infty)$ be the function which satisfies for all $A \in \mathbb{R}^{d \times d}$ that $\|\|A\|\| = \sup_{\{x \in \mathbb{R}^d: \|x\| \leq 1\}} \|Ax\|$, let $\mu: \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a function which satisfies for all $x \in \mathbb{R}^d$ that*

$$\|\mu(x)\| \leq C + c\|x\|, \tag{383}$$

and let $Y_n = (Y_n^{x,y})_{(x,y) \in \mathbb{R}^d \times (\mathbb{R}^d)^N}: \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$, $n \in \{0, 1, \dots, N\}$, be the functions which satisfy for all $n \in \{0, 1, \dots, N-1\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_0^{x,y} = x$ and

$$Y_{n+1}^{x,y} = Y_n^{x,y} + A_{n+1} \mu(Y_n^{x,y}) + y_{n+1}. \tag{384}$$

Then

(i) it holds for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that

$$Y_n^{x,y} = x + \sum_{k=0}^{n-1} [A_{k+1} \mu(Y_k^{x,y}) + y_{k+1}] \quad (385)$$

and

(ii) it holds for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} & \|Y_n^{x,y}\| \\ & \leq \left(\|x\| + C \left[\sum_{k=1}^n \|A_k\| \right] + \max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right) \exp \left(c \left[\sum_{k=1}^n \|A_k\| \right] \right). \end{aligned} \quad (386)$$

Proof of Lemma 3.11. We claim that for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$Y_n^{x,y} = x + \sum_{k=0}^{n-1} [A_{k+1} \mu(Y_k^{x,y}) + y_{k+1}]. \quad (387)$$

We now prove (387) by induction on $n \in \{0, 1, \dots, N\}$. Observe that the hypothesis that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that $Y_0^{x,y} = x$ proves (387) in the base case $n = 0$. For the induction step note that (384) implies that for all $n \in \{0, 1, \dots, N-1\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ with

$$Y_n^{x,y} = x + \sum_{k=0}^{n-1} [A_{k+1} \mu(Y_k^{x,y}) + y_{k+1}] \quad (388)$$

it holds that

$$\begin{aligned} Y_{n+1}^{x,y} &= Y_n^{x,y} + A_{n+1} \mu(Y_n^{x,y}) + y_{n+1} \\ &= x + \left[\sum_{k=0}^{n-1} (A_{k+1} \mu(Y_k^{x,y}) + y_{k+1}) \right] + (A_{n+1} \mu(Y_n^{x,y}) + y_{n+1}) \\ &= x + \left[\sum_{k=0}^n (A_{k+1} \mu(Y_k^{x,y}) + y_{k+1}) \right]. \end{aligned} \quad (389)$$

Induction thus proves (387). Observe that (387) establishes item (i). In addition, note that (387), the triangle inequality, and the fact that for all $A \in \mathbb{R}^{d \times d}$, $x \in \mathbb{R}^d$ it holds that $\|Ax\| \leq \|A\| \|x\|$ demonstrate that for all $m \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\|Y_m^{x,y}\| \leq \|x\| + \left[\sum_{k=0}^{m-1} \|A_{k+1}\| \|\mu(Y_k^{x,y})\| \right] + \left\| \sum_{k=0}^{m-1} y_{k+1} \right\|. \quad (390)$$

Combining this with (383) ensures that for all $n \in \{0, 1, \dots, N\}$, $m \in \{0, 1, \dots, n\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned}
\|Y_m^{x,y}\| &\leq \|x\| + \left[\sum_{k=0}^{m-1} \|A_{k+1}\| (C + c\|Y_k^{x,y}\|) \right] + \left\| \sum_{k=1}^m y_k \right\| \\
&= \|x\| + C \left[\sum_{k=1}^m \|A_k\| \right] + \left\| \sum_{k=1}^m y_k \right\| + c \left[\sum_{k=0}^{m-1} \|A_{k+1}\| \|Y_k^{x,y}\| \right] \\
&\leq \|x\| + C \left[\sum_{k=1}^n \|A_k\| \right] + \left[\max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right] + c \left[\sum_{k=0}^{m-1} \|A_{k+1}\| \|Y_k^{x,y}\| \right].
\end{aligned} \tag{391}$$

The time-discrete Gronwall inequality (cf., e.g., Hutzenthaler et al. [23, Lemma 2.1] (with $N = n$, $\alpha = (\|x\| + C[\sum_{k=1}^n \|A_k\|] + \max_{m \in \{0,1,\dots,n\}} \|\sum_{k=1}^m y_k\|)$, $\beta_0 = c\|A_1\|$, $\beta_1 = c\|A_2\|$, \dots , $\beta_{n-1} = c\|A_n\|$, $\epsilon_0 = \|Y_0^{x,y}\|$, $\epsilon_1 = \|Y_1^{x,y}\|$, \dots , $\epsilon_n = \|Y_n^{x,y}\|$ for $n \in \{1, 2, \dots, N\}$ in the notation of Hutzenthaler et al. [23, Lemma 2.1])) hence implies that for all $n \in \{1, 2, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\|Y_n^{x,y}\| \leq \left(\|x\| + C \left[\sum_{k=1}^n \|A_k\| \right] + \max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right) \exp \left(c \left[\sum_{k=0}^{n-1} \|A_{k+1}\| \right] \right). \tag{392}$$

The hypothesis that for all $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that $Y_0^{x,y} = x$ therefore assures that for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\|Y_n^{x,y}\| \leq \left(\|x\| + C \left[\sum_{k=1}^n \|A_k\| \right] + \max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right) \exp \left(c \left[\sum_{k=1}^n \|A_k\| \right] \right). \tag{393}$$

This establishes item (ii). The proof of Lemma 3.11 is thus completed. \square

3.3.5 A priori error estimates for space-time ANN approximations

Theorem 3.12. *Let $N, d \in \mathbb{N}$, $\mathfrak{C} \in [0, \infty)$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$, let $T \in (0, \infty)$, $(t_n)_{n \in \{0,1,\dots,N\}} \subseteq \mathbb{R}$ satisfy for all $n \in \{0, 1, \dots, N\}$ that $t_n = \frac{nT}{N}$, let $\mathfrak{D} \in [1, \infty)$, $\varepsilon \in (0, 1]$, $q \in (2, \infty)$ satisfy that*

$$\mathfrak{D} = \left\lceil \frac{720q}{(q-2)} \right\rceil \left[\log_2(\varepsilon^{-1}) + q + 1 \right] - 504, \tag{394}$$

let $\Phi \in \mathbf{N}$ satisfy for all $x \in \mathbb{R}^d$ that $\mathcal{I}(\Phi) = \mathcal{O}(\Phi) = d$ and $\|(\mathcal{R}_a(\Phi))(x)\| \leq \mathfrak{C}(1 + \|x\|)$, let $Y = (Y_t^{x,y})_{(t,x,y) \in [0,T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N} : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$ be the function which satisfies for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_0^{x,y} = x$ and

$$Y_t^{x,y} = Y_{t_n}^{x,y} + \left(\frac{tN}{T} - n \right) \left[\frac{T}{N} (\mathcal{R}_a(\Phi))(Y_{t_n}^{x,y}) + y_{n+1} \right], \tag{395}$$

and let $g_n: \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow [0, \infty)$, $n \in \{0, 1, \dots, N\}$, be the functions which satisfy for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that

$$g_n(x, y) = \left(\|x\| + \mathfrak{C}t_n + \max_{m \in \{0, 1, \dots, n\}} \left\| \sum_{k=1}^m y_k \right\| \right) \exp(\mathfrak{C}t_n) \quad (396)$$

(cf. Definition 2.1, Definition 2.3, and Definition 3.6). Then there exist $\Psi_y \in \mathbf{N}$, $y \in (\mathbb{R}^d)^N$, such that

(i) it holds for all $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_y) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(ii) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|Y_t^{x,y} - (\mathcal{R}_a(\Psi_y))(t, x)\| \leq \varepsilon(2\sqrt{d} + (g_n(x, y))^q + (g_{n+1}(x, y))^q), \quad (397)$$

(iii) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|(\mathcal{R}_a(\Psi_y))(t, x)\| \leq 6\sqrt{d} + 2((g_n(x, y))^2 + (g_{n+1}(x, y))^2), \quad (398)$$

(iv) it holds for all $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{P}(\Psi_y) \leq \frac{9}{2} N^6 d^{16} \left[2\mathcal{H}(\Phi) + \mathfrak{D} + (30 + 6\mathcal{H}(\Phi) + [4 + \mathcal{P}(\Phi)]^2)^2 \right]^2, \quad (399)$$

(v) it holds for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_y))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (400)$$

and

(vi) it holds for all $n \in \{0, 1, \dots, N\}$, $t \in [0, t_n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Psi_y))(t, x) = (\mathcal{R}_a(\Psi_z))(t, x). \quad (401)$$

Proof of Theorem 3.12. Throughout this proof let $\Psi_y \in \mathbf{N}$, $y \in (\mathbb{R}^d)^N$, satisfy that

(I) it holds for all $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_y) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(II) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|Y_t^{x,y} - (\mathcal{R}_a(\Psi_y))(t, x)\| \leq \varepsilon(2\sqrt{d} + \|Y_{t_n}^{x,y}\|^q + \|Y_{t_{n+1}}^{x,y}\|^q), \quad (402)$$

(III) it holds for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|(\mathcal{R}_a(\Psi_y))(t, x)\| \leq 6\sqrt{d} + 2(\|Y_{t_n}^{x,y}\|^2 + \|Y_{t_{n+1}}^{x,y}\|^2), \quad (403)$$

(IV) it holds for all $y \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} \mathcal{P}(\Psi_y) &\leq \frac{1}{2} \left[6d^2 N^2 \mathcal{H}(\Phi) \right. \\ &\quad \left. + 3N \left[d^2 \mathfrak{D} + (23 + 6N\mathcal{H}(\Phi) + 7d^2 + N[4d^2 + \mathcal{P}(\Phi)]^2)^2 \right] \right]^2, \end{aligned} \quad (404)$$

(V) it holds for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_y))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (405)$$

and

(VI) it holds for all $n \in \{0, 1, \dots, N\}$, $t \in [0, t_n]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Psi_y))(t, x) = (\mathcal{R}_a(\Psi_z))(t, x) \quad (406)$$

(cf. Proposition 3.10). Note that (IV) ensures for all $y \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} &\mathcal{P}(\Psi_y) \\ &\leq \frac{1}{2} \left[6d^2 N^2 \mathcal{H}(\Phi) + 3N \left[d^2 \mathfrak{D} + (23 + 6N\mathcal{H}(\Phi) + 7d^2 + Nd^4 [4 + \mathcal{P}(\Phi)]^2)^2 \right] \right]^2 \\ &\leq \frac{1}{2} \left[6d^2 N^2 \mathcal{H}(\Phi) + 3N \left[d^2 \mathfrak{D} + N^2 d^8 (30 + 6\mathcal{H}(\Phi) + [4 + \mathcal{P}(\Phi)]^2)^2 \right] \right]^2. \end{aligned} \quad (407)$$

Hence, we obtain that for all $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \mathcal{P}(\Psi_y) &\leq \frac{1}{2} \left[6d^2 N^2 \mathcal{H}(\Phi) + 3N^3 d^8 \left[\mathfrak{D} + (30 + 6\mathcal{H}(\Phi) + [4 + \mathcal{P}(\Phi)]^2)^2 \right] \right]^2 \\ &\leq \frac{9}{2} N^6 d^{16} \left[2\mathcal{H}(\Phi) + \mathfrak{D} + (30 + 6\mathcal{H}(\Phi) + [4 + \mathcal{P}(\Phi)]^2)^2 \right]^2. \end{aligned} \quad (408)$$

In addition, observe that Lemma 3.11 and the hypothesis that for all $n \in \{0, 1, \dots, N\}$ it holds that $t_n = \frac{nT}{N}$ demonstrate that for all $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \|Y_{t_n}^{x,y}\| &\leq \left[\|x\| + \frac{\mathfrak{C}nT}{N} + \max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right] \exp\left(\frac{\mathfrak{C}nT}{N}\right) \\ &= \left[\|x\| + \mathfrak{C}t_n + \max_{m \in \{0,1,\dots,n\}} \left\| \sum_{k=1}^m y_k \right\| \right] \exp(\mathfrak{C}t_n) = g_n(x, y). \end{aligned} \quad (409)$$

Combining this with (II) and (III) ensures that for all $n \in \{0, 1, \dots, N-1\}$, $t \in [t_n, t_{n+1}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \|Y_t^{x,y} - (\mathcal{R}_a(\Psi_y))(t, x)\| &\leq \varepsilon(2\sqrt{d} + \|Y_{t_n}^{x,y}\|^q + \|Y_{t_{n+1}}^{x,y}\|^q) \\ &\leq \varepsilon(2\sqrt{d} + (g_n(x, y))^q + (g_{n+1}(x, y))^q) \end{aligned} \quad (410)$$

and

$$\begin{aligned} \|(\mathcal{R}_a(\Psi_y))(t, x)\| &\leq 6\sqrt{d} + 2(\|Y_{t_n}^{x,y}\|^2 + \|Y_{t_{n+1}}^{x,y}\|^2) \\ &\leq 6\sqrt{d} + 2((g_n(x, y))^2 + (g_{n+1}(x, y))^2). \end{aligned} \quad (411)$$

This, (I), (V), (VI), and (408) establish items (i)–(vi). The proof of Theorem 3.12 is thus completed. \square

Corollary 3.13. *Let $\mathfrak{C}, T, \mathfrak{d} \in (0, \infty)$, $a \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, 0\}$, let $\Phi_d \in \mathbf{N}$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$ that $\mathcal{I}(\Phi_d) = \mathcal{O}(\Phi_d) = d$, $\|(\mathcal{R}_a(\Phi_d))(x)\| \leq \mathfrak{C}(1 + \|x\|)$, and $\mathcal{P}(\Phi_d) \leq \mathfrak{C}d^{\mathfrak{p}}$, let $Y^{d,N} = (Y_{t,x,y}^{d,N})_{(t,x,y) \in [0,T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N} : [0, T] \times \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow \mathbb{R}^d$, $N, d \in \mathbb{N}$, be the functions which satisfy for all $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that $Y_{0,x,y}^{d,N} = x$ and*

$$Y_{t,x,y}^{d,N} = Y_{\frac{nT}{N},x,y}^{d,N} + \left(\frac{tN}{T} - n\right) \left[\frac{T}{N}(\mathcal{R}_a(\Phi_d))(Y_{\frac{nT}{N},x,y}^{d,N}) + y_{n+1}\right]. \quad (412)$$

(cf. Definition 2.1, Definition 2.3, and Definition 3.6). Then there exist $C \in \mathbb{R}$ and $\Psi_{\varepsilon,d,N,y} \in \mathbf{N}$, $y \in (\mathbb{R}^d)^N$, $N, d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, such that

(i) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_{\varepsilon,d,N,y}) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(ii) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|Y_{t,x,y}^{d,N} - (\mathcal{R}_a(\Psi_{\varepsilon,d,N,y}))(t, x)\| \leq Cd^{1/2}N^{3/2}\varepsilon(1 + \|x\|^3 + \|y\|^3), \quad (413)$$

(iii) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|(\mathcal{R}_a(\Psi_{\varepsilon,d,N,y}))(t, x)\| \leq Cd^{1/2}N(1 + \|x\|^2 + \|y\|^2), \quad (414)$$

(iv) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$ that

$$\mathcal{P}(\Psi_{\varepsilon,d,N,y}) \leq Cd^{16+8\mathfrak{d}}N^6[1 + |\ln(\varepsilon)|^2], \quad (415)$$

(v) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_{\varepsilon,d,N,y}))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (416)$$

and

(vi) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N\}$, $t \in [0, \frac{nT}{N}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ that

$$(\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x) = (\mathcal{R}_a(\Psi_{\varepsilon, d, N, z}))(t, x). \quad (417)$$

Proof of Corollary 3.13. Throughout this proof let $\mathfrak{D}_{\varepsilon, q} \in [1, \infty)$, $q \in (2, \infty)$, $\varepsilon \in (0, 1]$, satisfy for all $\varepsilon \in (0, 1]$, $q \in (2, \infty)$ that

$$\mathfrak{D}_{\varepsilon, q} = \left\lceil \frac{720q}{(q-2)} \right\rceil \left[\log_2(\varepsilon^{-1}) + q + 1 \right] - 504, \quad (418)$$

let $c = \max\{\exp(\mathfrak{C}T), \mathfrak{D}_{1,3}, 62 + 6\mathfrak{C}(\mathfrak{C} + 1)\}$, and let $g_n^{d,N}: \mathbb{R}^d \times (\mathbb{R}^d)^N \rightarrow [0, \infty)$, $n \in \{0, 1, \dots, N\}$, $N, d \in \mathbb{N}$, be the functions which satisfy for all $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N\}$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ that

$$g_n^{d,N}(x, y) = \left(\|x\| + \frac{\mathfrak{C}nT}{N} + \max_{m \in \{0, 1, \dots, n\}} \left\| \sum_{k=1}^m y_k \right\| \right) \exp\left(\frac{\mathfrak{C}nT}{N}\right). \quad (419)$$

Note that Theorem 3.12 (with $N = N$, $d = d$, $\mathfrak{C} = \mathfrak{C}$, $a = a$, $T = T$, $t_n = \frac{nT}{N}$, $\mathfrak{D} = \mathfrak{D}_{\varepsilon, 3}$, $\varepsilon = \varepsilon$, $q = 3$, $\Phi = \Phi_d$, $Y = Y^{d,N}$, $g_n = g_n^{d,N}$ for $N, d \in \mathbb{N}$, $n \in \{0, 1, \dots, N\}$, $\varepsilon \in (0, 1]$ in the notation of Theorem 3.12) implies that there exist $\Psi_{\varepsilon, d, N, y} \in \mathbf{N}$, $y \in (\mathbb{R}^d)^N$, $N, d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, which satisfy that

(I) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$ that $\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}) \in C(\mathbb{R}^{d+1}, \mathbb{R}^d)$,

(II) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|Y_{t,x,y}^{d,N} - (\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x)\| \leq \varepsilon(2\sqrt{d} + (g_n^{d,N}(x, y))^3 + (g_{n+1}^{d,N}(x, y))^3), \quad (420)$$

(III) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$, $y \in (\mathbb{R}^d)^N$ that

$$\|(\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x)\| \leq 6\sqrt{d} + 2((g_n^{d,N}(x, y))^2 + (g_{n+1}^{d,N}(x, y))^2), \quad (421)$$

(IV) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$ that

$$\begin{aligned} & \mathcal{P}(\Psi_{\varepsilon, d, N, y}) \\ & \leq \frac{9}{2} N^6 d^{16} \left[2\mathcal{H}(\Phi_d) + \mathfrak{D}_{\varepsilon, 3} + (30 + 6\mathcal{H}(\Phi_d) + [4 + \mathcal{P}(\Phi_d)]^2)^2 \right]^2, \end{aligned} \quad (422)$$

(V) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$[(\mathbb{R}^d)^N \ni y \mapsto (\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x) \in \mathbb{R}^d] \in C((\mathbb{R}^d)^N, \mathbb{R}^d), \quad (423)$$

and

(VI) it holds for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N\}$, $t \in [0, \frac{nT}{N}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N)$, $z = (z_1, z_2, \dots, z_N) \in (\mathbb{R}^d)^N$ with $\forall k \in \mathbb{N} \cap [0, n]: y_k = z_k$ it holds that

$$(\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x) = (\mathcal{R}_a(\Psi_{\varepsilon, d, N, z}))(t, x). \quad (424)$$

Observe that Jensen's inequality implies that for all $n \in \mathbb{N}$, $p \in [1, \infty)$, $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ it holds that

$$|x_1 + \dots + x_n|^p \leq n^{p-1}(|x_1|^p + \dots + |x_n|^p). \quad (425)$$

Moreover, note that Hölder's inequality shows that for all $N \in \mathbb{N}$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\sum_{k=1}^N \|y_k\| = \sum_{k=1}^N (1\|y_k\|) \leq N^{1/2} \left(\sum_{k=1}^N \|y_k\|^2 \right)^{1/2} = N^{1/2} \|y\|. \quad (426)$$

Combining (425), (II), and (419) therefore ensures that for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \|Y_{t,x,y}^{d,N} - (\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x)\| &\leq 2d^{1/2}\varepsilon(1 + (g_N^{d,N}(x, y))^3) \\ &= 2d^{1/2}\varepsilon \left(1 + \left(\|x\| + \mathfrak{C}T + \max_{m \in \{0, 1, \dots, N\}} \left\| \sum_{k=1}^m y_k \right\| \right)^3 \exp(3\mathfrak{C}T) \right) \\ &\leq 2d^{1/2}\varepsilon \left(1 + 9 \left(\|x\|^3 + c^3 + \left(\sum_{k=1}^N \|y_k\| \right)^3 \right) c^3 \right) \\ &\leq 2d^{1/2}\varepsilon (1 + 9(\|x\|^3 + c^3 + N^{3/2}\|y\|^3)c^3) \\ &\leq 2c^6 d^{1/2} N^{3/2} \varepsilon (1 + 9(\|x\|^3 + 1 + \|y\|^3)) \\ &= 2c^6 d^{1/2} N^{3/2} \varepsilon (10 + 9\|x\|^3 + 9\|y\|^3) \\ &\leq 20c^6 d^{1/2} N^{3/2} \varepsilon (1 + \|x\|^3 + \|y\|^3). \end{aligned} \quad (427)$$

Next note that (III), (419), (425), and (426) imply that for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $n \in \{0, 1, \dots, N-1\}$, $t \in [\frac{nT}{N}, \frac{(n+1)T}{N}]$, $x \in \mathbb{R}^d$, $y = (y_1, y_2, \dots, y_N) \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \|(\mathcal{R}_a(\Psi_{\varepsilon, d, N, y}))(t, x)\| &\leq 6\sqrt{d} + 4(g_N^{d,N}(x, y))^2 \\ &= 6\sqrt{d} + 4 \left(\|x\| + \mathfrak{C}T + \max_{m \in \{0, 1, \dots, N\}} \left\| \sum_{k=1}^m y_k \right\| \right)^2 \exp(2\mathfrak{C}T) \\ &\leq 6\sqrt{d} + 12 \left(\|x\|^2 + c^2 + \left(\sum_{k=1}^N \|y_k\| \right)^2 \right) c^2 \\ &\leq 6\sqrt{d} + 12(\|x\|^2 + c^2 + N\|y\|^2)c^2 \\ &\leq 18c^4 \sqrt{d} N (1 + \|x\|^2 + \|y\|^2). \end{aligned} \quad (428)$$

Furthermore, observe that (418) shows that for all $\varepsilon \in (0, 1]$ it holds that

$$\begin{aligned} (\mathfrak{D}_{\varepsilon,3})^2 &= \left(\frac{2160}{\ln(2)}\ln(\varepsilon^{-1}) + \mathfrak{D}_{1,3}\right)^2 \leq (\mathfrak{D}_{1,3})^2(\ln(\varepsilon^{-1}) + 1)^2 \\ &\leq c^2|1 - \ln(\varepsilon)|^2 \leq 2c^2(1 + |\ln(\varepsilon)|^2). \end{aligned} \quad (429)$$

This, (IV), the hypothesis that for all $d \in \mathbb{N}$ it holds that $\mathcal{P}(\Phi_d) \leq \mathfrak{C}d^\mathfrak{D}$, and (425) assure that for all $\varepsilon \in (0, 1]$, $d, N \in \mathbb{N}$, $y \in (\mathbb{R}^d)^N$ it holds that

$$\begin{aligned} \mathcal{P}(\Psi_{\varepsilon,d,N,y}) &\leq \frac{9}{2}N^6d^{16} \left[2\mathfrak{C}d^\mathfrak{D} + \mathfrak{D}_{\varepsilon,3} + (30 + 6\mathfrak{C}d^\mathfrak{D} + [4 + \mathfrak{C}d^\mathfrak{D}]^2)^2\right]^2 \\ &\leq \frac{27}{2}N^6d^{16} \left[4\mathfrak{C}^2d^{2\mathfrak{D}} + (\mathfrak{D}_{\varepsilon,3})^2 + (30 + 6\mathfrak{C}d^\mathfrak{D} + 2[16 + \mathfrak{C}^2d^{2\mathfrak{D}}])^4\right] \\ &\leq \frac{27}{2}N^6d^{16} \left[4\mathfrak{C}^2d^{2\mathfrak{D}} + 2c^2(1 + |\ln(\varepsilon)|^2) + (62 + 6\mathfrak{C}d^\mathfrak{D} + 2\mathfrak{C}^2d^{2\mathfrak{D}})^4\right] \\ &\leq \frac{27}{2}N^6d^{16} \left[4\mathfrak{C}^2d^{2\mathfrak{D}} + 2c^2(1 + |\ln(\varepsilon)|^2) + (62 + 6\mathfrak{C}(\mathfrak{C} + 1)d^{2\mathfrak{D}})^4\right] \quad (430) \\ &\leq \frac{27}{2}N^6d^{16} \left[cd^{2\mathfrak{D}} + 2c^2(1 + |\ln(\varepsilon)|^2) + (cd^{2\mathfrak{D}})^4\right] \\ &\leq 27N^6d^{16} \left[c^2(1 + |\ln(\varepsilon)|^2) + (cd^{2\mathfrak{D}})^4\right] \\ &\leq 54c^4N^6d^{16+8\mathfrak{D}} \left[1 + |\ln(\varepsilon)|^2\right]. \end{aligned}$$

Combining (I), (427), (428), (430), (V), and (VI) establishes items (i)-(vi). The proof of Corollary 3.13 is thus completed. \square

References

- [1] BECK, C., BECKER, S., CHERIDITO, P., JENTZEN, A., AND NEUFELD, A. Deep splitting method for parabolic PDEs. *arXiv:1907.03452* (2019), 40 pages.
- [2] BECK, C., BECKER, S., GROHS, P., JAAFARI, N., AND JENTZEN, A. Solving stochastic differential equations and Kolmogorov equations by means of deep learning. *arXiv:1806.00421* (2018), 56 pages.
- [3] BECK, C., E, W., AND JENTZEN, A. Machine Learning Approximation Algorithms for High-Dimensional Fully Nonlinear Partial Differential Equations and Second-order Backward Stochastic Differential Equations. *Journal of Nonlinear Science* (Jan 2019). doi: 10.1007/s00332-018-9525-3.
- [4] BECKER, S., CHERIDITO, P., AND JENTZEN, A. Deep Optimal Stopping. *Journal of Machine Learning Research* 20, 74 (2019), 1–25. doi: 10.3929/ethz-b-000344707.

- [5] BECKER, S., CHERIDITO, P., JENTZEN, A., AND WELTI, T. Solving high-dimensional optimal stopping problems using deep learning. *arXiv:1908.01602* (2019), 42 pages.
- [6] BERG, J., AND NYSTRÖM, K. A unified deep artificial neural network approach to partial differential equations in complex geometries. *Neurocomputing* 317 (2018), 28–41. doi:10.1016/j.neucom.2018.06.056.
- [7] BERNER, J., GROHS, P., AND JENTZEN, A. Analysis of the generalization error: Empirical risk minimization over deep artificial neural networks overcomes the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. *arXiv:1809.03062* (2018), 35 pages.
- [8] CHAN-WAI-NAM, Q., MIKAEL, J., AND WARIN, X. Machine Learning for Semi Linear PDEs. *Journal of Scientific Computing* 79, 3 (Jun 2019), 1667–1712. doi: 10.1007/s10915-019-00908-3.
- [9] E, W., HAN, J., AND JENTZEN, A. Deep Learning-Based Numerical Methods for High-Dimensional Parabolic Partial Differential Equations and Backward Stochastic Differential Equations. *Commun. Math. Stat.* 5, 4 (2017), 349–380. doi: 10.1007/s40304-017-0117-6.
- [10] E, W., AND YU, B. The deep Ritz method: A deep learning-based numerical algorithm for solving variational problems. *arXiv:1710.00211* (2017), 14 pages.
- [11] ELBRÄCHTER, D., GROHS, P., JENTZEN, A., AND SCHWAB, C. DNN Expression Rate Analysis of High-dimensional PDEs: Application to Option Pricing. *arXiv:1809.07669v2* (2018), 50 pages.
- [12] FARAHMAND, A.-M., NABI, S., AND NIKOVSKI, D. N. Deep reinforcement learning for partial differential equation control. *American Control Conference (ACC)* (2017), 3120–3127. doi: 10.23919/ACC.2017.7963427.
- [13] FUJII, M., TAKAHASHI, A., AND TAKAHASHI, M. Asymptotic Expansion as Prior Knowledge in Deep Learning Method for High dimensional BSDEs. *Asia-Pacific Financial Markets* (Mar 2019). doi: 10.1007/s10690-019-09271-7.
- [14] GOODFELLOW, I., BENGIO, Y., AND COURVILLE, A. *Deep Learning*. MIT Press, 2016. <http://www.deeplearningbook.org>.
- [15] GOUDENEGE, L., MOLENT, A., AND ZANETTE, A. Machine Learning for Pricing American Options in High Dimension. *arXiv:1903.11275* (2019), 11 pages.

- [16] GROHS, P., HORNING, F., JENTZEN, A., AND VON WURSTEMBERGER, P. A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. *arXiv:1809.02362* (2018), 124 pages.
- [17] GROHS, P., PEREKRESTENKO, D., ELBRÄCHTER, D., AND BÖLCSKEI, H. Deep Neural Network Approximation Theory. *arXiv:1901.02220* (2019), 60 pages.
- [18] HAN, J., JENTZEN, A., AND E, W. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences* 115, 34 (2018), 8505–8510. doi: 10.1073/pnas.1718942115.
- [19] HAN, J., AND LONG, J. Convergence of the deep BSDE method for coupled FBSDEs. *arXiv:1811.01165* (2018), 26 pages.
- [20] HENRY-LABORDERE, P. Deep Primal-Dual Algorithm for BSDEs: Applications of Machine Learning to CVA and IM. *Preprint, SSRN-id3071506* (2017), 16 pages.
- [21] HURÉ, C., PHAM, H., AND WARIN, X. Some machine learning schemes for high-dimensional nonlinear PDEs. *arXiv:1902.01599* (2019), 33 pages.
- [22] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., AND NGUYEN, T. A. A proof that rectified deep neural networks overcome the curse of dimensionality in the numerical approximation of semilinear heat equations. *arXiv:1901.10854* (2019), 24 pages.
- [23] HUTZENTHALER, M., JENTZEN, A., AND VON WURSTEMBERGER, P. Overcoming the curse of dimensionality in the approximative pricing of financial derivatives with default risks. *arXiv:1903.05985* (2019), 71 pages.
- [24] JACQUIER, A., AND OUMGARI, M. Deep PPDEs for rough local stochastic volatility. *arXiv:1906.02551* (2019), 21 pages.
- [25] JENTZEN, A., SALIMOVA, D., AND WELTI, T. A proof that deep artificial neural networks overcome the curse of dimensionality in the numerical approximation of Kolmogorov partial differential equations with constant diffusion and nonlinear drift coefficients. *arXiv:1809.07321* (2018), 48 pages.
- [26] KUTYNIOK, G., PETERSEN, P., RASLAN, M., AND SCHNEIDER, R. A theoretical analysis of deep neural networks and parametric PDEs. *arXiv:1904.00377* (2019), 42 pages.

- [27] LI, B., SHANSHAN, T., AND YU, H. Better Approximations of High Dimensional Smooth Functions by Deep Neural Networks with Rectified Power Units. *arXiv:1903.05858v3* (2019), 21 pages.
- [28] LONG, Z., LU, Y., MA, X., AND DONG, B. PDE-Net: Learning PDEs from Data. *arXiv:1710.09668* (2017), 15 pages.
- [29] LYE, K. O., MISHRA, S., AND RAY, D. Deep learning observables in computational fluid dynamics. *arXiv:1903.03040* (2019), 57 pages.
- [30] MAGILL, M., QURESHI, F., AND DE HAAN, H. Neural Networks Trained to Solve Differential Equations Learn General Representations. *Advances in Neural Information Processing Systems* (2018), 4075–4085.
- [31] PETERSEN, P., RASLAN, M., AND VOIGTLAENDER, F. Topological properties of the set of functions generated by neural networks of fixed size. *arXiv:1806.08459v2* (2018), 56 pages.
- [32] PETERSEN, P., AND VOIGTLAENDER, F. Optimal approximation of piecewise smooth functions using deep ReLU neural networks. *arXiv:1709.05289* (2017), 54 pages.
- [33] PHAM, H., AND WARIN, X. Neural networks-based backward scheme for fully nonlinear PDEs. *arXiv:1908.00412v1* (2019), 15 pages.
- [34] RAISSI, M. Deep Hidden Physics Models: Deep Learning of Nonlinear Partial Differential Equations. *arXiv:1801.06637* (2018), 26 pages.
- [35] REISINGER, C., AND ZHANG, Y. Rectified deep neural networks overcome the curse of dimensionality for nonsmooth value functions in zero-sum games of nonlinear stiff systems. *arXiv:1903.06652* (2019), 34 pages.
- [36] SIRIGNANO, J., AND SPILIOPOULOS, K. DGM: A deep learning algorithm for solving partial differential equations. *Journal of Computational Physics* 375 (2018), 1339–1364. doi: 10.1016/j.jcp.2018.08.029.
- [37] YAROTSKY, D. Error bounds for approximations with deep ReLU networks. *Neural Networks* 94 (2017), 103–114. doi: 10.1016/j.neunet.2017.07.002.