

MLP starting ideas

July 15, 2022

Abstract

Abstract goes here...

Contents

1	Introduction	1
2	Multilevel Picard approximations for the heat equation	1
3	Stochastic solutions to parabolic partial differential equations	2

1 Introduction

Add an appropriate introduction...

2

Theorem 2.1. *Let $T, \kappa, \delta \in (0, \infty)$, $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$, let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $t \in [0, T]$, $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ that*

$$|u_d(t, x)| \leq \kappa d^\kappa \left(1 + \sum_{k=1}^d |x_k|\right)^\kappa \quad \text{and} \quad \left(\frac{\partial}{\partial t} u_d\right)(t, x) = (\Delta_x u_d)(t, x), \quad (2.1)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W^{d,\theta}: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, $\theta \in \Theta$, be independent standard Brownian motions, let $U_m^{d,\theta}: [0, T] \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$, $d, m \in \mathbb{Z}$, $\theta \in \Theta$, satisfy for all $d, m \in \mathbb{N}$, $\theta \in \Theta$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$U_m^{d,\theta}(t, x) = \frac{1}{m} \left[\sum_{k=1}^m u_d(0, x + \sqrt{2} W_t^{d,(\theta,0,-k)}) \right],$$

and for every $d, n, m \in \mathbb{N}$ let $\mathfrak{C}_{d,n,m} \in \mathbb{N}$ be the number of function evaluations of $u_d(0, \cdot)$ and the number of realizations of scalar random variables which are used to compute one realization of $U_m^{d,0}(T, 0): \Omega \rightarrow \mathbb{R}$. Then there exist $c \in \mathbb{R}$ and $\mathfrak{n}: \mathbb{N} \times (0, 1] \rightarrow \mathbb{N}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\left(\mathbb{E} \left[|u_d(T, 0) - U_{\mathfrak{n}(d,\varepsilon)}^{d,0}(T, 0)|^2 \right] \right)^{1/2} \leq \varepsilon \quad \text{and} \quad \mathfrak{C}_{d,\mathfrak{n}(d,\varepsilon),\mathfrak{n}(d,\varepsilon)} \leq cd^c \varepsilon^{-(2+\delta)}. \quad (2.2)$$

3 Stochastic solutions to parabolic partial differential equations

Lemma 3.1. *Let $T \in (0, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that*

$$\left(\frac{\partial}{\partial t} u_d\right)(t, x) + (\Delta_x u_d)(t, x) = 0, \quad (3.1)$$

let $W^d: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, be standard Brownian motions, and let $\mathcal{X}^{d,t,x}: [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, be a stochastic process with continuous sample paths satisfying that for all $d \in \mathbb{N}$, $t \in [0, T]$, $s \in [t, T]$, $x \in \mathbb{R}^d$ we have \mathbb{P} -a.s. that

$$\mathcal{X}_s^{d,t,x} = x + \int_t^s \sqrt{2} dW_r^d = x + \sqrt{2} W_{t-s}^d. \quad (3.2)$$

Then for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$u_d(t, x) = \mathbb{E}\left[u_d(T, \mathcal{X}_T^{d,t,x})\right]. \quad (3.3)$$

Proof of Lemma 3.1. The proof of Lemma 3.1 is thus complete. \square

Lemma 3.2. *Let $T \in (0, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\sigma_d: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$, $d \in \mathbb{N}$, be infinitely often differentiable functions, let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that*

$$\left(\frac{\partial}{\partial t} u_d\right)(t, x) + \text{Trace}\left(\sigma(x)[\sigma(x)]^*(\text{Hess}_x u_d)(t, x)\right) = 0, \quad (3.4)$$

let $W^d: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, be standard Brownian motions, and let $\mathcal{X}^{d,t,x}: [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, be a stochastic process with continuous sample paths satisfying that for all $d \in \mathbb{N}$, $t \in [0, T]$, $s \in [t, T]$, $x \in \mathbb{R}^d$ we have \mathbb{P} -a.s. that

$$\mathcal{X}_s^{d,t,x} = x + \int_s^t \sqrt{2} \sigma(\mathcal{X}_r^{d,t,x}) dW_r^d. \quad (3.5)$$

Then for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$u_d(t, x) = \mathbb{E}\left[u_d(T, \mathcal{X}_T^{d,t,x})\right]. \quad (3.6)$$

Proof of Lemma 3.2. The proof of Lemma 3.2 is thus complete. \square

Lemma 3.3. *Let $T \in (0, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\mu_d \in \mathbb{R}^d \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, be infinitely often differentiable functions, let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that*

$$\left(\frac{\partial}{\partial t} u_d\right)(t, x) + (\Delta_x u_d)(t, x) + [\mu_d(x)]^*(\nabla_x u_d)(t, x) = 0, \quad (3.7)$$

let $W^d: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, be standard Brownian motions, and let $\mathcal{X}^{d,t,x}: [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, be a stochastic process with continuous sample paths satisfying that for all $d \in \mathbb{N}$, $t \in [0, T]$, $s \in [t, T]$, $x \in \mathbb{R}^d$ we have \mathbb{P} -a.s. that

$$\mathcal{X}_s^{d,t,x} = x + \int_s^t \mu_d(\mathcal{X}_r^{d,t,x}) dr + \int_s^t \sqrt{2} dW_r^d. \quad (3.8)$$

Then for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$u_d(t, x) = \mathbb{E}\left[u_d(T, \mathcal{X}_T^{d,t,x})\right]. \quad (3.9)$$

Proof of Lemma 3.3. The proof of Lemma 3.3 is thus complete. \square

Lemma 3.4. Let $T \in (0, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $\alpha_d \in \mathbb{R}^d \rightarrow \mathbb{R}$, $d \in \mathbb{N}$, be infinitely often differentiable functions, let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$, $d \in \mathbb{N}$, satisfy for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\left(\frac{\partial}{\partial t}u_d\right)(t, x) + (\Delta_x u_d)(t, x) + \alpha_d(x)u_d(t, x) = 0, \quad (3.10)$$

let $W^d: [0, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, be standard Brownian motions, and let $\mathcal{X}^{d,t,x}: [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, be a stochastic process with continuous sample paths satisfying that for all $d \in \mathbb{N}$, $t \in [0, T]$, $s \in [t, T]$, $x \in \mathbb{R}^d$ we have \mathbb{P} -a.s. that

$$\mathcal{X}_s^{d,t,x} = x + \int_s^t \sqrt{2} dW_r^d. \quad (3.11)$$

Then for all $d \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$u_d(t, x) = \mathbb{E}\left[\exp\left(\int_t^T \alpha_d(\mathcal{X}_r^{d,t,x}) dr\right)u_d(T, \mathcal{X}_T^{d,t,x})\right]. \quad (3.12)$$

Proof of Lemma 3.4. The proof of Lemma 3.4 is thus complete. \square